

PART-B

Name of the Student: V. Hemant Kumar Reddy

Signature of the Student: Hemant Roll No.: 230123070

Signature of the Invigilator: [Signature]

INSTRUCTIONS

1. Check that you have **10 printed pages**. The PART-B of the examination has **4 questions**, for a total of **20 points**.
2. Attempt **all questions**.
3. There is **no credit** for a solution if the appropriate work is not shown, even if the answer is correct.
4. Write your answers in this booklet and only in the **space marked for answers**. Any answer written in any space other than the designated space will **not be evaluated**.
5. The pages numbered **8 to 10** are kept as **additional pages**. If you cannot able to fit a complete answer in the space provided just after the question, you can use these three pages by referring the page number of additional page in the original space.
6. At the beginning of the examination, a supplementary sheet has been provided only for rough work. Should you need more, please ask the invigilator.
7. No questions require any clarification from the instructor or invigilators, even if a question has an error or incomplete data. The students are advised to write answer according to their understanding or write reasons for why it is not possible to solve partially or completely that question by citing errors/insufficient data.

FOR OFFICE USE ONLY

Question:	1	2	3	4	Total
Points:	6	7	5	2	20
Score:	0	7	4	1	12

QUESTIONS

1. (6 points) Let

$$S = \begin{pmatrix} S_{yy} & S_{yx} \\ S_{yx} & S_{xx} \end{pmatrix}$$

be the partitioned sample covariance matrix of $Y_{1 \times 1}$ and $X_{q \times 1}$ based on a centered (with respect to sample mean) sample of size n . Prove that the squared sample first canonical correlation r_1^2 is identical to the coefficient of determination (R^2) obtained from the multiple linear regression of Y on the vector X .

Sol.
p.

We know that

Sample first canonical correlation is given by ρ_1^*

where $\rho_1^{*2}, \rho_2^{*2}, \rho_3^{*2}, \dots$ are eigen values of the

matrix $(S_{xx}^{-1/2}) S_{yx} (S_{yy}^{-1}) (S_{yx}^T) (S_{xx}^{-1/2})$

and eigen vectors $e_1, e_2, \dots$ respectively.

$$\Rightarrow \frac{(S_{xx}^{-1/2}) S_{yx} S_{yy}^{-1} S_{yx}^T (S_{xx}^{-1/2})}{(S_{yy})} \cdot e_1 = (\rho_1^*)^2 \cdot e_1$$

$$\Rightarrow (\rho_1^*)^2 = \frac{e_1^T (S_{xx}^{-1/2} \cdot S_{yx} \cdot S_{yy}^{-1} \cdot S_{yx}^T \cdot S_{xx}^{-1/2}) e_1}{S_{yy}} = r_1^2$$

We have maximum co-variance / co-variation

$$\Downarrow$$

$$[U = e_1^T (S_{xx}^{-1}) \cdot X]$$

Coefficient of determination $R^2 = \frac{1}{S_{yy}} (S_{xx}^{-1/2} \cdot S_{yx} \cdot S_{yy}^{-1} \cdot S_{yx}^T \cdot S_{xx}^{-1/2})$

Hence proved:

normal, $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ is a basis for $\mathbb{R}^3$ and $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ is a basis for $\mathbb{R}^3$.
 Find the matrix A such that $A\mathbf{e}_1 = \mathbf{e}_2, A\mathbf{e}_2 = \mathbf{e}_3, A\mathbf{e}_3 = \mathbf{e}_1$.

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

$$[A\mathbf{e}_1, A\mathbf{e}_2, A\mathbf{e}_3] = [\mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1]$$

$$[A\mathbf{e}_1, A\mathbf{e}_2, A\mathbf{e}_3] = [\mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1]$$

$$\begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad (3)$$

$$[A\mathbf{e}_1, A\mathbf{e}_2, A\mathbf{e}_3] = [\mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1] \text{ we have}$$

$$A\mathbf{e}_1 = \mathbf{e}_2, A\mathbf{e}_2 = \mathbf{e}_3, A\mathbf{e}_3 = \mathbf{e}_1$$

$$A\mathbf{e}_1 = \mathbf{e}_2, A\mathbf{e}_2 = \mathbf{e}_3, A\mathbf{e}_3 = \mathbf{e}_1$$

$$A\mathbf{e}_1 = \mathbf{e}_2, A\mathbf{e}_2 = \mathbf{e}_3, A\mathbf{e}_3 = \mathbf{e}_1 \Rightarrow A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

$$A\mathbf{e}_1 = \mathbf{e}_2, A\mathbf{e}_2 = \mathbf{e}_3, A\mathbf{e}_3 = \mathbf{e}_1$$

2. Suppose you are performing a factor analysis on a 4×4 sample covariance matrix $\mathbf{S}$. The eigenvalues of $\mathbf{S}$ are $\lambda_1 = 0.8$, $\lambda_2 = 2.1$, $\lambda_3 = 4.2$, and $\lambda_4 = 0.2$. The corresponding normalized eigenvectors are $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$, and $\mathbf{e}_4$. You decide to extract $m = 2$ factors using the principal component method.
- (2 points) Write down the explicit mathematical expression for the estimated factor loading matrix $\tilde{\mathbf{L}}$ in terms of the given eigenvalues and eigenvectors.
 - (1 point) Let the first row of $\tilde{\mathbf{L}}$ be $\tilde{\mathbf{l}}_1^T = [0.8, -0.4]$. Calculate the estimated communality $\tilde{h}_1^2$ for the first variable.
 - (4 points) Prove that the sum of all estimated communalities $\sum_{i=1}^p \tilde{h}_i^2$ is exactly equal to the sum of the largest $m = 2$ eigenvalues of $\mathbf{S}$.

Sol. eigen values of $\mathbf{S}$ are $\lambda_1 = 0.8, \lambda_2 = 2.1, \lambda_3 = 4.2, \lambda_4 = 0.2$
 $(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_4) \Rightarrow$ corresponding normalized eigen vectors.

(a)

$$\lambda_3 > \lambda_2 > \lambda_1 > \lambda_4$$

hence

$$\tilde{\mathbf{L}} = [\sqrt{\lambda_3} \mathbf{e}_3 \mid \sqrt{\lambda_2} \mathbf{e}_2]$$

02

$$= [\sqrt{4.2} \mathbf{e}_3 \mid \sqrt{2.1} \mathbf{e}_2]$$

(b)

$$\text{let } \mathbf{e}_3 = \begin{pmatrix} e_{31} \\ e_{32} \\ e_{33} \\ e_{34} \end{pmatrix}, \quad \mathbf{e}_2 = \begin{pmatrix} e_{21} \\ e_{22} \\ e_{23} \\ e_{24} \end{pmatrix}$$

$$\text{first row } [\sqrt{4.2} e_{31}, \sqrt{2.1} e_{21}] = [0.8, -0.4]$$

$$\Rightarrow e_{31} = \frac{0.8}{\sqrt{4.2}} = \frac{0.8}{2.05} = \frac{8}{20.5} = \frac{8}{20.5}$$

$$\Rightarrow e_{21} = \frac{-0.4}{\sqrt{2.1}} = \frac{-0.4}{1.45} = \frac{-4}{14.5} = \frac{-4}{14.5}$$

01

$$\tilde{h}_1^2 = (\sqrt{4.2} e_{31})^2 + (\sqrt{2.1} e_{21})^2 = (0.8)^2 + (-0.4)^2 = 0.8$$

$$= \left(\frac{8}{20.5} \right)^2 \times 4.2 + \left(\frac{-4}{14.5} \right)^2 \times 2.1 = 0.8$$

(c) $\sum_{i=1}^R \tilde{h}_i^2$

$$h_1^2 = \lambda_3 e_{31}^2 + \lambda_2 e_{21}^2$$

$$h_2^2 = \lambda_3 e_{32}^2 + \lambda_2 e_{22}^2$$

$$h_3^2 = \lambda_3 e_{33}^2 + \lambda_2 e_{23}^2$$

$$h_4^2 = \lambda_3 e_{34}^2 + \lambda_2 e_{24}^2$$

$$\sum_{i=1}^R \tilde{h}_i^2 = \lambda_3 \left(\sum_{i=1}^4 e_{3i}^2 \right) + \lambda_2 \left(\sum_{i=1}^4 e_{2i}^2 \right)$$

w.k.t ' $e_3, e_2 \Rightarrow$ normalised eigen vectors

$$e_3^T e_3 = 1 \Rightarrow \sum_{i=1}^4 e_{3i}^2 = 1$$

04

Similarly

$$\sum_{i=1}^4 e_{2i}^2 = 1$$

Hence,

$$\sum_{i=1}^R \tilde{h}_i^2 =$$

$$\lambda_3 + \lambda_2$$

Sum of largest two eigen values

3. (5 points) Let $X_1, X_2, \dots, X_n$ be a random sample of size n from a distribution with mean $\mu \in \mathbb{R}$ and variance μ^2 . Let $\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$. Derive a variance-stabilizing transformation $g(\bar{X}_n)$ such that the asymptotic variance of $g(\bar{X}_n)$ does not depend on μ .

Sol.
p.

$$X_i \sim N(\mu, \mu^2), \mu \in \mathbb{R}, \forall i \in \{1, 2, \dots, n\}$$

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i$$

$$E(\bar{X}_n) = \frac{1}{n} \sum_{i=1}^n E(X_i) = \frac{1}{n} \cdot \mu n = \mu$$

$$\text{Var}(\bar{X}_n) = \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i) = \left(\frac{\mu^2 \cdot n}{n^2} \right) = \frac{\mu^2}{n} \quad \left(\begin{array}{l} \text{since random sample} \\ \text{Cov}(X_i, X_j) = 0 \quad \forall i \neq j \end{array} \right)$$

$$\bar{X}_n \sim N\left(\mu, \frac{\mu^2}{n}\right)$$

$$\sqrt{n}(\bar{X}_n - \mu) \sim N(0, \mu^2)$$

From Taylor's expansion

$$g(\bar{X}_n) \approx g(\mu) + (\bar{X}_n - \mu) \cdot g'(\mu)$$

$$(g(\bar{X}_n) - g(\mu)) \approx (\bar{X}_n - \mu) g'(\mu)$$

$$\text{as } \sqrt{n}(\bar{X}_n - \mu) \xrightarrow{d} N(0, \mu^2), \text{ so } \sqrt{n}(g(\bar{X}_n) - g(\mu)) \xrightarrow{d} N(0, \mu^2 [g'(\mu)]^2)$$


$$\mu^2 [g'(\mu)]^2 = \text{Constant} \Rightarrow g'(\mu) \mu = \sqrt{k}$$

$$\Rightarrow g(\mu) = \sqrt{k} \ln \mu \Rightarrow \boxed{g(x) = \sqrt{k} \ln x}$$

$$\boxed{g(\bar{X}_n) = \sqrt{k} (\ln \bar{X}_n)}$$

where k is some constant.

4. (2 points) The figure below displays a Normal Q-Q plot for a dataset of size $n = 50$. Interpret this plot to determine if the data follows a normal distribution. If the normality assumption is violated, specify which class of distributions would more accurately characterize the data.



- * From the data it is clear that points at the extreme moving away largely from the line.
- * Hence, the data follows log-log distribution → What is it??
- * Hence, normality assumption is violated.

01

ADDITIONAL PAGES FOR ANSWERS

