

MA 101 (Mathematics-I)

Subhamay Saha and Ayon Ganguly
Department of Mathematics
IIT Guwahati

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ be such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$.

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ be such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$.

If $f : D \rightarrow \mathbb{R}$, then $\ell \in \mathbb{R}$ is said to be the limit of f at x_0 if for each $\varepsilon > 0$, there exists $\delta > 0$ such that

whenever $x \in D$ and $0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$.

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ be such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$.

If $f : D \rightarrow \mathbb{R}$, then $\ell \in \mathbb{R}$ is said to be the limit of f at x_0 if for each $\varepsilon > 0$, there exists $\delta > 0$ such that

whenever $x \in D$ and $0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$.

We write: $\lim_{x \rightarrow x_0} f(x) = \ell$.

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ be such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$.

If $f : D \rightarrow \mathbb{R}$, then $\ell \in \mathbb{R}$ is said to be the limit of f at x_0 if for each $\varepsilon > 0$, there exists $\delta > 0$ such that

$$\text{whenever } x \in D \text{ and } 0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon.$$

We write: $\lim_{x \rightarrow x_0} f(x) = \ell$.

Result: If limit exists, then it is unique.

Example: $\lim_{x \rightarrow 1} \left(\frac{3x}{2} - 1 \right) = \frac{1}{2}$. Let $\varepsilon > 0$. We have to find $\delta > 0$ such that $0 < |x - 1| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ holds with $\ell = 1/2$. Working backwards,

$$\frac{3}{2}|x - 1| < \varepsilon \text{ whenever } |x - 1| < \delta := \frac{2}{3}\varepsilon.$$

Example: $\lim_{x \rightarrow 1} \left(\frac{3x}{2} - 1 \right) = \frac{1}{2}$. Let $\varepsilon > 0$. We have to find $\delta > 0$ such that $0 < |x - 1| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ holds with $\ell = 1/2$. Working backwards,

$$\frac{3}{2}|x - 1| < \varepsilon \text{ whenever } |x - 1| < \delta := \frac{2}{3}\varepsilon.$$


Theorem (Sequential criterion)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$. Let $f : D \rightarrow \mathbb{R}$. Then the following are equivalent.


- (a) $\lim_{x \rightarrow x_0} f(x) = \ell$.
- (b) For any sequence (x_n) in D with $x_n \neq x_0$ for all $n \geq 1$ and $x_n \rightarrow x_0$, the sequence $(f(x_n))$ converges to ℓ .

Example: $\lim_{x \rightarrow 0} \sin \frac{1}{x}$ does not exist.

Example: $\lim_{x \rightarrow 0} \sin \frac{1}{x}$ does not exist.




Example: $\lim_{x \rightarrow 0} \sin \frac{1}{x}$ does not exist.



Solution: Let $x_n = \frac{2}{(4n+1)\pi}$ and $y_n = \frac{1}{n\pi}$ for all $n \in \mathbb{N}$. Then $x_n \rightarrow 0$ and $y_n \rightarrow 0$.


Example: $\lim_{x \rightarrow 0} \sin \frac{1}{x}$ does not exist.



Solution: Let $x_n = \frac{2}{(4n+1)\pi}$ and $y_n = \frac{1}{n\pi}$ for all $n \in \mathbb{N}$. Then $x_n \rightarrow 0$ and $y_n \rightarrow 0$.

Since $\sin \frac{1}{x_n} = 1$ and $\sin \frac{1}{y_n} = 0$ for all $n \in \mathbb{N}$, we get $\sin \frac{1}{x_n} \rightarrow 1$ and $\sin \frac{1}{y_n} \rightarrow 0$.

Example: $\lim_{x \rightarrow 0} \sin \frac{1}{x}$ does not exist.



Solution: Let $x_n = \frac{2}{(4n+1)\pi}$ and $y_n = \frac{1}{n\pi}$ for all $n \in \mathbb{N}$. Then $x_n \rightarrow 0$ and $y_n \rightarrow 0$.

Since $\sin \frac{1}{x_n} = 1$ and $\sin \frac{1}{y_n} = 0$ for all $n \in \mathbb{N}$, we get $\sin \frac{1}{x_n} \rightarrow 1$ and $\sin \frac{1}{y_n} \rightarrow 0$.

Therefore by the sequential criterion for limit, $\lim_{x \rightarrow 0} \sin \frac{1}{x}$ does not exist.

A function $f : D \rightarrow \mathbb{R}$ is called bounded if there exists $M > 0$ such that $|f(x)| < M$ for all $x \in D$.

A function $f : D \rightarrow \mathbb{R}$ is called bounded if there exists $M > 0$ such that $|f(x)| < M$ for all $x \in D$.

Result: Let $f : D \rightarrow \mathbb{R}$. Suppose that $\lim_{x \rightarrow x_0} f(x) = \ell$. Then there exists some $\delta > 0$ such that f is bounded on $(x_0 - \delta, x_0 + \delta) \setminus \{x_0\}$. That is, there exists $M > 0$ such that $|f(x)| < M$ for all $x \in (x_0 - \delta, x_0 + \delta)$ with $x \neq x_0$.

Theorem (Limit Theorems)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$. Let $f, g, j : D \rightarrow \mathbb{R}$. Suppose that $\lim_{x \rightarrow x_0} f(x) = \ell$ and $\lim_{x \rightarrow x_0} g(x) = m$. Then

Theorem (Limit Theorems)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$. Let $f, g, j : D \rightarrow \mathbb{R}$. Suppose that $\lim_{x \rightarrow x_0} f(x) = \ell$ and $\lim_{x \rightarrow x_0} g(x) = m$. Then

(1) $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$.

Theorem (Limit Theorems)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$. Let $f, g, j : D \rightarrow \mathbb{R}$. Suppose that $\lim_{x \rightarrow x_0} f(x) = \ell$ and $\lim_{x \rightarrow x_0} g(x) = m$. Then

- (1) $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$.
- (2) If $f(x) \leq g(x)$ for all $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$, then $\ell \leq m$.

Theorem (Limit Theorems)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$. Let $f, g, j : D \rightarrow \mathbb{R}$. Suppose that $\lim_{x \rightarrow x_0} f(x) = \ell$ and $\lim_{x \rightarrow x_0} g(x) = m$. Then

- (1) $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$.
- (2) If $f(x) \leq g(x)$ for all $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$, then $\ell \leq m$.
- (3) $\lim_{x \rightarrow x_0} (fg)(x) = \ell m$ and if $m \neq 0$ and $g(x) \neq 0$ for all $x \in D$, then $\lim_{x \rightarrow x_0} \frac{1}{g(x)} = \frac{1}{m}$.

Theorem (Limit Theorems)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$. Let $f, g, j : D \rightarrow \mathbb{R}$. Suppose that $\lim_{x \rightarrow x_0} f(x) = \ell$ and $\lim_{x \rightarrow x_0} g(x) = m$. Then

- (1) $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$.
- (2) If $f(x) \leq g(x)$ for all $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$, then $\ell \leq m$.
- (3) $\lim_{x \rightarrow x_0} (fg)(x) = \ell m$ and if $m \neq 0$ and $g(x) \neq 0$ for all $x \in D$, then $\lim_{x \rightarrow x_0} \frac{1}{g(x)} = \frac{1}{m}$.
- (4) If $f(x) \leq j(x) \leq g(x)$ for all $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$ and $\ell = m$, then $\lim_{x \rightarrow x_0} j(x) = \ell$.

Result: Suppose that $f(x)$ is bounded in $(x_0 - h, x_0 + h) \setminus \{x_0\}$ for some $h > 0$ and $\lim_{x \rightarrow x_0} g(x) = 0$.
Then $\lim_{x \rightarrow x_0} f(x)g(x) = 0$.


Result: Suppose that $f(x)$ is bounded in $(x_0 - h, x_0 + h) \setminus \{x_0\}$ for some $h > 0$ and $\lim_{x \rightarrow x_0} g(x) = 0$.

Then $\lim_{x \rightarrow x_0} f(x)g(x) = 0$.

Example: $\lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0$.

Result: Suppose that $f(x)$ is bounded in $(x_0 - h, x_0 + h) \setminus \{x_0\}$ for some $h > 0$ and $\lim_{x \rightarrow x_0} g(x) = 0$. Then $\lim_{x \rightarrow x_0} f(x)g(x) = 0$.

Example: $\lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0.$



One sided limits

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$,
 $(x_0, x_0 + h) \subseteq D$.

One sided limits

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0, x_0 + h) \subseteq D$.

If $f : D \rightarrow \mathbb{R}$, then $\ell \in \mathbb{R}$ is said to be the right hand limit of f at x_0 if for each $\varepsilon > 0$, there exists $\delta > 0$ such that

whenever $x \in D$ and $0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon$.

One sided limits

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0, x_0 + h) \subseteq D$.

If $f : D \rightarrow \mathbb{R}$, then $\ell \in \mathbb{R}$ is said to be the right hand limit of f at x_0 if for each $\varepsilon > 0$, there exists $\delta > 0$ such that

whenever $x \in D$ and $0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon$.

We write: $\lim_{x \rightarrow x_0+} f(x) = \ell$.

One sided limits

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0, x_0 + h) \subseteq D$.

If $f : D \rightarrow \mathbb{R}$, then $\ell \in \mathbb{R}$ is said to be the right hand limit of f at x_0 if for each $\varepsilon > 0$, there exists $\delta > 0$ such that

$$\text{whenever } x \in D \text{ and } 0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon.$$

We write: $\lim_{x \rightarrow x_0+} f(x) = \ell$.

Similarly one defines left hand limit of f at x_0 and is denoted by $\lim_{x \rightarrow x_0-} f(x)$.

One sided limits

Let $D \subseteq \mathbb{R}$ and let $x_0 \in \mathbb{R}$ such that for some $h > 0$, $(x_0, x_0 + h) \subseteq D$.

If $f : D \rightarrow \mathbb{R}$, then $\ell \in \mathbb{R}$ is said to be the right hand limit of f at x_0 if for each $\varepsilon > 0$, there exists $\delta > 0$ such that

whenever $x \in D$ and $0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon$.

We write: $\lim_{x \rightarrow x_0+} f(x) = \ell$.


Similarly one defines left hand limit of f at x_0 and is denoted by $\lim_{x \rightarrow x_0-} f(x)$.

Result: $\lim_{x \rightarrow x_0} f(x) = \ell \Leftrightarrow \lim_{x \rightarrow x_0+} f(x) = \lim_{x \rightarrow x_0-} f(x) = \ell$.

Example: Show that $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$.

Example: Show that $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$.

We will use Sandwich Thm for limit
(Limit Thm (4))



Area of $\Delta OAP < \text{Area of the circular section } OAPQ < \text{Area of } \Delta OAT$

Limits at infinity and infinite limits

Definition: $f(x)$ has limit ℓ as x approaches $+\infty$, if for any given $\varepsilon > 0$, there exists $M > 0$ such that

$$x > M \implies |f(x) - \ell| < \varepsilon.$$

Similarly, one can define limit of $f(x)$ as x approaches $-\infty$.

Limits at infinity and infinite limits

Definition: $f(x)$ has limit ℓ as x approaches $+\infty$, if for any given $\varepsilon > 0$, there exists $M > 0$ such that

$$x > M \implies |f(x) - \ell| < \varepsilon.$$

Similarly, one can define limit of $f(x)$ as x approaches $-\infty$.

Example: (i) $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, (ii) $\lim_{x \rightarrow -\infty} \frac{1}{x} = 0$,
 (iii) $\lim_{x \rightarrow \infty} \sin x$ does not exist.

Limits at infinity and infinite limits

Definition: $f(x)$ has limit ℓ as x approaches $+\infty$, if for any given $\varepsilon > 0$, there exists $M > 0$ such that

$$x > M \implies |f(x) - \ell| < \varepsilon.$$

Similarly, one can define limit of $f(x)$ as x approaches $-\infty$.

Example: (i) $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$, (ii) $\lim_{x \rightarrow -\infty} \frac{1}{x} = 0$,
(iii) $\lim_{x \rightarrow \infty} \sin x$ does not exist.

Definition: A function $f(x)$ approaches ∞ ($f(x) \rightarrow \infty$) as $x \rightarrow x_0$ if, for every real $M > 0$, there exists $\delta > 0$ such that

$$0 < |x - x_0| < \delta \implies f(x) > M.$$

Similarly, one can define limit of $f(x)$ approaching $-\infty$.

Example: (i) $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$, (ii) $\lim_{x \rightarrow 0} \frac{1}{x^2} \sin(1/x)$ does not exist.

Example: (i) $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$, (ii) $\lim_{x \rightarrow 0} \frac{1}{x^2} \sin(1/x)$ does not exist.

For (ii), let $x_n = \frac{1}{\frac{\pi}{2} + 2n\pi}$ and $y_n = \frac{1}{n\pi}$. Then $x_n, y_n \rightarrow 0$ as $n \rightarrow \infty$.

But $\lim_{n \rightarrow \infty} f(x_n) = \frac{1}{x_n^2} \rightarrow \infty$ and $\lim_{n \rightarrow \infty} f(y_n) = 0$.

Example: (i) $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$, (ii) $\lim_{x \rightarrow 0} \frac{1}{x^2} \sin(1/x)$ does not exist.

For (ii), let $x_n = \frac{1}{\frac{\pi}{2} + 2n\pi}$ and $y_n = \frac{1}{n\pi}$. Then $x_n, y_n \rightarrow 0$ as $n \rightarrow \infty$.

But $\lim_{n \rightarrow \infty} f(x_n) = \frac{1}{x_n^2} \rightarrow \infty$ and $\lim_{n \rightarrow \infty} f(y_n) = 0$.

Theorem

Suppose that $\lim_{x \rightarrow x_0} f(x) = \ell$. If $\ell \neq 0$, then there exists some δ such that $f(x) \neq 0$ for all $x \in (x_0 - \delta, x_0 + \delta) \setminus \{x_0\}$.

Continuous functions

Let D be a nonempty subset of \mathbb{R} and $f : D \rightarrow \mathbb{R}$. We say that f is continuous at $x_0 \in D$ if for each $\varepsilon > 0$, there exists $\delta > 0$ such that $|f(x) - f(x_0)| < \varepsilon$ for all $x \in D$ satisfying $|x - x_0| < \delta$.

Continuous functions

Let D be a nonempty subset of \mathbb{R} and $f : D \rightarrow \mathbb{R}$. We say that f is continuous at $x_0 \in D$ if for each $\varepsilon > 0$, there exists $\delta > 0$ such that $|f(x) - f(x_0)| < \varepsilon$ for all $x \in D$ satisfying $|x - x_0| < \delta$.

Let $f : [a, b] \rightarrow \mathbb{R}$. Then f is continuous at $c \in (a, b)$ if $\lim_{x \rightarrow c} f(x) = f(c)$. f is continuous at a if $\lim_{x \rightarrow a+} f(x) = f(a)$. Similarly, f is continuous at b if $\lim_{x \rightarrow b-} f(x) = f(b)$.

Continuous functions

Let D be a nonempty subset of \mathbb{R} and $f : D \rightarrow \mathbb{R}$. We say that f is continuous at $x_0 \in D$ if for each $\varepsilon > 0$, there exists $\delta > 0$ such that $|f(x) - f(x_0)| < \varepsilon$ for all $x \in D$ satisfying $|x - x_0| < \delta$.

Let $f : [a, b] \rightarrow \mathbb{R}$. Then f is continuous at $c \in (a, b)$ if $\lim_{x \rightarrow c} f(x) = f(c)$. f is continuous at a if $\lim_{x \rightarrow a+} f(x) = f(a)$. Similarly, f is continuous at b if $\lim_{x \rightarrow b-} f(x) = f(b)$.

Sequential criterion of continuity: $f : D \rightarrow \mathbb{R}$ is continuous at $x_0 \in D$ if and only if for every sequence (x_n) in D such that $x_n \rightarrow x_0$, we have $f(x_n) \rightarrow f(x_0)$.

Continuous functions

Let D be a nonempty subset of \mathbb{R} and $f : D \rightarrow \mathbb{R}$. We say that f is continuous at $x_0 \in D$ if for each $\varepsilon > 0$, there exists $\delta > 0$ such that $|f(x) - f(x_0)| < \varepsilon$ for all $x \in D$ satisfying $|x - x_0| < \delta$.

Let $f : [a, b] \rightarrow \mathbb{R}$. Then f is continuous at $c \in (a, b)$ if $\lim_{x \rightarrow c} f(x) = f(c)$. f is continuous at a if $\lim_{x \rightarrow a+} f(x) = f(a)$. Similarly, f is continuous at b if $\lim_{x \rightarrow b-} f(x) = f(b)$.

Sequential criterion of continuity: $f : D \rightarrow \mathbb{R}$ is continuous at $x_0 \in D$ if and only if for every sequence (x_n) in D such that $x_n \rightarrow x_0$, we have $f(x_n) \rightarrow f(x_0)$.

We say that $f : D \rightarrow \mathbb{R}$ is continuous if f is continuous at each $x_0 \in D$.

Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(3) f(x) = \begin{cases} \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(3) f(x) = \begin{cases} \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(4) f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(3) f(x) = \begin{cases} \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(4) f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

$$(5) f(x) = \begin{cases} x & \text{if } x \in \mathbb{Q}, \\ -x & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

Result: Let $f, g : D \rightarrow \mathbb{R}$ be continuous at $x_0 \in D$. Then

Result: Let $f, g : D \rightarrow \mathbb{R}$ be continuous at $x_0 \in D$. Then
(a) $f \pm g$, fg and $|f|$ are continuous at x_0 .

Result: Let $f, g : D \rightarrow \mathbb{R}$ be continuous at $x_0 \in D$. Then

(a) $f \pm g$, fg and $|f|$ are continuous at x_0 .

(b) f/g is continuous at x_0 if $g(x) \neq 0$ for all $x \in D$.

Result: Let $f, g : D \rightarrow \mathbb{R}$ be continuous at $x_0 \in D$. Then

(a) $f \pm g$, fg and $|f|$ are continuous at x_0 .

(b) f/g is continuous at x_0 if $g(x) \neq 0$ for all $x \in D$.

Result: Composition of two continuous functions is continuous.

Result: Let $f, g : D \rightarrow \mathbb{R}$ be continuous at $x_0 \in D$. Then

(a) $f \pm g$, fg and $|f|$ are continuous at x_0 .

(b) f/g is continuous at x_0 if $g(x) \neq 0$ for all $x \in D$.

Result: Composition of two continuous functions is continuous.

Further examples of continuous functions:

Polynomial function, Rational function, sine function, cosine function, exponential function, etc.

Result: Let $f, g : D \rightarrow \mathbb{R}$ be continuous at $x_0 \in D$. Then

(a) $f \pm g$, fg and $|f|$ are continuous at x_0 .

(b) f/g is continuous at x_0 if $g(x) \neq 0$ for all $x \in D$.

Result: Composition of two continuous functions is continuous.

Further examples of continuous functions:

Polynomial function, Rational function, sine function, cosine function, exponential function, etc.

Result: If $f : D \rightarrow \mathbb{R}$ is continuous at $x_0 \in D$ and $f(x_0) \neq 0$, then there exists $\delta > 0$ such that $f(x) \neq 0$ for all $x \in D$ satisfying $|x - x_0| < \delta$.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if $f(a) \cdot f(b) < 0$, then there exists $c \in (a, b)$ such that $f(c) = 0$.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if $f(a) \cdot f(b) < 0$, then there exists $c \in (a, b)$ such that $f(c) = 0$.

Intermediate value theorem: Let I be an interval of \mathbb{R} and let $f : I \rightarrow \mathbb{R}$ be continuous. If $a, b \in I$ with $a < b$ and if $f(a) < k < f(b)$, then there exists $c \in (a, b)$ such that $f(c) = k$.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if $f(a) \cdot f(b) < 0$, then there exists $c \in (a, b)$ such that $f(c) = 0$.

Intermediate value theorem: Let I be an interval of \mathbb{R} and let $f : I \rightarrow \mathbb{R}$ be continuous. If $a, b \in I$ with $a < b$ and if $f(a) < k < f(b)$, then there exists $c \in (a, b)$ such that $f(c) = k$.

Examples:

- a The equation $x^2 = x \sin x + \cos x$ has at least two real roots.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if $f(a) \cdot f(b) < 0$, then there exists $c \in (a, b)$ such that $f(c) = 0$.

Intermediate value theorem: Let I be an interval of \mathbb{R} and let $f : I \rightarrow \mathbb{R}$ be continuous. If $a, b \in I$ with $a < b$ and if $f(a) < k < f(b)$, then there exists $c \in (a, b)$ such that $f(c) = k$.

Examples:

- a The equation $x^2 = x \sin x + \cos x$ has at least two real roots.
- b (Fixed point) If $f : [a, b] \rightarrow [a, b]$ is continuous, then there exists $c \in [a, b]$ such that $f(c) = c$.

Recall that a function $f : D \rightarrow \mathbb{R}$ is called bounded if there exists $M > 0$ such that $|f(x)| < M$ for all $x \in D$.

Recall that a function $f : D \rightarrow \mathbb{R}$ is called bounded if there exists $M > 0$ such that $|f(x)| < M$ for all $x \in D$.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous, then f is bounded.

Recall that a function $f : D \rightarrow \mathbb{R}$ is called bounded if there exists $M > 0$ such that $|f(x)| < M$ for all $x \in D$.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous, then f is bounded.

Example: There does not exist any continuous function from $[0, 1]$ onto $(0, \infty)$.

Recall that a function $f : D \rightarrow \mathbb{R}$ is called bounded if there exists $M > 0$ such that $|f(x)| < M$ for all $x \in D$.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous, then f is bounded.

Example: There does not exist any continuous function from $[0, 1]$ onto $(0, \infty)$.

Result: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous, then the supremum and the infimum of $f(x)$ are attained in $[a, b]$. That is, there exist $x_0, y_0 \in [a, b]$ such that $f(x_0) \leq f(x) \leq f(y_0)$ for all $x \in [a, b]$.

Limit point of a set: Let $A \subseteq \mathbb{R}$. A real number x is called a **limit point** of A if there exists a sequence (x_n) in A converging to x .

Limit point of a set: Let $A \subseteq \mathbb{R}$. A real number x is called a **limit point** of A if there exists a sequence (x_n) in A converging to x .

Let $A \subseteq \mathbb{R}$. Then A is called a closed set if A contains all its limit points. That is, if (x_n) is a sequence in A converging to x , then $x \in A$.

Limit point of a set: Let $A \subseteq \mathbb{R}$. A real number x is called a **limit point** of A if there exists a sequence (x_n) in A converging to x .

Let $A \subseteq \mathbb{R}$. Then A is called a closed set if A contains all its limit points. That is, if (x_n) is a sequence in A converging to x , then $x \in A$.

For example, \mathbb{R} , $[a, b]$, $\{x_1, x_2, \dots, x_n\}$, \mathbb{N} are closed sets. But, (a, b) , \mathbb{Q} are not closed sets.

Limit point of a set: Let $A \subseteq \mathbb{R}$. A real number x is called a **limit point** of A if there exists a sequence (x_n) in A converging to x .

Let $A \subseteq \mathbb{R}$. Then A is called a closed set if A contains all its limit points. That is, if (x_n) is a sequence in A converging to x , then $x \in A$.

For example, \mathbb{R} , $[a, b]$, $\{x_1, x_2, \dots, x_n\}$, \mathbb{N} are closed sets. But, (a, b) , \mathbb{Q} are not closed sets.

Result: Let A be a closed and bounded subset of \mathbb{R} . If $f : A \rightarrow \mathbb{R}$ is continuous, then f is bounded.

Limit point of a set: Let $A \subseteq \mathbb{R}$. A real number x is called a **limit point** of A if there exists a sequence (x_n) in A converging to x .

Let $A \subseteq \mathbb{R}$. Then A is called a closed set if A contains all its limit points. That is, if (x_n) is a sequence in A converging to x , then $x \in A$.

For example, \mathbb{R} , $[a, b]$, $\{x_1, x_2, \dots, x_n\}$, \mathbb{N} are closed sets. But, (a, b) , \mathbb{Q} are not closed sets.

Result: Let A be a closed and bounded subset of \mathbb{R} . If $f : A \rightarrow \mathbb{R}$ is continuous, then f is bounded.

Remark: The above result is not true if A is bounded but not closed. For example $f(x) = 1/x$ on $(0, 1)$. Also, the result is not true if A is closed but not bounded. For example, $f(x) = x$ on \mathbb{R} .