

# MA 101 (Mathematics-I)

Subhamay Saha and Ayon Ganguly  
Department of Mathematics  
IIT Guwahati

# Introduction

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  be such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ .

# Introduction

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  be such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ .

If  $f : D \rightarrow \mathbb{R}$ , then  $\ell \in \mathbb{R}$  is said to be the limit of  $f$  at  $x_0$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

whenever  $x \in D$  and  $0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ .

# Introduction

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  be such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ .

If  $f : D \rightarrow \mathbb{R}$ , then  $\ell \in \mathbb{R}$  is said to be the limit of  $f$  at  $x_0$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

whenever  $x \in D$  and  $0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ .

We write:  $\lim_{x \rightarrow x_0} f(x) = \ell$ .

# Introduction

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  be such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ .

If  $f : D \rightarrow \mathbb{R}$ , then  $\ell \in \mathbb{R}$  is said to be the limit of  $f$  at  $x_0$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$\text{whenever } x \in D \text{ and } 0 < |x - x_0| < \delta \Rightarrow |f(x) - \ell| < \varepsilon.$$

We write:  $\lim_{x \rightarrow x_0} f(x) = \ell$ .

**Result:** If limit exists, then it is unique.

**Example:**  $\lim_{x \rightarrow 1} \left( \frac{3x}{2} - 1 \right) = \frac{1}{2}.$

Let  $\varepsilon > 0$ . We have to find  $\delta > 0$  such that

$0 < |x - 1| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$  holds with  $\ell = 1/2$ .

Working backwards,

$$\frac{3}{2}|x - 1| < \varepsilon \text{ whenever } |x - 1| < \delta := \frac{2}{3}\varepsilon.$$

**Example:**  $\lim_{x \rightarrow 1} \left( \frac{3x}{2} - 1 \right) = \frac{1}{2}.$

Let  $\varepsilon > 0$ . We have to find  $\delta > 0$  such that

$0 < |x - 1| < \delta \Rightarrow |f(x) - \ell| < \varepsilon$  holds with  $\ell = 1/2$ .

Working backwards,

$$\frac{3}{2}|x - 1| < \varepsilon \text{ whenever } |x - 1| < \delta := \frac{2}{3}\varepsilon.$$

## Theorem (Sequential criterion)


Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ . Let  $f : D \rightarrow \mathbb{R}$ . Then the following are equivalent.

- (a)  $\lim_{x \rightarrow x_0} f(x) = \ell.$
- (b) For any sequence  $(x_n)$  in  $D$  with  $x_n \neq x_0$  for all  $n \geq 1$  and  $x_n \rightarrow x_0$ , the sequence  $(f(x_n))$  converges to  $\ell$ .


**Example:**  $\lim_{x \rightarrow 0} \sin \frac{1}{x}$  does not exist.



**Example:**  $\lim_{x \rightarrow 0} \sin \frac{1}{x}$  does not exist.




**Example:**  $\lim_{x \rightarrow 0} \sin \frac{1}{x}$  does not exist.



**Solution:** Let  $x_n = \frac{2}{(4n+1)\pi}$  and  $y_n = \frac{1}{n\pi}$  for all  $n \in \mathbb{N}$ . Then  $x_n \rightarrow 0$  and  $y_n \rightarrow 0$ .


**Example:**  $\lim_{x \rightarrow 0} \sin \frac{1}{x}$  does not exist.



**Solution:** Let  $x_n = \frac{2}{(4n+1)\pi}$  and  $y_n = \frac{1}{n\pi}$  for all  $n \in \mathbb{N}$ . Then  $x_n \rightarrow 0$  and  $y_n \rightarrow 0$ .

Since  $\sin \frac{1}{x_n} = 1$  and  $\sin \frac{1}{y_n} = 0$  for all  $n \in \mathbb{N}$ , we get  $\sin \frac{1}{x_n} \rightarrow 1$  and  $\sin \frac{1}{y_n} \rightarrow 0$ .

**Example:**  $\lim_{x \rightarrow 0} \sin \frac{1}{x}$  does not exist.



**Solution:** Let  $x_n = \frac{2}{(4n+1)\pi}$  and  $y_n = \frac{1}{n\pi}$  for all  $n \in \mathbb{N}$ . Then  $x_n \rightarrow 0$  and  $y_n \rightarrow 0$ .

Since  $\sin \frac{1}{x_n} = 1$  and  $\sin \frac{1}{y_n} = 0$  for all  $n \in \mathbb{N}$ , we get  $\sin \frac{1}{x_n} \rightarrow 1$  and  $\sin \frac{1}{y_n} \rightarrow 0$ .

Therefore by the sequential criterion for limit,  $\lim_{x \rightarrow 0} \sin \frac{1}{x}$  does not exist.

A function  $f : D \rightarrow \mathbb{R}$  is called bounded if there exists  $M > 0$  such that  $|f(x)| < M$  for all  $x \in D$ .

A function  $f : D \rightarrow \mathbb{R}$  is called bounded if there exists  $M > 0$  such that  $|f(x)| < M$  for all  $x \in D$ .

**Result:** Let  $f : D \rightarrow \mathbb{R}$ . Suppose that  $\lim_{x \rightarrow x_0} f(x) = \ell$ . Then there exists some  $\delta > 0$  such that  $f$  is bounded on  $(x_0 - \delta, x_0 + \delta) \setminus \{x_0\}$ . That is, there exists  $M > 0$  such that  $|f(x)| < M$  for all  $x \in (x_0 - \delta, x_0 + \delta)$  with  $x \neq x_0$ .

## Theorem (Limit Theorems)

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ . Let  $f, g, j : D \rightarrow \mathbb{R}$ . Suppose that  $\lim_{x \rightarrow x_0} f(x) = \ell$  and  $\lim_{x \rightarrow x_0} g(x) = m$ . Then

## Theorem (Limit Theorems)

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ . Let  $f, g, j : D \rightarrow \mathbb{R}$ . Suppose that  $\lim_{x \rightarrow x_0} f(x) = \ell$  and  $\lim_{x \rightarrow x_0} g(x) = m$ . Then

(1)  $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$ .



## Theorem (Limit Theorems)

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ . Let  $f, g, j : D \rightarrow \mathbb{R}$ . Suppose that  $\lim_{x \rightarrow x_0} f(x) = \ell$  and  $\lim_{x \rightarrow x_0} g(x) = m$ . Then

- (1)  $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$ .
- (2) If  $f(x) \leq g(x)$  for all  $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$ , then  $\ell \leq m$ .

## Theorem (Limit Theorems)

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ . Let  $f, g, j : D \rightarrow \mathbb{R}$ . Suppose that  $\lim_{x \rightarrow x_0} f(x) = \ell$  and  $\lim_{x \rightarrow x_0} g(x) = m$ . Then

- (1)  $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$ .
- (2) If  $f(x) \leq g(x)$  for all  $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$ , then  $\ell \leq m$ .
- (3)  $\lim_{x \rightarrow x_0} (fg)(x) = \ell m$  and if  $m \neq 0$  and  $g(x) \neq 0$  for all  $x \in D$ , then  $\lim_{x \rightarrow x_0} \frac{1}{g(x)} = \frac{1}{m}$ .

## Theorem (Limit Theorems)

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0 - h, x_0 + h) \setminus \{x_0\} \subseteq D$ . Let  $f, g, j : D \rightarrow \mathbb{R}$ . Suppose that  $\lim_{x \rightarrow x_0} f(x) = \ell$  and  $\lim_{x \rightarrow x_0} g(x) = m$ . Then

- (1)  $\lim_{x \rightarrow x_0} (f(x) \pm g(x)) = \ell \pm m$ .
- (2) If  $f(x) \leq g(x)$  for all  $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$ , then  $\ell \leq m$ .
- (3)  $\lim_{x \rightarrow x_0} (fg)(x) = \ell m$  and if  $m \neq 0$  and  $g(x) \neq 0$  for all  $x \in D$ , then  $\lim_{x \rightarrow x_0} \frac{1}{g(x)} = \frac{1}{m}$ .
- (4) If  $f(x) \leq j(x) \leq g(x)$  for all  $x \in (x_0 - h, x_0 + h) \setminus \{x_0\}$  and  $\ell = m$ , then  $\lim_{x \rightarrow x_0} j(x) = \ell$ .

**Result:** Suppose that  $f(x)$  is bounded in  $(x_0 - h, x_0 + h) \setminus \{x_0\}$  for some  $h > 0$  and  $\lim_{x \rightarrow x_0} g(x) = 0$ .  
Then  $\lim_{x \rightarrow x_0} f(x)g(x) = 0$ .

**Result:** Suppose that  $f(x)$  is bounded in  $(x_0 - h, x_0 + h) \setminus \{x_0\}$  for some  $h > 0$  and  $\lim_{x \rightarrow x_0} g(x) = 0$ .


Then  $\lim_{x \rightarrow x_0} f(x)g(x) = 0$ .

**Example:**  $\lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0$ .

**Result:** Suppose that  $f(x)$  is bounded in  $(x_0 - h, x_0 + h) \setminus \{x_0\}$  for some  $h > 0$  and  $\lim_{x \rightarrow x_0} g(x) = 0$ .

Then  $\lim_{x \rightarrow x_0} f(x)g(x) = 0$ .

**Example:**  $\lim_{x \rightarrow 0} x \sin \frac{1}{x} = 0$ .



## One sided limits

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  
 $(x_0, x_0 + h) \subseteq D$ .

## One sided limits

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0, x_0 + h) \subseteq D$ .

If  $f : D \rightarrow \mathbb{R}$ , then  $\ell \in \mathbb{R}$  is said to be the right hand limit of  $f$  at  $x_0$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

whenever  $x \in D$  and  $0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ .



## One sided limits

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0, x_0 + h) \subseteq D$ .

If  $f : D \rightarrow \mathbb{R}$ , then  $\ell \in \mathbb{R}$  is said to be the right hand limit of  $f$  at  $x_0$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

whenever  $x \in D$  and  $0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon$ .

We write:  $\lim_{x \rightarrow x_0+} f(x) = \ell$ .

## One sided limits

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0, x_0 + h) \subseteq D$ .

If  $f : D \rightarrow \mathbb{R}$ , then  $\ell \in \mathbb{R}$  is said to be the right hand limit of  $f$  at  $x_0$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$\text{whenever } x \in D \text{ and } 0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon.$$

We write:  $\lim_{x \rightarrow x_0+} f(x) = \ell$ .

Similarly one defines left hand limit of  $f$  at  $x_0$  and is denoted by  $\lim_{x \rightarrow x_0-} f(x)$ .

## One sided limits

Let  $D \subseteq \mathbb{R}$  and let  $x_0 \in \mathbb{R}$  such that for some  $h > 0$ ,  $(x_0, x_0 + h) \subseteq D$ .

If  $f : D \rightarrow \mathbb{R}$ , then  $\ell \in \mathbb{R}$  is said to be the right hand limit of  $f$  at  $x_0$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$$\text{whenever } x \in D \text{ and } 0 < x - x_0 < \delta \Rightarrow |f(x) - \ell| < \varepsilon.$$

We write:  $\lim_{x \rightarrow x_0+} f(x) = \ell$ .


Similarly one defines left hand limit of  $f$  at  $x_0$  and is denoted by  $\lim_{x \rightarrow x_0-} f(x)$ .

**Result:**  $\lim_{x \rightarrow x_0} f(x) = \ell \Leftrightarrow \lim_{x \rightarrow x_0+} f(x) = \lim_{x \rightarrow x_0-} f(x) = \ell$ .

**Example:** Show that  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$ .

**Example:** Show that  $\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} = 1$ .

We will use Sandwich Thm for limit  
(Limit Thm (4))



**Area of  $\triangle OAP$  < Area of the circular section OAPQ < Area of  $\triangle OAT$**

## Limits at infinity and infinite limits

**Definition:**  $f(x)$  has limit  $\ell$  as  $x$  approaches  $+\infty$ , if for any given  $\varepsilon > 0$ , there exists  $M > 0$  such that

$$x > M \implies |f(x) - \ell| < \varepsilon.$$

Similarly, one can define limit of  $f(x)$  as  $x$  approaches  $-\infty$ .

# Limits at infinity and infinite limits

**Definition:**  $f(x)$  has limit  $\ell$  as  $x$  approaches  $+\infty$ , if for any given  $\varepsilon > 0$ , there exists  $M > 0$  such that

$$x > M \implies |f(x) - \ell| < \varepsilon.$$

Similarly, one can define limit of  $f(x)$  as  $x$  approaches  $-\infty$ .

**Example:** (i)  $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$ , (ii)  $\lim_{x \rightarrow -\infty} \frac{1}{x} = 0$ ,  
 (iii)  $\lim_{x \rightarrow \infty} \sin x$  does not exist.

## Limits at infinity and infinite limits

**Definition:**  $f(x)$  has limit  $\ell$  as  $x$  approaches  $+\infty$ , if for any given  $\varepsilon > 0$ , there exists  $M > 0$  such that

$$x > M \implies |f(x) - \ell| < \varepsilon.$$

Similarly, one can define limit of  $f(x)$  as  $x$  approaches  $-\infty$ .

**Example:** (i)  $\lim_{x \rightarrow \infty} \frac{1}{x} = 0$ , (ii)  $\lim_{x \rightarrow -\infty} \frac{1}{x} = 0$ ,  
 (iii)  $\lim_{x \rightarrow \infty} \sin x$  does not exist.

**Definition:** A function  $f(x)$  approaches  $\infty$  ( $f(x) \rightarrow \infty$ ) as  $x \rightarrow x_0$  if, for every real  $M > 0$ , there exists  $\delta > 0$  such that

$$0 < |x - x_0| < \delta \implies f(x) > M.$$

Similarly, one can define limit of  $f(x)$  approaching  $-\infty$ .



**Example:** (i)  $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$ , (ii)  $\lim_{x \rightarrow 0} \frac{1}{x^2} \sin(1/x)$  does not exist.

**Example:** (i)  $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$ , (ii)  $\lim_{x \rightarrow 0} \frac{1}{x^2} \sin(1/x)$  does not exist.

For (ii), let  $x_n = \frac{1}{\frac{\pi}{2} + 2n\pi}$  and  $y_n = \frac{1}{n\pi}$ . Then  $x_n, y_n \rightarrow 0$  as  $n \rightarrow \infty$ .

But  $\lim_{n \rightarrow \infty} f(x_n) = \frac{1}{x_n^2} \rightarrow \infty$  and  $\lim_{n \rightarrow \infty} f(y_n) = 0$ .

**Example:** (i)  $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$ , (ii)  $\lim_{x \rightarrow 0} \frac{1}{x^2} \sin(1/x)$  does not exist.

For (ii), let  $x_n = \frac{1}{\frac{\pi}{2} + 2n\pi}$  and  $y_n = \frac{1}{n\pi}$ . Then  $x_n, y_n \rightarrow 0$  as  $n \rightarrow \infty$ .

But  $\lim_{n \rightarrow \infty} f(x_n) = \frac{1}{x_n^2} \rightarrow \infty$  and  $\lim_{n \rightarrow \infty} f(y_n) = 0$ .

## Theorem

*Suppose that  $\lim_{x \rightarrow x_0} f(x) = \ell$ . If  $\ell \neq 0$ , then there exists some  $\delta$  such that  $f(x) \neq 0$  for all  $x \in (x_0 - \delta, x_0 + \delta) \setminus \{x_0\}$ .*

## Continuous functions

Let  $D$  be a nonempty subset of  $\mathbb{R}$  and  $f : D \rightarrow \mathbb{R}$ . We say that  $f$  is continuous at  $x_0 \in D$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $|f(x) - f(x_0)| < \varepsilon$  for all  $x \in D$  satisfying  $|x - x_0| < \delta$ .

## Continuous functions

Let  $D$  be a nonempty subset of  $\mathbb{R}$  and  $f : D \rightarrow \mathbb{R}$ . We say that  $f$  is continuous at  $x_0 \in D$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $|f(x) - f(x_0)| < \varepsilon$  for all  $x \in D$  satisfying  $|x - x_0| < \delta$ .

Let  $f : [a, b] \rightarrow \mathbb{R}$ . Then  $f$  is continuous at  $c \in (a, b)$  if  $\lim_{x \rightarrow c} f(x) = f(c)$ .  $f$  is continuous at  $a$  if  $\lim_{x \rightarrow a+} f(x) = f(a)$ . Similarly,  $f$  is continuous at  $b$  if  $\lim_{x \rightarrow b-} f(x) = f(b)$ .

## Continuous functions

Let  $D$  be a nonempty subset of  $\mathbb{R}$  and  $f : D \rightarrow \mathbb{R}$ . We say that  $f$  is continuous at  $x_0 \in D$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $|f(x) - f(x_0)| < \varepsilon$  for all  $x \in D$  satisfying  $|x - x_0| < \delta$ .

Let  $f : [a, b] \rightarrow \mathbb{R}$ . Then  $f$  is continuous at  $c \in (a, b)$  if  $\lim_{x \rightarrow c} f(x) = f(c)$ .  $f$  is continuous at  $a$  if  $\lim_{x \rightarrow a+} f(x) = f(a)$ . Similarly,  $f$  is continuous at  $b$  if  $\lim_{x \rightarrow b-} f(x) = f(b)$ .

**Sequential criterion of continuity:**  $f : D \rightarrow \mathbb{R}$  is continuous at  $x_0 \in D$  if and only if for every sequence  $(x_n)$  in  $D$  such that  $x_n \rightarrow x_0$ , we have  $f(x_n) \rightarrow f(x_0)$ .

## Continuous functions

Let  $D$  be a nonempty subset of  $\mathbb{R}$  and  $f : D \rightarrow \mathbb{R}$ . We say that  $f$  is continuous at  $x_0 \in D$  if for each  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $|f(x) - f(x_0)| < \varepsilon$  for all  $x \in D$  satisfying  $|x - x_0| < \delta$ .

Let  $f : [a, b] \rightarrow \mathbb{R}$ . Then  $f$  is continuous at  $c \in (a, b)$  if  $\lim_{x \rightarrow c} f(x) = f(c)$ .  $f$  is continuous at  $a$  if  $\lim_{x \rightarrow a+} f(x) = f(a)$ . Similarly,  $f$  is continuous at  $b$  if  $\lim_{x \rightarrow b-} f(x) = f(b)$ .

**Sequential criterion of continuity:**  $f : D \rightarrow \mathbb{R}$  is continuous at  $x_0 \in D$  if and only if for every sequence  $(x_n)$  in  $D$  such that  $x_n \rightarrow x_0$ , we have  $f(x_n) \rightarrow f(x_0)$ .

We say that  $f : D \rightarrow \mathbb{R}$  is continuous if  $f$  is continuous at each  $x_0 \in D$ .

**Result:** Let  $f, g : D \rightarrow \mathbb{R}$  be continuous at  $x_0 \in D$ . Then



**Result:** Let  $f, g : D \rightarrow \mathbb{R}$  be continuous at  $x_0 \in D$ . Then  
(a)  $f \pm g$ ,  $fg$  and  $|f|$  are continuous at  $x_0$ .

**Result:** Let  $f, g : D \rightarrow \mathbb{R}$  be continuous at  $x_0 \in D$ . Then

(a)  $f \pm g$ ,  $fg$  and  $|f|$  are continuous at  $x_0$ .

(b)  $f/g$  is continuous at  $x_0$  if  $g(x) \neq 0$  for all  $x \in D$ .

**Result:** Let  $f, g : D \rightarrow \mathbb{R}$  be continuous at  $x_0 \in D$ . Then

(a)  $f \pm g$ ,  $fg$  and  $|f|$  are continuous at  $x_0$ .

(b)  $f/g$  is continuous at  $x_0$  if  $g(x) \neq 0$  for all  $x \in D$ .

**Result:** Composition of two continuous functions is continuous.

**Result:** Let  $f, g : D \rightarrow \mathbb{R}$  be continuous at  $x_0 \in D$ . Then

(a)  $f \pm g$ ,  $fg$  and  $|f|$  are continuous at  $x_0$ .

(b)  $f/g$  is continuous at  $x_0$  if  $g(x) \neq 0$  for all  $x \in D$ .

**Result:** Composition of two continuous functions is continuous.

Standard examples of continuous functions:

Polynomial function, Rational function, sine function, cosine function, exponential function, etc.

**Result:** Let  $f, g : D \rightarrow \mathbb{R}$  be continuous at  $x_0 \in D$ . Then

(a)  $f \pm g$ ,  $fg$  and  $|f|$  are continuous at  $x_0$ .

(b)  $f/g$  is continuous at  $x_0$  if  $g(x) \neq 0$  for all  $x \in D$ .

**Result:** Composition of two continuous functions is continuous.

Standard examples of continuous functions:

Polynomial function, Rational function, sine function, cosine function, exponential function, etc.

**Result:** If  $f : D \rightarrow \mathbb{R}$  is continuous at  $x_0 \in D$  and  $f(x_0) \neq 0$ , then there exists  $\delta > 0$  such that  $f(x) \neq 0$  for all  $x \in D$  satisfying  $|x - x_0| < \delta$ .

# Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

## Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

# Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(3) f(x) = \begin{cases} \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$



## Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(3) f(x) = \begin{cases} \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(4) f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

## Examples

$$(1) f(x) = \begin{cases} 3x + 2 & \text{if } x < 1, \\ 4x^2 & \text{if } x \geq 1. \end{cases}$$

$$(2) f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(3) f(x) = \begin{cases} \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

$$(4) f(x) = \begin{cases} 1 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

$$(5) f(x) = \begin{cases} x & \text{if } x \in \mathbb{Q}, \\ -x & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$$

## Intermediate value theorem

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous and if  $f(a) \cdot f(b) < 0$ , then there exists  $c \in (a, b)$  such that  $f(c) = 0$ .

## Intermediate value theorem

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous and if  $f(a) \cdot f(b) < 0$ , then there exists  $c \in (a, b)$  such that  $f(c) = 0$ .

**Intermediate value theorem:** Let  $I$  be an interval of  $\mathbb{R}$  and let  $f : I \rightarrow \mathbb{R}$  be continuous. If  $a, b \in I$  with  $a < b$  and if  $f(a) < k < f(b)$ , then there exists  $c \in (a, b)$  such that  $f(c) = k$ .

## Intermediate value theorem

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous and if  $f(a) \cdot f(b) < 0$ , then there exists  $c \in (a, b)$  such that  $f(c) = 0$ .

**Intermediate value theorem:** Let  $I$  be an interval of  $\mathbb{R}$  and let  $f : I \rightarrow \mathbb{R}$  be continuous. If  $a, b \in I$  with  $a < b$  and if  $f(a) < k < f(b)$ , then there exists  $c \in (a, b)$  such that  $f(c) = k$ .

**Examples:**

- a The equation  $x^2 = x \sin x + \cos x$  has at least two real roots.

## Intermediate value theorem

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous and if  $f(a) \cdot f(b) < 0$ , then there exists  $c \in (a, b)$  such that  $f(c) = 0$ .

**Intermediate value theorem:** Let  $I$  be an interval of  $\mathbb{R}$  and let  $f : I \rightarrow \mathbb{R}$  be continuous. If  $a, b \in I$  with  $a < b$  and if  $f(a) < k < f(b)$ , then there exists  $c \in (a, b)$  such that  $f(c) = k$ .

**Examples:**

- a The equation  $x^2 = x \sin x + \cos x$  has at least two real roots.
- b (Fixed point) If  $f : [a, b] \rightarrow [a, b]$  is continuous, then there exists  $c \in [a, b]$  such that  $f(c) = c$ .

Recall that a function  $f : D \rightarrow \mathbb{R}$  is called bounded if there exists  $M > 0$  such that  $|f(x)| < M$  for all  $x \in D$ .

Recall that a function  $f : D \rightarrow \mathbb{R}$  is called bounded if there exists  $M > 0$  such that  $|f(x)| < M$  for all  $x \in D$ .

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f$  is bounded.



Recall that a function  $f : D \rightarrow \mathbb{R}$  is called bounded if there exists  $M > 0$  such that  $|f(x)| < M$  for all  $x \in D$ .

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f$  is bounded.

**Example:** There does not exist any continuous function from  $[0, 1]$  onto  $(0, \infty)$ .

Recall that a function  $f : D \rightarrow \mathbb{R}$  is called bounded if there exists  $M > 0$  such that  $|f(x)| < M$  for all  $x \in D$ .

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous, then  $f$  is bounded.

**Example:** There does not exist any continuous function from  $[0, 1]$  onto  $(0, \infty)$ .

**Result:** If  $f : [a, b] \rightarrow \mathbb{R}$  is continuous, then the supremum and the infimum of  $f(x)$  are attained in  $[a, b]$ . That is, there exist  $x_0, y_0 \in [a, b]$  such that  $f(x_0) \leq f(x) \leq f(y_0)$  for all  $x \in [a, b]$ .

**Limit point of a set:** Let  $A \subseteq \mathbb{R}$ . A real number  $x$  is called a **limit point** of  $A$  if there exists a sequence  $(x_n)$  in  $A$  converging to  $x$ .

**Limit point of a set:** Let  $A \subseteq \mathbb{R}$ . A real number  $x$  is called a **limit point** of  $A$  if there exists a sequence  $(x_n)$  in  $A$  converging to  $x$ .

Let  $A \subseteq \mathbb{R}$ . Then  $A$  is called a closed set if  $A$  contains all its limit points. That is, if  $(x_n)$  is a sequence in  $A$  converging to  $x$ , then  $x \in A$ .

**Limit point of a set:** Let  $A \subseteq \mathbb{R}$ . A real number  $x$  is called a **limit point** of  $A$  if there exists a sequence  $(x_n)$  in  $A$  converging to  $x$ .

Let  $A \subseteq \mathbb{R}$ . Then  $A$  is called a closed set if  $A$  contains all its limit points. That is, if  $(x_n)$  is a sequence in  $A$  converging to  $x$ , then  $x \in A$ .

For example,  $\mathbb{R}$ ,  $[a, b]$ ,  $\{x_1, x_2, \dots, x_n\}$ ,  $\mathbb{N}$  are closed sets. But,  $(a, b)$ ,  $\mathbb{Q}$  are not closed sets.

**Limit point of a set:** Let  $A \subseteq \mathbb{R}$ . A real number  $x$  is called a **limit point** of  $A$  if there exists a sequence  $(x_n)$  in  $A$  converging to  $x$ .

Let  $A \subseteq \mathbb{R}$ . Then  $A$  is called a closed set if  $A$  contains all its limit points. That is, if  $(x_n)$  is a sequence in  $A$  converging to  $x$ , then  $x \in A$ .

For example,  $\mathbb{R}$ ,  $[a, b]$ ,  $\{x_1, x_2, \dots, x_n\}$ ,  $\mathbb{N}$  are closed sets. But,  $(a, b)$ ,  $\mathbb{Q}$  are not closed sets.

**Result:** Let  $A$  be a closed and bounded subset of  $\mathbb{R}$ . If  $f : A \rightarrow \mathbb{R}$  is continuous, then  $f$  is bounded.

**Limit point of a set:** Let  $A \subseteq \mathbb{R}$ . A real number  $x$  is called a **limit point** of  $A$  if there exists a sequence  $(x_n)$  in  $A$  converging to  $x$ .

Let  $A \subseteq \mathbb{R}$ . Then  $A$  is called a closed set if  $A$  contains all its limit points. That is, if  $(x_n)$  is a sequence in  $A$  converging to  $x$ , then  $x \in A$ .

For example,  $\mathbb{R}$ ,  $[a, b]$ ,  $\{x_1, x_2, \dots, x_n\}$ ,  $\mathbb{N}$  are closed sets. But,  $(a, b)$ ,  $\mathbb{Q}$  are not closed sets.

**Result:** Let  $A$  be a closed and bounded subset of  $\mathbb{R}$ . If  $f : A \rightarrow \mathbb{R}$  is continuous, then  $f$  is bounded.

**Remark:** The above result is not true if  $A$  is bounded but not closed. For example  $f(x) = 1/x$  on  $(0, 1)$ . Also, the result is not true if  $A$  is closed but not bounded. For example,  $f(x) = x$  on  $\mathbb{R}$ .