

# MA 101 (Mathematics-I)

# Introduction

Consider the “sequence” of rationals

1.4, 1.41, 1.414, 1.4142, 1.41421, ...

# Introduction

Consider the “sequence” of rationals

$$1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$$

Their squares are:

$$1.96, 1.9881, 1.99396, 1.99996164, 1.9999899241, \dots$$

# Introduction

Consider the “sequence” of rationals

$$1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$$

Their squares are:

$$1.96, 1.9881, 1.99396, 1.99996164, 1.9999899241, \dots$$

It seems that the above numbers are getting closer and closer to 2.

# Introduction

Consider the “sequence” of rationals

$$1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$$

Their squares are:

$$1.96, 1.9881, 1.99396, 1.99996164, 1.9999899241, \dots$$

It seems that the above numbers are getting closer and closer to 2.

Therefore, the rationals  $1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$  are getting closer and closer to  $\sqrt{2}$ .

## Definition (Sequence)

A sequence of real numbers or a sequence in  $\mathbb{R}$  is a mapping  $f : \mathbb{N} \rightarrow \mathbb{R}$ . We write  $x_n$  for  $f(n)$ ,  $n \in \mathbb{N}$  and it is customary to denote a sequence as  $\langle x_n \rangle$  or  $(x_n)$  or  $\{x_n\}$ .

## Definition (Sequence)

A sequence of real numbers or a sequence in  $\mathbb{R}$  is a mapping  $f : \mathbb{N} \rightarrow \mathbb{R}$ . We write  $x_n$  for  $f(n)$ ,  $n \in \mathbb{N}$  and it is customary to denote a sequence as  $\langle x_n \rangle$  or  $(x_n)$  or  $\{x_n\}$ .

## Example

*There are different ways of expressing a sequence. For example:*

- ① *Constant sequence:  $(a, a, a, \dots)$ , where  $a \in \mathbb{R}$*
- ② *Sequence defined by listing:  $(1, 4, 8, 11, 52, \dots)$*
- ③ *Sequence defined by rule:  $(x_n)$ , where  $x_n = 3n^2$  for all  $n \in \mathbb{N}$*
- ④ *Sequence defined recursively:  $(x_n)$ , where  $x_1 = 4$  and  $x_{n+1} = 2x_n - 5$  for all  $n \in \mathbb{N}$*

## Convergence: What does it mean?

Think of the examples:

①  $(2, 2, 2, \dots)$

②  $(\frac{1}{n})$

③  $((-1)^n \frac{1}{n})$

④  $(1, 2, 1, 2, \dots)$

⑤  $(\sqrt{n})$

⑥  $((-1)^n(1 - \frac{1}{n}))$



## Definition (Convergent sequence)

A sequence  $(x_n)$  is said to be convergent if there exists  $\ell \in \mathbb{R}$  such that for every  $\varepsilon > 0$ , there exists  $n_0 \in \mathbb{N}$  satisfying  $|x_n - \ell| < \varepsilon$  for all  $n \geq n_0$ . We say that  $\ell$  is a limit of  $(x_n)$ .

## Definition (Convergent sequence)

A sequence  $(x_n)$  is said to be convergent if there exists  $\ell \in \mathbb{R}$  such that for every  $\varepsilon > 0$ , there exists  $n_0 \in \mathbb{N}$  satisfying  $|x_n - \ell| < \varepsilon$  for all  $n \geq n_0$ . We say that  $\ell$  is a limit of  $(x_n)$ .

**Notation:** We write  $\lim_{n \rightarrow \infty} x_n = \ell$  or  $x_n \rightarrow \ell$ .

## Definition (Convergent sequence)

A sequence  $(x_n)$  is said to be convergent if there exists  $\ell \in \mathbb{R}$  such that for every  $\varepsilon > 0$ , there exists  $n_0 \in \mathbb{N}$  satisfying  $|x_n - \ell| < \varepsilon$  for all  $n \geq n_0$ . We say that  $\ell$  is a limit of  $(x_n)$ .

**Notation:** We write  $\lim_{n \rightarrow \infty} x_n = \ell$  or  $x_n \rightarrow \ell$ .

## Theorem

*Limit of a convergent sequence is unique.*

## Example

*Using the definition of convergence of a sequence, show that*

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

## Example

*Using the definition of convergence of a sequence, show that*

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

## Example

*If  $p > 0$ , then  $\frac{1}{n^p} \rightarrow 0$ .*

## Example

*Using the definition of convergence of a sequence, show that*

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

## Example

*If  $p > 0$ , then  $\frac{1}{n^p} \rightarrow 0$ .*

## Example

*Consider the sequence  $(x_n)$  where  $x_n = (-1)^n$ . The terms of the sequence are  $-1, 1, -1, 1, -1, 1, \dots$ . It is intuitively clear that this sequence does not approach to any real number. Therefore, the sequence does not converge. We now establish this fact by using the definition.*

**Bounded sequence:** Given a sequence  $(x_n)$ , we can ask whether the set  $\{x_1, x_2, x_3, \dots\}$  is bounded or not. If this set is bounded then we call that the sequence  $(x_n)$  is bounded. Equivalently, the sequence  $(x_n)$  is bounded if there is a positive number  $M$  such that  $|x_n| \leq M$  for all  $n \in \mathbb{N}$ . If  $(x_n)$  is not bounded then it is said to be unbounded.

**Bounded sequence:** Given a sequence  $(x_n)$ , we can ask whether the set  $\{x_1, x_2, x_3, \dots\}$  is bounded or not. If this set is bounded then we call that the sequence  $(x_n)$  is bounded. Equivalently, the sequence  $(x_n)$  is bounded if there is a positive number  $M$  such that  $|x_n| \leq M$  for all  $n \in \mathbb{N}$ . If  $(x_n)$  is not bounded then it is said to be unbounded.

## Theorem

*Every convergent sequence is bounded.*



**Bounded sequence:** Given a sequence  $(x_n)$ , we can ask whether the set  $\{x_1, x_2, x_3, \dots\}$  is bounded or not. If this set is bounded then we call that the sequence  $(x_n)$  is bounded. Equivalently, the sequence  $(x_n)$  is bounded if there is a positive number  $M$  such that  $|x_n| \leq M$  for all  $n \in \mathbb{N}$ . If  $(x_n)$  is not bounded then it is said to be unbounded.

## Theorem

*Every convergent sequence is bounded.*

## Remark

- *From the above theorem, it follows that if a sequence is not bounded then it is not convergent. For example, the sequence  $(\sqrt{n})$  is unbounded and hence is not convergent.*
- *Every bounded sequence is not convergent. For example,  $((-1)^n)$  is a bounded sequence but it does not converge.*

# Limit rules for convergent sequences

## Theorem

Let  $x_n \rightarrow x$  and  $y_n \rightarrow y$ . Then

- (a)  $x_n + y_n \rightarrow x + y$ .
- (b)  $\alpha x_n \rightarrow \alpha x$  for all  $\alpha \in \mathbb{R}$ .
- (c)  $|x_n| \rightarrow |x|$ .
- (d)  $x_n y_n \rightarrow xy$ .
- (e)  $\frac{x_n}{y_n} \rightarrow \frac{x}{y}$  if  $y_n \neq 0$  for all  $n \in \mathbb{N}$  and  $y \neq 0$ .

# Limit rules for convergent sequences

## Theorem

Let  $x_n \rightarrow x$  and  $y_n \rightarrow y$ . Then

- (a)  $x_n + y_n \rightarrow x + y$ .
- (b)  $\alpha x_n \rightarrow \alpha x$  for all  $\alpha \in \mathbb{R}$ .
- (c)  $|x_n| \rightarrow |x|$ .
- (d)  $x_n y_n \rightarrow xy$ .
- (e)  $\frac{x_n}{y_n} \rightarrow \frac{x}{y}$  if  $y_n \neq 0$  for all  $n \in \mathbb{N}$  and  $y \neq 0$ .

## Example

The sequence  $\left(\frac{2n^2-3n}{3n^2+5n+3}\right)$  is convergent with limit  $\frac{2}{3}$ .

# Sandwich theorem

## Theorem (Sandwich theorem)

*Let  $(x_n)$ ,  $(y_n)$ ,  $(z_n)$  be sequences such that  $x_n \leq y_n \leq z_n$  for all  $n \geq n_0$  for some  $n_0 \in \mathbb{N}$ . If both  $(x_n)$  and  $(z_n)$  converge to the same limit  $\ell$ , then  $(y_n)$  also converges to  $\ell$ .*

# Sandwich theorem

## Theorem (Sandwich theorem)

*Let  $(x_n)$ ,  $(y_n)$ ,  $(z_n)$  be sequences such that  $x_n \leq y_n \leq z_n$  for all  $n \geq n_0$  for some  $n_0 \in \mathbb{N}$ . If both  $(x_n)$  and  $(z_n)$  converge to the same limit  $\ell$ , then  $(y_n)$  also converges to  $\ell$ .*

## Example

$$\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$$

# Sandwich theorem

## Theorem (Sandwich theorem)

*Let  $(x_n)$ ,  $(y_n)$ ,  $(z_n)$  be sequences such that  $x_n \leq y_n \leq z_n$  for all  $n \geq n_0$  for some  $n_0 \in \mathbb{N}$ . If both  $(x_n)$  and  $(z_n)$  converge to the same limit  $\ell$ , then  $(y_n)$  also converges to  $\ell$ .*

## Example

$$\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$$

## Example

*The sequence  $(\sqrt{n+1} - \sqrt{n})$  is convergent with limit 0.*

# Sandwich theorem

## Theorem (Sandwich theorem)

Let  $(x_n)$ ,  $(y_n)$ ,  $(z_n)$  be sequences such that  $x_n \leq y_n \leq z_n$  for all  $n \geq n_0$  for some  $n_0 \in \mathbb{N}$ . If both  $(x_n)$  and  $(z_n)$  converge to the same limit  $\ell$ , then  $(y_n)$  also converges to  $\ell$ .

## Example

$$\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$$

## Example

The sequence  $(\sqrt{n+1} - \sqrt{n})$  is convergent with limit 0.

## Example

If  $\alpha > 0$ , then the sequence  $(\alpha^{\frac{1}{n}})$  converges to 1.

## Example

*The sequence  $(n^{\frac{1}{n}})$  converges to 1.*



### Example

*The sequence  $(n^{\frac{1}{n}})$  converges to 1.*

### Example

*If  $p > 0$  and  $\alpha \in \mathbb{R}$ , then  $\frac{n^\alpha}{(1+p)^n} \rightarrow 0$ .*

### Example

*The sequence  $(n^{\frac{1}{n}})$  converges to 1.*

### Example

*If  $p > 0$  and  $\alpha \in \mathbb{R}$ , then  $\frac{n^\alpha}{(1+p)^n} \rightarrow 0$ .*

### Theorem

*Let  $r \in \mathbb{R}$ . Then there exists a sequence  $(x_n)$  of rational numbers such that  $\lim_{n \rightarrow \infty} x_n = r$ .*

# Divergent sequences

A sequence  $(x_n)$  is said to be **divergent** if it has no limit.

# Divergent sequences

A sequence  $(x_n)$  is said to be **divergent** if it has no limit.

## Example

- *If  $(x_n)$  is unbounded then it is divergent.*
- *$(\sqrt{n})$ ,  $(3n^2)$ ,  $((-1)^n n^3)$  are all divergent.*

## Example

*The sequence  $((-1)^n)$  is not convergent, and so it is a divergent sequence although it is bounded.*

## Definition

A sequence  $(x_n)$  is said to approach infinity or diverges to infinity if for any real number  $M > 0$ , there is a positive integer  $n_0$  such that  $x_n \geq M$  for all  $n \geq n_0$ . Similarly,  $(x_n)$  is said to approach  $-\infty$  or diverges to  $-\infty$  if for any real number  $M > 0$ , there is a positive integer  $n_0$  such that  $x_n \leq -M$  for all  $n \geq n_0$ .

# Monotone sequence

## Definition

A sequence  $(x_n)$  is said to be increasing if  $x_{n+1} \geq x_n$  for all  $n \in \mathbb{N}$ . Similarly,  $(x_n)$  is said to be decreasing if  $x_{n+1} \leq x_n$  for all  $n \in \mathbb{N}$ . We say that  $(x_n)$  is monotonic if it is either increasing or decreasing.

# Monotone sequence

## Definition

A sequence  $(x_n)$  is said to be increasing if  $x_{n+1} \geq x_n$  for all  $n \in \mathbb{N}$ . Similarly,  $(x_n)$  is said to be decreasing if  $x_{n+1} \leq x_n$  for all  $n \in \mathbb{N}$ . We say that  $(x_n)$  is monotonic if it is either increasing or decreasing.

- 1 The sequence  $(\frac{1}{n})$  is decreasing.

# Monotone sequence

## Definition

A sequence  $(x_n)$  is said to be increasing if  $x_{n+1} \geq x_n$  for all  $n \in \mathbb{N}$ . Similarly,  $(x_n)$  is said to be decreasing if  $x_{n+1} \leq x_n$  for all  $n \in \mathbb{N}$ . We say that  $(x_n)$  is monotonic if it is either increasing or decreasing.

- 1 The sequence  $(\frac{1}{n})$  is decreasing.
- 2 The sequence  $(n + \frac{1}{n})$  is increasing.



# Monotone sequence

## Definition

A sequence  $(x_n)$  is said to be increasing if  $x_{n+1} \geq x_n$  for all  $n \in \mathbb{N}$ . Similarly,  $(x_n)$  is said to be decreasing if  $x_{n+1} \leq x_n$  for all  $n \in \mathbb{N}$ . We say that  $(x_n)$  is monotonic if it is either increasing or decreasing.

- 1 The sequence  $(\frac{1}{n})$  is decreasing.
- 2 The sequence  $(n + \frac{1}{n})$  is increasing.
- 3 The sequence  $(\cos \frac{n\pi}{3})$  is not monotonic.

# Monotone sequence

## Definition

A sequence  $(x_n)$  is said to be increasing if  $x_{n+1} \geq x_n$  for all  $n \in \mathbb{N}$ . Similarly,  $(x_n)$  is said to be decreasing if  $x_{n+1} \leq x_n$  for all  $n \in \mathbb{N}$ . We say that  $(x_n)$  is monotonic if it is either increasing or decreasing.

- 1 The sequence  $(\frac{1}{n})$  is decreasing.
- 2 The sequence  $(n + \frac{1}{n})$  is increasing.
- 3 The sequence  $(\cos \frac{n\pi}{3})$  is not monotonic.
- 4 The sequence  $((-1)^n)$  is not monotonic.

# Convergence of Monotone sequences

## Theorem

*If  $(x_n)$  is increasing and not bounded above then  $(x_n)$  diverges to  $\infty$ . If  $(x_n)$  is decreasing and not bounded below then  $(x_n)$  diverges to  $-\infty$ .*

# Convergence of Monotone sequences

## Theorem

*If  $(x_n)$  is increasing and not bounded above then  $(x_n)$  diverges to  $\infty$ . If  $(x_n)$  is decreasing and not bounded below then  $(x_n)$  diverges to  $-\infty$ .*

## Theorem (Monotone convergence theorem)

*Let  $(x_n)$  be a sequence of real numbers.*

- (a) If  $(x_n)$  is increasing and bounded above then  $(x_n)$  converges to  $\sup\{x_n : n \in \mathbb{N}\}$ .*
- (b) If  $(x_n)$  is decreasing and bounded below then  $(x_n)$  converges to  $\inf\{x_n : n \in \mathbb{N}\}$ .*
- (c) A monotonic sequence converges if and only if it is bounded.*

## Example

Let  $x_1 = 1$  and  $x_{n+1} = \frac{1}{3}(x_n + 1)$  for all  $n \in \mathbb{N}$ . Then the sequence  $(x_n)$  is convergent and  $\lim_{n \rightarrow \infty} x_n = \frac{1}{2}$ .

### Example

Let  $x_1 = 1$  and  $x_{n+1} = \frac{1}{3}(x_n + 1)$  for all  $n \in \mathbb{N}$ . Then the sequence  $(x_n)$  is convergent and  $\lim_{n \rightarrow \infty} x_n = \frac{1}{2}$ .

### Example

Let  $x_1 = 1$  and  $x_{n+1} = \frac{3}{x_n}$  for all  $n \geq 1$ . Then  $(x_n)$  diverges.

**Example:** The sequence  $((1 + 1/n)^n)$  is convergent.

**Example:** The sequence  $((1 + 1/n)^n)$  is convergent.

**Solution:** Let  $x_n = (1 + 1/n)^n$ . Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$



**Example:** The sequence  $((1 + 1/n)^n)$  is convergent.

**Solution:** Let  $x_n = (1 + 1/n)^n$ . Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$

$x_{n+1}$

$$\begin{aligned} &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n+1}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n-1}{n+1}\right) \\ &\quad + \frac{1}{(n+1)!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n}{n+1}\right). \end{aligned}$$

**Example:** The sequence  $((1 + 1/n)^n)$  is convergent.

**Solution:** Let  $x_n = (1 + 1/n)^n$ . Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$

$x_{n+1}$

$$\begin{aligned} &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n+1}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n-1}{n+1}\right) \\ &\quad + \frac{1}{(n+1)!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n}{n+1}\right). \end{aligned}$$

Therefore, we have

$$2 \leq x_1 < x_2 < \cdots < x_n < x_{n+1} < \cdots$$

**Example:** The sequence  $((1 + 1/n)^n)$  is convergent.

**Solution:** Let  $x_n = (1 + 1/n)^n$ . Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$

$x_{n+1}$

$$\begin{aligned} &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n+1}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n-1}{n+1}\right) \\ &\quad + \frac{1}{(n+1)!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n}{n+1}\right). \end{aligned}$$

Therefore, we have

$$2 \leq x_1 < x_2 < \cdots < x_n < x_{n+1} < \cdots$$

For  $n > 1$ , we have

$$2 < x_n < 1 + 1 + \left(\frac{1}{2} + \frac{1}{2^2} + \cdots + \frac{1}{2^{n-1}}\right) < 2 + \left(1 - \frac{1}{2^{n-1}}\right) < 3.$$

## Theorem

Let  $x_n \neq 0$  for all  $n \in \mathbb{N}$  and let  $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$  exist.

- 1 If  $L < 1$ , then  $x_n \rightarrow 0$ .
- 2 If  $L > 1$ , then  $(x_n)$  is divergent.

## Theorem

Let  $x_n \neq 0$  for all  $n \in \mathbb{N}$  and let  $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$  exist.

- ① If  $L < 1$ , then  $x_n \rightarrow 0$ .
- ② If  $L > 1$ , then  $(x_n)$  is divergent.

## Remark

If  $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right| = 1$ , then  $(x_n)$  may converge or diverge. For example, the sequence  $((-1)^n)$  diverges and  $L = 1$ . For any nonzero constant sequence,  $L = 1$  and constant sequences are convergent.

## Theorem

Let  $x_n \neq 0$  for all  $n \in \mathbb{N}$  and let  $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$  exist.

- ① If  $L < 1$ , then  $x_n \rightarrow 0$ .
- ② If  $L > 1$ , then  $(x_n)$  is divergent.

## Remark

If  $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right| = 1$ , then  $(x_n)$  may converge or diverge. For example, the sequence  $((-1)^n)$  diverges and  $L = 1$ . For any nonzero constant sequence,  $L = 1$  and constant sequences are convergent.

## Example

If  $\alpha \in \mathbb{R}$ , then the sequence  $\left(\frac{\alpha^n}{n!}\right)$  is convergent.

## Theorem

Let  $x_n \neq 0$  for all  $n \in \mathbb{N}$  and let  $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$  exist.

- ① If  $L < 1$ , then  $x_n \rightarrow 0$ .
- ② If  $L > 1$ , then  $(x_n)$  is divergent.

## Remark

If  $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right| = 1$ , then  $(x_n)$  may converge or diverge. For example, the sequence  $((-1)^n)$  diverges and  $L = 1$ . For any nonzero constant sequence,  $L = 1$  and constant sequences are convergent.

## Example

If  $\alpha \in \mathbb{R}$ , then the sequence  $\left(\frac{\alpha^n}{n!}\right)$  is convergent.

## Example

The sequence  $\left(\frac{2^n}{n^4}\right)$  is not convergent.

# Subsequence

## Definition (Subsequence)

Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . If  $(n_k)$  is a sequence of positive integers such that  $n_1 < n_2 < n_3 < \cdots$ , then  $(x_{n_k})$  is called a subsequence of  $(x_n)$ .



# Subsequence

## Definition (Subsequence)

Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . If  $(n_k)$  is a sequence of positive integers such that  $n_1 < n_2 < n_3 < \cdots$ , then  $(x_{n_k})$  is called a subsequence of  $(x_n)$ .

**Example:** Think of some divergent sequences and their convergent subsequences.

# Subsequence

## Definition (Subsequence)

Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . If  $(n_k)$  is a sequence of positive integers such that  $n_1 < n_2 < n_3 < \cdots$ , then  $(x_{n_k})$  is called a subsequence of  $(x_n)$ .

**Example:** Think of some divergent sequences and their convergent subsequences.

**Theorem:** If a sequence  $(x_n)$  converges to  $\ell$ , then every subsequence of  $(x_n)$  must converge to  $\ell$ .

# Subsequence

## Definition (Subsequence)

Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . If  $(n_k)$  is a sequence of positive integers such that  $n_1 < n_2 < n_3 < \cdots$ , then  $(x_{n_k})$  is called a subsequence of  $(x_n)$ .

**Example:** Think of some divergent sequences and their convergent subsequences.

**Theorem:** If a sequence  $(x_n)$  converges to  $\ell$ , then every subsequence of  $(x_n)$  must converge to  $\ell$ .

**Remark:** From the above theorem, we have the following:

- If  $(x_n)$  has a subsequence  $(x_{n_k})$  such that  $x_{n_k} \not\rightarrow \ell$ , then  $x_n \not\rightarrow \ell$ .
- If  $(x_n)$  has two subsequences converging to two different limits, then  $(x_n)$  cannot be convergent.

# Subsequence

Example: We have

# Subsequence

**Example:** We have

- 1 If  $x_n = (-1)^n(1 - \frac{1}{n})$  for all  $n \in \mathbb{N}$ , then  $(x_n)$  is not convergent.

# Subsequence

**Example:** We have

- ① If  $x_n = (-1)^n(1 - \frac{1}{n})$  for all  $n \in \mathbb{N}$ , then  $(x_n)$  is not convergent.
- ② Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . Then  $(x_{2n})$  and  $(x_{2n-1})$  are two subsequences of  $(x_n)$ . Suppose that  $x_{2n} \rightarrow \ell \in \mathbb{R}$  and  $x_{2n-1} \rightarrow \ell$ . Then  $x_n \rightarrow \ell$ .

# Subsequence

**Example:** We have

- 1 If  $x_n = (-1)^n(1 - \frac{1}{n})$  for all  $n \in \mathbb{N}$ , then  $(x_n)$  is not convergent.
- 2 Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . Then  $(x_{2n})$  and  $(x_{2n-1})$  are two subsequences of  $(x_n)$ . Suppose that  $x_{2n} \rightarrow \ell \in \mathbb{R}$  and  $x_{2n-1} \rightarrow \ell$ . Then  $x_n \rightarrow \ell$ .
- 3 The sequence  $(1, \frac{1}{2}, 1, \frac{2}{3}, 1, \frac{3}{4}, \dots)$  converges to 1.

# Subsequence

**Example:** We have

- ① If  $x_n = (-1)^n(1 - \frac{1}{n})$  for all  $n \in \mathbb{N}$ , then  $(x_n)$  is not convergent.
- ② Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . Then  $(x_{2n})$  and  $(x_{2n-1})$  are two subsequences of  $(x_n)$ . Suppose that  $x_{2n} \rightarrow \ell \in \mathbb{R}$  and  $x_{2n-1} \rightarrow \ell$ . Then  $x_n \rightarrow \ell$ .
- ③ The sequence  $(1, \frac{1}{2}, 1, \frac{2}{3}, 1, \frac{3}{4}, \dots)$  converges to 1.

## Theorem

*Every sequence of real numbers has a monotone subsequence.*



# Subsequence

**Example:** We have

- 1 If  $x_n = (-1)^n(1 - \frac{1}{n})$  for all  $n \in \mathbb{N}$ , then  $(x_n)$  is not convergent.
- 2 Let  $(x_n)$  be a sequence in  $\mathbb{R}$ . Then  $(x_{2n})$  and  $(x_{2n-1})$  are two subsequences of  $(x_n)$ . Suppose that  $x_{2n} \rightarrow \ell \in \mathbb{R}$  and  $x_{2n-1} \rightarrow \ell$ . Then  $x_n \rightarrow \ell$ .
- 3 The sequence  $(1, \frac{1}{2}, 1, \frac{2}{3}, 1, \frac{3}{4}, \dots)$  converges to 1.

## Theorem

*Every sequence of real numbers has a monotone subsequence.*

## Theorem (Bolzano-Weierstrass Theorem)

*Every bounded sequence in  $\mathbb{R}$  has a convergent subsequence.*

# Cauchy sequence

## Definition (Cauchy sequence)

A sequence  $(x_n)$  is called a Cauchy sequence if for each  $\varepsilon > 0$ , there exists  $n_0 \in \mathbb{N}$  such that  $|x_m - x_n| < \varepsilon$  for all  $m, n \geq n_0$ .

# Cauchy sequence

## Definition (Cauchy sequence)

A sequence  $(x_n)$  is called a Cauchy sequence if for each  $\varepsilon > 0$ , there exists  $n_0 \in \mathbb{N}$  such that  $|x_m - x_n| < \varepsilon$  for all  $m, n \geq n_0$ .

**Theorem:** Every Cauchy sequence is bounded.

# Cauchy sequence

## Definition (Cauchy sequence)

A sequence  $(x_n)$  is called a Cauchy sequence if for each  $\varepsilon > 0$ , there exists  $n_0 \in \mathbb{N}$  such that  $|x_m - x_n| < \varepsilon$  for all  $m, n \geq n_0$ .

**Theorem:** Every Cauchy sequence is bounded.

## Theorem (Cauchy's criterion for convergence)

*A sequence in  $\mathbb{R}$  is convergent if and only if it is a Cauchy sequence.*

# Cauchy sequence

## Theorem

Let  $(x_n)$  satisfy *either* of the following conditions:

- 1  $|x_{n+1} - x_n| \leq \alpha^n$  for all  $n \in \mathbb{N}$
- 2  $|x_{n+2} - x_{n+1}| \leq \alpha|x_{n+1} - x_n|$  for all  $n \in \mathbb{N}$ ,

where  $0 < \alpha < 1$ . Then  $(x_n)$  is a Cauchy sequence.

# Cauchy sequence

## Theorem

Let  $(x_n)$  satisfy *either* of the following conditions:

- ①  $|x_{n+1} - x_n| \leq \alpha^n$  for all  $n \in \mathbb{N}$
- ②  $|x_{n+2} - x_{n+1}| \leq \alpha|x_{n+1} - x_n|$  for all  $n \in \mathbb{N}$ ,

where  $0 < \alpha < 1$ . Then  $(x_n)$  is a Cauchy sequence.

## Proof of (1).

For all  $m, n \in \mathbb{N}$  with  $m > n$ , we have

$$\begin{aligned} |x_m - x_n| &\leq |x_n - x_{n+1}| + |x_{n+1} - x_{n+2}| + \cdots + |x_{m-1} - x_m| \\ &\leq \alpha^n + \alpha^{n+1} + \cdots + \alpha^{m-1} \\ &= \frac{\alpha^n}{1 - \alpha} (1 - \alpha^{m-n}) < \frac{\alpha^n}{1 - \alpha} \end{aligned}$$



# Cauchy sequence

**Proof of (2)** For all  $m, n \in \mathbb{N}$  with  $m > n$ , we have

$$\begin{aligned} |x_m - x_n| &\leq |x_n - x_{n+1}| + |x_{n+1} - x_{n+2}| + \cdots + |x_{m-1} - x_m| \\ &\leq (\alpha^{n-1} + \alpha^n + \cdots + \alpha^{m-2}) |x_2 - x_1| \\ &= \frac{\alpha^{n-1}}{1 - \alpha} (1 - \alpha^{m-n}) |x_2 - x_1| \leq \frac{\alpha^{n-1}}{1 - \alpha} |x_2 - x_1| \end{aligned}$$

# Cauchy sequence

**Proof of (2)** For all  $m, n \in \mathbb{N}$  with  $m > n$ , we have

$$\begin{aligned} |x_m - x_n| &\leq |x_n - x_{n+1}| + |x_{n+1} - x_{n+2}| + \cdots + |x_{m-1} - x_m| \\ &\leq (\alpha^{n-1} + \alpha^n + \cdots + \alpha^{m-2})|x_2 - x_1| \\ &= \frac{\alpha^{n-1}}{1 - \alpha}(1 - \alpha^{m-n})|x_2 - x_1| \leq \frac{\alpha^{n-1}}{1 - \alpha}|x_2 - x_1| \end{aligned}$$

**Example:** Let  $(x_n)$  be a sequence defined as  $x_1 = 1$  and  $x_{n+1} = 1 + \frac{1}{x_n}$  for  $n \in \mathbb{N}$ . Then  $x_{n+1}x_n = 1 + x_n > 2$ . Now,

$$|x_{n+2} - x_{n+1}| = \left| \frac{x_{n+1} - x_n}{x_{n+1}x_n} \right| < \frac{1}{2}|x_{n+1} - x_n|.$$

Hence,  $(x_n)$  is a Cauchy sequence.



## Limit superior

Let  $(x_n)$  be a bounded sequence. Let  $y_1 = \sup\{x_1, x_2, \dots\}$ ,  $y_2 = \sup\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

## Limit superior

Let  $(x_n)$  be a bounded sequence. Let  $y_1 = \sup\{x_1, x_2, \dots\}$ ,  $y_2 = \sup\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

## Limit superior

Let  $(x_n)$  be a bounded sequence. Let  $y_1 = \sup\{x_1, x_2, \dots\}$ ,  $y_2 = \sup\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

By Monotone convergence theorem,  $(y_n)$  is convergent and converges to the infimum of  $\{y_1, y_2, \dots\}$ .

## Limit superior

Let  $(x_n)$  be a bounded sequence. Let  $y_1 = \sup\{x_1, x_2, \dots\}$ ,  $y_2 = \sup\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

By Monotone convergence theorem,  $(y_n)$  is convergent and converges to the infimum of  $\{y_1, y_2, \dots\}$ .

The limit of the sequence  $(y_n)$  is called the limit superior of the sequence  $(x_n)$ , and is denoted by  $\limsup x_n$ .

## Limit superior

Let  $(x_n)$  be a bounded sequence. Let  $y_1 = \sup\{x_1, x_2, \dots\}$ ,  $y_2 = \sup\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

By Monotone convergence theorem,  $(y_n)$  is convergent and converges to the infimum of  $\{y_1, y_2, \dots\}$ .

The limit of the sequence  $(y_n)$  is called the limit superior of the sequence  $(x_n)$ , and is denoted by  $\limsup x_n$ .

Thus,

$$\limsup x_n := \lim_{n \rightarrow \infty} y_n = \inf\{y_1, y_2, \dots\} = \inf_n \sup_{k \geq n} x_k.$$

## Limit inferior

Similarly, let  $z_1 = \inf\{x_1, x_2, \dots\}$ ,  $z_2 = \inf\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

## Limit inferior

Similarly, let  $z_1 = \inf\{x_1, x_2, \dots\}$ ,  $z_2 = \inf\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

## Limit inferior

Similarly, let  $z_1 = \inf\{x_1, x_2, \dots\}$ ,  $z_2 = \inf\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

By Monotone convergence theorem,  $(z_n)$  is convergent and converges to the supremum of  $\{z_1, z_2, \dots\}$ .



## Limit inferior

Similarly, let  $z_1 = \inf\{x_1, x_2, \dots\}$ ,  $z_2 = \inf\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

By Monotone convergence theorem,  $(z_n)$  is convergent and converges to the supremum of  $\{z_1, z_2, \dots\}$ .

The limit of the sequence  $(z_n)$  is called the limit inferior of the sequence  $(x_n)$ , and is denoted by  $\liminf x_n$ .

## Limit inferior

Similarly, let  $z_1 = \inf\{x_1, x_2, \dots\}$ ,  $z_2 = \inf\{x_2, x_3, \dots\}$ , and so on. That is, for  $n \in \mathbb{N}$ ,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

By Monotone convergence theorem,  $(z_n)$  is convergent and converges to the supremum of  $\{z_1, z_2, \dots\}$ .

The limit of the sequence  $(z_n)$  is called the limit inferior of the sequence  $(x_n)$ , and is denoted by  $\liminf x_n$ .

Thus,

$$\liminf x_n := \lim_{n \rightarrow \infty} z_n = \sup\{z_1, z_2, \dots\} = \sup_n \inf_{k \geq n} x_k.$$

**Remark:** Suppose that  $|x_n| < M$  for  $n \in \mathbb{N}$ . Then  $-M \leq z_n \leq y_n \leq M$  for all  $n$ . Hence,

$$-M \leq \liminf x_n \leq \limsup x_n \leq M.$$

**Remark:** Suppose that  $|x_n| < M$  for  $n \in \mathbb{N}$ . Then  $-M \leq z_n \leq y_n \leq M$  for all  $n$ . Hence,

$$-M \leq \liminf x_n \leq \limsup x_n \leq M.$$

**Example:** Consider the sequence  $(x_n)$ , where  $x_n = (-1)^n$ . Clearly, for any  $n$ ,  $y_n = \sup\{x_n, x_{n+1}, \dots\} = 1$  and  $z_n = \inf\{x_n, x_{n+1}, \dots\} = -1$ . Hence,

$$\limsup x_n = 1 \quad \text{and} \quad \liminf x_n = -1.$$

**Remark:** Suppose that  $|x_n| < M$  for  $n \in \mathbb{N}$ . Then  $-M \leq z_n \leq y_n \leq M$  for all  $n$ . Hence,

$$-M \leq \liminf x_n \leq \limsup x_n \leq M.$$

**Example:** Consider the sequence  $(x_n)$ , where  $x_n = (-1)^n$ . Clearly, for any  $n$ ,  $y_n = \sup\{x_n, x_{n+1}, \dots\} = 1$  and  $z_n = \inf\{x_n, x_{n+1}, \dots\} = -1$ . Hence,

$$\limsup x_n = 1 \quad \text{and} \quad \liminf x_n = -1.$$

**Example:** Consider the sequence  $(x_n)$ , where  $x_n = \frac{1}{n}$ .

Clearly, for any  $n$ ,  $y_n = \sup\{\frac{1}{k} : k \geq n\} = \frac{1}{n}$  and

$$z_n = \inf\{\frac{1}{k} : k \geq n\} = 0.$$

Hence,  $\limsup x_n = \lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} \frac{1}{n} = 0$  and  $\liminf x_n = 0$ .

## Theorem

*Let  $(a_n)$  and  $(b_n)$  be two bounded sequences.*

- ①  $\liminf a_n \leq \limsup a_n$ .
- ② *If  $a_n \leq b_n$  for all  $n \in \mathbb{N}$ , then  $\limsup a_n \leq \limsup b_n$  and  $\liminf a_n \leq \liminf b_n$ .*
- ③  $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$  and  $\liminf(a_n + b_n) \geq \liminf a_n + \liminf b_n$ .

## Theorem

Let  $(a_n)$  and  $(b_n)$  be two bounded sequences.

- ①  $\liminf a_n \leq \limsup a_n$ .
- ② If  $a_n \leq b_n$  for all  $n \in \mathbb{N}$ , then  $\limsup a_n \leq \limsup b_n$  and  $\liminf a_n \leq \liminf b_n$ .
- ③  $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$  and  $\liminf(a_n + b_n) \geq \liminf a_n + \liminf b_n$ .

## Theorem

If  $(a_n)$  is a convergent sequence, then

$$\liminf a_n = \lim_{n \rightarrow \infty} a_n = \limsup a_n.$$

## Theorem

Let  $(a_n)$  and  $(b_n)$  be two bounded sequences.

- ①  $\liminf a_n \leq \limsup a_n$ .
- ② If  $a_n \leq b_n$  for all  $n \in \mathbb{N}$ , then  $\limsup a_n \leq \limsup b_n$  and  $\liminf a_n \leq \liminf b_n$ .
- ③  $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$  and  $\liminf(a_n + b_n) \geq \liminf a_n + \liminf b_n$ .

## Theorem

If  $(a_n)$  is a convergent sequence, then

$$\liminf a_n = \lim_{n \rightarrow \infty} a_n = \limsup a_n.$$

## Theorem

Let  $(a_n)$  be a bounded sequence. If  $\limsup a_n = \liminf a_n$ , then  $(a_n)$  is convergent and  $\lim_{n \rightarrow \infty} a_n = \limsup a_n$ .