

MA 101 (Mathematics-I)

Introduction

Consider the “sequence” of rationals

1.4, 1.41, 1.414, 1.4142, 1.41421, ...

Introduction

Consider the “sequence” of rationals

$$1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$$

Their squares are:

$$1.96, 1.9881, 1.99396, 1.99996164, 1.9999899241, \dots$$

Introduction

Consider the “sequence” of rationals

$$1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$$

Their squares are:

$$1.96, 1.9881, 1.99396, 1.99996164, 1.9999899241, \dots$$

It seems that the above numbers are getting closer and closer to 2.

Introduction

Consider the “sequence” of rationals

$$1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$$

Their squares are:

$$1.96, 1.9881, 1.99396, 1.99996164, 1.9999899241, \dots$$

It seems that the above numbers are getting closer and closer to 2.

Therefore, the rationals $1.4, 1.41, 1.414, 1.4142, 1.41421, \dots$ are getting closer and closer to $\sqrt{2}$.

Definition (Sequence)

A sequence of real numbers or a sequence in \mathbb{R} is a mapping $f : \mathbb{N} \rightarrow \mathbb{R}$. We write x_n for $f(n)$, $n \in \mathbb{N}$ and it is customary to denote a sequence as $\langle x_n \rangle$ or (x_n) or $\{x_n\}$.

Definition (Sequence)

A sequence of real numbers or a sequence in \mathbb{R} is a mapping $f : \mathbb{N} \rightarrow \mathbb{R}$. We write x_n for $f(n)$, $n \in \mathbb{N}$ and it is customary to denote a sequence as $\langle x_n \rangle$ or (x_n) or $\{x_n\}$.

Example

There are different ways of expressing a sequence. For example:

- ① *Constant sequence: (a, a, a, \dots) , where $a \in \mathbb{R}$*
- ② *Sequence defined by listing: $(1, 4, 8, 11, 52, \dots)$*
- ③ *Sequence defined by rule: (x_n) , where $x_n = 3n^2$ for all $n \in \mathbb{N}$*
- ④ *Sequence defined recursively: (x_n) , where $x_1 = 4$ and $x_{n+1} = 2x_n - 5$ for all $n \in \mathbb{N}$*

Convergence: What does it mean?

Think of the examples:

① $(2, 2, 2, \dots)$

② $(\frac{1}{n})$

③ $((-1)^n \frac{1}{n})$

④ $(1, 2, 1, 2, \dots)$

⑤ (\sqrt{n})

⑥ $((-1)^n(1 - \frac{1}{n}))$

Definition (Convergent sequence)

A sequence (x_n) is said to be convergent if there exists $\ell \in \mathbb{R}$ such that for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ satisfying $|x_n - \ell| < \varepsilon$ for all $n \geq n_0$. We say that ℓ is a limit of (x_n) .

Definition (Convergent sequence)

A sequence (x_n) is said to be convergent if there exists $\ell \in \mathbb{R}$ such that for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ satisfying $|x_n - \ell| < \varepsilon$ for all $n \geq n_0$. We say that ℓ is a limit of (x_n) .

Notation: We write $\lim_{n \rightarrow \infty} x_n = \ell$ or $x_n \rightarrow \ell$.

Definition (Convergent sequence)

A sequence (x_n) is said to be convergent if there exists $\ell \in \mathbb{R}$ such that for every $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ satisfying $|x_n - \ell| < \varepsilon$ for all $n \geq n_0$. We say that ℓ is a limit of (x_n) .

Notation: We write $\lim_{n \rightarrow \infty} x_n = \ell$ or $x_n \rightarrow \ell$.

Theorem

Limit of a convergent sequence is unique.

Example

Using the definition of convergence of a sequence, show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

Example

Using the definition of convergence of a sequence, show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

Example

Consider the sequence (x_n) where $x_n = (-1)^n$. The terms of the sequence are $-1, 1, -1, 1, -1, 1, \dots$. It is intuitively clear that this sequence does not approach to any real number. Therefore, the sequence does not converge. We now establish this fact by using the definition.

Example

Using the definition of convergence of a sequence, show that

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

Example

Consider the sequence (x_n) where $x_n = (-1)^n$. The terms of the sequence are $-1, 1, -1, 1, -1, 1, \dots$. It is intuitively clear that this sequence does not approach to any real number. Therefore, the sequence does not converge. We now establish this fact by using the definition.

Exercise (Tutorial problem)

If $|\alpha| < 1$, then the sequence (α^n) converges to 0.

Bounded sequence: Given a sequence (x_n) , we can ask whether the set $\{x_1, x_2, x_3, \dots\}$ is bounded or not. If this set is bounded then we call that the sequence (x_n) is bounded. Equivalently, the sequence (x_n) is bounded if there is a positive number M such that $|x_n| \leq M$ for all $n \in \mathbb{N}$. If (x_n) is not bounded then it is said to be unbounded.

Bounded sequence: Given a sequence (x_n) , we can ask whether the set $\{x_1, x_2, x_3, \dots\}$ is bounded or not. If this set is bounded then we call that the sequence (x_n) is bounded. Equivalently, the sequence (x_n) is bounded if there is a positive number M such that $|x_n| \leq M$ for all $n \in \mathbb{N}$. If (x_n) is not bounded then it is said to be unbounded.

Theorem

Every convergent sequence is bounded.

Bounded sequence: Given a sequence (x_n) , we can ask whether the set $\{x_1, x_2, x_3, \dots\}$ is bounded or not. If this set is bounded then we call that the sequence (x_n) is bounded. Equivalently, the sequence (x_n) is bounded if there is a positive number M such that $|x_n| \leq M$ for all $n \in \mathbb{N}$. If (x_n) is not bounded then it is said to be unbounded.

Theorem

Every convergent sequence is bounded.

Remark

- *From the above theorem, it follows that if a sequence is not bounded then it is not convergent. For example, the sequence (\sqrt{n}) is unbounded and hence is not convergent.*
- *Every bounded sequence is not convergent. For example, $((-1)^n)$ is a bounded sequence but it does not converge.*

Limit rules for convergent sequences

Theorem

Let $x_n \rightarrow x$ and $y_n \rightarrow y$. Then

- (a) $x_n + y_n \rightarrow x + y$.
- (b) $\alpha x_n \rightarrow \alpha x$ for all $\alpha \in \mathbb{R}$.
- (c) $|x_n| \rightarrow |x|$.
- (d) $x_n y_n \rightarrow xy$.
- (e) $\frac{x_n}{y_n} \rightarrow \frac{x}{y}$ if $y_n \neq 0$ for all $n \in \mathbb{N}$ and $y \neq 0$.

Limit rules for convergent sequences

Theorem

Let $x_n \rightarrow x$ and $y_n \rightarrow y$. Then

- (a) $x_n + y_n \rightarrow x + y$.
- (b) $\alpha x_n \rightarrow \alpha x$ for all $\alpha \in \mathbb{R}$.
- (c) $|x_n| \rightarrow |x|$.
- (d) $x_n y_n \rightarrow xy$.
- (e) $\frac{x_n}{y_n} \rightarrow \frac{x}{y}$ if $y_n \neq 0$ for all $n \in \mathbb{N}$ and $y \neq 0$.

Example

The sequence $\left(\frac{2n^2-3n}{3n^2+5n+3}\right)$ is convergent with limit $\frac{2}{3}$.

Limit rules for convergent sequences

Theorem

Let $x_n \rightarrow x$ and $y_n \rightarrow y$. Then

- (a) $x_n + y_n \rightarrow x + y$.
- (b) $\alpha x_n \rightarrow \alpha x$ for all $\alpha \in \mathbb{R}$.
- (c) $|x_n| \rightarrow |x|$.
- (d) $x_n y_n \rightarrow xy$.
- (e) $\frac{x_n}{y_n} \rightarrow \frac{x}{y}$ if $y_n \neq 0$ for all $n \in \mathbb{N}$ and $y \neq 0$.

Example

The sequence $\left(\frac{2n^2-3n}{3n^2+5n+3}\right)$ is convergent with limit $\frac{2}{3}$.

Example

The sequence $(\sqrt{n+1} - \sqrt{n})$ is convergent with limit 0.

Sandwich theorem

Theorem (Sandwich theorem)

Let (x_n) , (y_n) , (z_n) be sequences such that $x_n \leq y_n \leq z_n$ for all $n \geq n_0$ for some $n_0 \in \mathbb{N}$. If both (x_n) and (z_n) converge to the same limit ℓ , then (y_n) also converges to ℓ .

Sandwich theorem

Theorem (Sandwich theorem)

Let (x_n) , (y_n) , (z_n) be sequences such that $x_n \leq y_n \leq z_n$ for all $n \geq n_0$ for some $n_0 \in \mathbb{N}$. If both (x_n) and (z_n) converge to the same limit ℓ , then (y_n) also converges to ℓ .

Example

$$\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$$

Sandwich theorem

Theorem (Sandwich theorem)

Let (x_n) , (y_n) , (z_n) be sequences such that $x_n \leq y_n \leq z_n$ for all $n \geq n_0$ for some $n_0 \in \mathbb{N}$. If both (x_n) and (z_n) converge to the same limit ℓ , then (y_n) also converges to ℓ .

Example

$$\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$$

Example

If $p > 0$, then $\frac{1}{n^p} \rightarrow 0$.

Sandwich theorem

Theorem (Sandwich theorem)

Let (x_n) , (y_n) , (z_n) be sequences such that $x_n \leq y_n \leq z_n$ for all $n \geq n_0$ for some $n_0 \in \mathbb{N}$. If both (x_n) and (z_n) converge to the same limit ℓ , then (y_n) also converges to ℓ .

Example

$$\lim_{n \rightarrow \infty} \frac{\cos n}{n} = 0.$$

Example

If $p > 0$, then $\frac{1}{n^p} \rightarrow 0$.

Example

If $\alpha > 0$, then the sequence $(\alpha^{\frac{1}{n}})$ converges to 1.

Example

The sequence $(n^{\frac{1}{n}})$ converges to 1.

Example

The sequence $(n^{\frac{1}{n}})$ converges to 1.

Example

If $p > 0$ and $\alpha \in \mathbb{R}$, then $\frac{n^\alpha}{(1+p)^n} \rightarrow 0$.

Example

The sequence $(n^{\frac{1}{n}})$ converges to 1.

Example

If $p > 0$ and $\alpha \in \mathbb{R}$, then $\frac{n^\alpha}{(1+p)^n} \rightarrow 0$.

Theorem

Let $r \in \mathbb{R}$. Then there exists a sequence (x_n) of rational numbers such that $\lim_{n \rightarrow \infty} x_n = r$.

Divergent sequences

A sequence (x_n) is said to be **divergent** if it has no limit.

Divergent sequences

A sequence (x_n) is said to be **divergent** if it has no limit.

Example

- *If (x_n) is unbounded then it is divergent.*
- *(\sqrt{n}) , $(3n^2)$, $((-1)^n n^3)$ are all divergent.*

Example

The sequence $((-1)^n)$ is not convergent, and so it is a divergent sequence although it is bounded.

Definition

A sequence (x_n) is said to approach infinity or diverges to infinity if for any real number $M > 0$, there is a positive integer n_0 such that $a_n \geq M$ for all $n \geq n_0$. Similarly, (x_n) is said to approach $-\infty$ or diverges to $-\infty$ if for any real number $M > 0$, there is a positive integer n_0 such that $a_n \leq -M$ for all $n \geq n_0$.

Monotone sequence

Definition

A sequence (x_n) is said to be increasing if $x_{n+1} \geq x_n$ for all $n \in \mathbb{N}$. Similarly, (x_n) is said to be decreasing if $x_{n+1} \leq x_n$ for all $n \in \mathbb{N}$. We say that (x_n) is monotonic if it is either increasing or decreasing.

Monotone sequence

Definition

A sequence (x_n) is said to be increasing if $x_{n+1} \geq x_n$ for all $n \in \mathbb{N}$. Similarly, (x_n) is said to be decreasing if $x_{n+1} \leq x_n$ for all $n \in \mathbb{N}$. We say that (x_n) is monotonic if it is either increasing or decreasing.

- 1 The sequence $(\frac{1}{n})$ is decreasing.

Monotone sequence

Definition

A sequence (x_n) is said to be increasing if $x_{n+1} \geq x_n$ for all $n \in \mathbb{N}$. Similarly, (x_n) is said to be decreasing if $x_{n+1} \leq x_n$ for all $n \in \mathbb{N}$. We say that (x_n) is monotonic if it is either increasing or decreasing.

- 1 The sequence $(\frac{1}{n})$ is decreasing.
- 2 The sequence $(n + \frac{1}{n})$ is increasing.

Monotone sequence

Definition

A sequence (x_n) is said to be increasing if $x_{n+1} \geq x_n$ for all $n \in \mathbb{N}$. Similarly, (x_n) is said to be decreasing if $x_{n+1} \leq x_n$ for all $n \in \mathbb{N}$. We say that (x_n) is monotonic if it is either increasing or decreasing.

- 1 The sequence $(\frac{1}{n})$ is decreasing.
- 2 The sequence $(n + \frac{1}{n})$ is increasing.
- 3 The sequence $(\cos \frac{n\pi}{3})$ is not monotonic.

Monotone sequence

Definition

A sequence (x_n) is said to be increasing if $x_{n+1} \geq x_n$ for all $n \in \mathbb{N}$. Similarly, (x_n) is said to be decreasing if $x_{n+1} \leq x_n$ for all $n \in \mathbb{N}$. We say that (x_n) is monotonic if it is either increasing or decreasing.

- 1 The sequence $(\frac{1}{n})$ is decreasing.
- 2 The sequence $(n + \frac{1}{n})$ is increasing.
- 3 The sequence $(\cos \frac{n\pi}{3})$ is not monotonic.
- 4 The sequence $((-1)^n)$ is not monotonic.

Convergence of Monotone sequences

Theorem

If (x_n) is increasing and not bounded above then (x_n) diverges to ∞ . If (x_n) is decreasing and not bounded below then (x_n) diverges to $-\infty$.

Convergence of Monotone sequences

Theorem

If (x_n) is increasing and not bounded above then (x_n) diverges to ∞ . If (x_n) is decreasing and not bounded below then (x_n) diverges to $-\infty$.

Theorem (Monotone convergence theorem)

Let (x_n) be a sequence of real numbers.

- (a) If (x_n) is increasing and bounded above then (x_n) converges to $\sup\{x_n : n \in \mathbb{N}\}$.*
- (b) If (x_n) is decreasing and bounded below then (x_n) converges to $\inf\{x_n : n \in \mathbb{N}\}$.*
- (c) A monotonic sequence converges if and only if it is bounded.*

Example

Let $x_1 = 1$ and $x_{n+1} = \frac{1}{3}(x_n + 1)$ for all $n \in \mathbb{N}$. Then the sequence (x_n) is convergent and $\lim_{n \rightarrow \infty} x_n = \frac{1}{2}$.

Example

Let $x_1 = 1$ and $x_{n+1} = \frac{1}{3}(x_n + 1)$ for all $n \in \mathbb{N}$. Then the sequence (x_n) is convergent and $\lim_{n \rightarrow \infty} x_n = \frac{1}{2}$.

Example

Let $x_1 = 1$ and $x_{n+1} = \frac{3}{x_n}$ for all $n \geq 1$. Then (x_n) diverges.

Example: The sequence $((1 + 1/n)^n)$ is convergent.

Example: The sequence $((1 + 1/n)^n)$ is convergent.

Solution: Let $a_n = (1 + 1/n)^n$. Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$

Example: The sequence $((1 + 1/n)^n)$ is convergent.

Solution: Let $a_n = (1 + 1/n)^n$. Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$

x_{n+1}

$$\begin{aligned} &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n+1}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n-1}{n+1}\right) \\ &\quad + \frac{1}{(n+1)!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n}{n+1}\right). \end{aligned}$$

Example: The sequence $((1 + 1/n)^n)$ is convergent.

Solution: Let $a_n = (1 + 1/n)^n$. Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$

x_{n+1}

$$\begin{aligned} &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n+1}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n-1}{n+1}\right) \\ &\quad + \frac{1}{(n+1)!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n}{n+1}\right). \end{aligned}$$

Therefore, we have

$$2 \leq x_1 < x_2 < \cdots < x_n < x_{n+1} < \cdots$$

Example: The sequence $((1 + 1/n)^n)$ is convergent.

Solution: Let $a_n = (1 + 1/n)^n$. Then

$$\begin{aligned} x_n &= \sum_{k=0}^n \binom{n}{k} \left(\frac{1}{n}\right)^k = 1 + \frac{n}{1} \cdot \frac{1}{n} + \frac{n(n-1)}{2!} \cdot \frac{1}{n^2} + \cdots + \frac{n(n-1) \cdots 2 \cdot 1}{n!} \cdot \frac{1}{n^n} \\ &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n}\right) \left(1 - \frac{2}{n}\right) \cdots \left(1 - \frac{n-1}{n}\right). \end{aligned}$$

x_{n+1}

$$\begin{aligned} &= 1 + 1 + \frac{1}{2!} \left(1 - \frac{1}{n+1}\right) + \cdots + \frac{1}{n!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n-1}{n+1}\right) \\ &\quad + \frac{1}{(n+1)!} \left(1 - \frac{1}{n+1}\right) \left(1 - \frac{2}{n+1}\right) \cdots \left(1 - \frac{n}{n+1}\right). \end{aligned}$$

Therefore, we have

$$2 \leq x_1 < x_2 < \cdots < x_n < x_{n+1} < \cdots$$

For $n > 1$, we have

$$2 < x_n < 1 + 1 + \left(\frac{1}{2} + \frac{1}{2^2} + \cdots + \frac{1}{2^{n-1}}\right) < 2 + \left(1 - \frac{1}{2^{n-1}}\right) < 3.$$

Theorem

Let $x_n \neq 0$ for all $n \in \mathbb{N}$ and let $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$ exist.

- 1 If $L < 1$, then $x_n \rightarrow 0$.
- 2 If $L > 1$, then (x_n) is divergent.

Theorem

Let $x_n \neq 0$ for all $n \in \mathbb{N}$ and let $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$ exist.

- ① If $L < 1$, then $x_n \rightarrow 0$.
- ② If $L > 1$, then (x_n) is divergent.

Remark

If $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right| = 1$, then (x_n) may converge or diverge. For example, the sequence $((-1)^n)$ diverges and $L = 1$. For any nonzero constant sequence, $L = 1$ and constant sequences are convergent.

Theorem

Let $x_n \neq 0$ for all $n \in \mathbb{N}$ and let $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$ exist.

- ① If $L < 1$, then $x_n \rightarrow 0$.
- ② If $L > 1$, then (x_n) is divergent.

Remark

If $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right| = 1$, then (x_n) may converge or diverge. For example, the sequence $((-1)^n)$ diverges and $L = 1$. For any nonzero constant sequence, $L = 1$ and constant sequences are convergent.

Example

If $\alpha \in \mathbb{R}$, then the sequence $\left(\frac{\alpha^n}{n!}\right)$ is convergent.

Theorem

Let $x_n \neq 0$ for all $n \in \mathbb{N}$ and let $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right|$ exist.

- ① If $L < 1$, then $x_n \rightarrow 0$.
- ② If $L > 1$, then (x_n) is divergent.

Remark

If $L = \lim_{n \rightarrow \infty} \left| \frac{x_{n+1}}{x_n} \right| = 1$, then (x_n) may converge or diverge. For example, the sequence $((-1)^n)$ diverges and $L = 1$. For any nonzero constant sequence, $L = 1$ and constant sequences are convergent.

Example

If $\alpha \in \mathbb{R}$, then the sequence $\left(\frac{\alpha^n}{n!}\right)$ is convergent.

Example

The sequence $\left(\frac{2^n}{n^4}\right)$ is not convergent.

Subsequence

Definition (Subsequence)

Let (x_n) be a sequence in \mathbb{R} . If (n_k) is a sequence of positive integers such that $n_1 < n_2 < n_3 < \cdots$, then (x_{n_k}) is called a subsequence of (x_n) .

Subsequence

Definition (Subsequence)

Let (x_n) be a sequence in \mathbb{R} . If (n_k) is a sequence of positive integers such that $n_1 < n_2 < n_3 < \cdots$, then (x_{n_k}) is called a subsequence of (x_n) .

Example: Think of some divergent sequences and their convergent subsequences.

Subsequence

Definition (Subsequence)

Let (x_n) be a sequence in \mathbb{R} . If (n_k) is a sequence of positive integers such that $n_1 < n_2 < n_3 < \cdots$, then (x_{n_k}) is called a subsequence of (x_n) .

Example: Think of some divergent sequences and their convergent subsequences.

Theorem: If a sequence (x_n) converges to ℓ , then every subsequence of (x_n) must converge to ℓ .

Subsequence

Definition (Subsequence)

Let (x_n) be a sequence in \mathbb{R} . If (n_k) is a sequence of positive integers such that $n_1 < n_2 < n_3 < \cdots$, then (x_{n_k}) is called a subsequence of (x_n) .

Example: Think of some divergent sequences and their convergent subsequences.

Theorem: If a sequence (x_n) converges to ℓ , then every subsequence of (x_n) must converge to ℓ .

Remark: From the above theorem, we have the following:

- If (x_n) has a subsequence (x_{n_k}) such that $x_{n_k} \not\rightarrow \ell$, then $x_n \not\rightarrow \ell$.
- If (x_n) has two subsequences converging to two different limits, then (x_n) cannot be convergent.

Subsequence

Example: We have

Subsequence

Example: We have

- 1 If $x_n = (-1)^n(1 - \frac{1}{n})$ for all $n \in \mathbb{N}$, then (x_n) is not convergent.

Subsequence

Example: We have

- ① If $x_n = (-1)^n(1 - \frac{1}{n})$ for all $n \in \mathbb{N}$, then (x_n) is not convergent.
- ② Let (x_n) be a sequence in \mathbb{R} . Then (x_{2n}) and (x_{2n-1}) are two subsequences of (x_n) . Suppose that $x_{2n} \rightarrow \ell \in \mathbb{R}$ and $x_{2n-1} \rightarrow \ell$. Then $x_n \rightarrow \ell$.

Subsequence

Example: We have

- 1 If $x_n = (-1)^n(1 - \frac{1}{n})$ for all $n \in \mathbb{N}$, then (x_n) is not convergent.
- 2 Let (x_n) be a sequence in \mathbb{R} . Then (x_{2n}) and (x_{2n-1}) are two subsequences of (x_n) . Suppose that $x_{2n} \rightarrow \ell \in \mathbb{R}$ and $x_{2n-1} \rightarrow \ell$. Then $x_n \rightarrow \ell$.
- 3 The sequence $(1, \frac{1}{2}, 1, \frac{2}{3}, 1, \frac{3}{4}, \dots)$ converges to 1.

Subsequence

Example: We have

- 1 If $x_n = (-1)^n(1 - \frac{1}{n})$ for all $n \in \mathbb{N}$, then (x_n) is not convergent.
- 2 Let (x_n) be a sequence in \mathbb{R} . Then (x_{2n}) and (x_{2n-1}) are two subsequences of (x_n) . Suppose that $x_{2n} \rightarrow \ell \in \mathbb{R}$ and $x_{2n-1} \rightarrow \ell$. Then $x_n \rightarrow \ell$.
- 3 The sequence $(1, \frac{1}{2}, 1, \frac{2}{3}, 1, \frac{3}{4}, \dots)$ converges to 1.

Theorem

Every sequence of real numbers has a monotone subsequence.

Subsequence

Example: We have

- 1 If $x_n = (-1)^n(1 - \frac{1}{n})$ for all $n \in \mathbb{N}$, then (x_n) is not convergent.
- 2 Let (x_n) be a sequence in \mathbb{R} . Then (x_{2n}) and (x_{2n-1}) are two subsequences of (x_n) . Suppose that $x_{2n} \rightarrow \ell \in \mathbb{R}$ and $x_{2n-1} \rightarrow \ell$. Then $x_n \rightarrow \ell$.
- 3 The sequence $(1, \frac{1}{2}, 1, \frac{2}{3}, 1, \frac{3}{4}, \dots)$ converges to 1.

Theorem

Every sequence of real numbers has a monotone subsequence.

Theorem (Bolzano-Weierstrass Theorem)

Every bounded sequence in \mathbb{R} has a convergent subsequence.

Cauchy sequence

Definition (Cauchy sequence)

A sequence (x_n) is called a Cauchy sequence if for each $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that $|x_m - x_n| < \varepsilon$ for all $m, n \geq n_0$.

Cauchy sequence

Definition (Cauchy sequence)

A sequence (x_n) is called a Cauchy sequence if for each $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that $|x_m - x_n| < \varepsilon$ for all $m, n \geq n_0$.

Theorem: Every Cauchy sequence is bounded.

Cauchy sequence

Definition (Cauchy sequence)

A sequence (x_n) is called a Cauchy sequence if for each $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that $|x_m - x_n| < \varepsilon$ for all $m, n \geq n_0$.

Theorem: Every Cauchy sequence is bounded.

Theorem (Cauchy's criterion for convergence)

A sequence in \mathbb{R} is convergent if and only if it is a Cauchy sequence.

Cauchy sequence

Theorem

Let (x_n) satisfy *either* of the following conditions:

- 1 $|x_{n+1} - x_n| \leq \alpha^n$ for all $n \in \mathbb{N}$
- 2 $|x_{n+2} - x_{n+1}| \leq \alpha|x_{n+1} - x_n|$ for all $n \in \mathbb{N}$,

where $0 < \alpha < 1$. Then (x_n) is a Cauchy sequence.

Cauchy sequence

Theorem

Let (x_n) satisfy *either* of the following conditions:

- ① $|x_{n+1} - x_n| \leq \alpha^n$ for all $n \in \mathbb{N}$
- ② $|x_{n+2} - x_{n+1}| \leq \alpha |x_{n+1} - x_n|$ for all $n \in \mathbb{N}$,

where $0 < \alpha < 1$. Then (x_n) is a Cauchy sequence.

Proof of (1).

For all $m, n \in \mathbb{N}$ with $m > n$, we have

$$\begin{aligned} |x_m - x_n| &\leq |x_n - x_{n+1}| + |x_{n+1} - x_{n+2}| + \cdots + |x_{m-1} - x_m| \\ &\leq \alpha^n + \alpha^{n+1} + \cdots + \alpha^{m-1} \\ &= \frac{\alpha^n}{1 - \alpha} (1 - \alpha^{m-n}) < \frac{\alpha^n}{1 - \alpha} \end{aligned}$$



Cauchy sequence

Proof of (2) For all $m, n \in \mathbb{N}$ with $m > n$, we have

$$\begin{aligned} |x_m - x_n| &\leq |x_n - x_{n+1}| + |x_{n+1} - x_{n+2}| + \cdots + |x_{m-1} - x_m| \\ &\leq (\alpha^{n-1} + \alpha^n + \cdots + \alpha^{m-2}) |x_2 - x_1| \\ &= \frac{\alpha^{n-1}}{1 - \alpha} (1 - \alpha^{m-n}) |x_2 - x_1| \leq \frac{\alpha^{n-1}}{1 - \alpha} |x_2 - x_1| \end{aligned}$$

Cauchy sequence

Proof of (2) For all $m, n \in \mathbb{N}$ with $m > n$, we have

$$\begin{aligned} |x_m - x_n| &\leq |x_n - x_{n+1}| + |x_{n+1} - x_{n+2}| + \cdots + |x_{m-1} - x_m| \\ &\leq (\alpha^{n-1} + \alpha^n + \cdots + \alpha^{m-2})|x_2 - x_1| \\ &= \frac{\alpha^{n-1}}{1 - \alpha}(1 - \alpha^{m-n})|x_2 - x_1| \leq \frac{\alpha^{n-1}}{1 - \alpha}|x_2 - x_1| \end{aligned}$$

Example: Let (x_n) be a sequence defined as $x_1 = 1$ and $x_{n+1} = 1 + \frac{1}{x_n}$ for $n \in \mathbb{N}$. Then $x_{n+1}x_n = 1 + x_n > 2$. Now,

$$|x_{n+2} - x_{n+1}| = \left| \frac{x_{n+1} - x_n}{x_{n+1}x_n} \right| < \frac{1}{2}|x_{n+1} - x_n|.$$

Hence, (x_n) is a Cauchy sequence.

Limit superior

Let (x_n) be a bounded sequence. Let $y_1 = \sup\{x_1, x_2, \dots\}$, $y_2 = \sup\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

Limit superior

Let (x_n) be a bounded sequence. Let $y_1 = \sup\{x_1, x_2, \dots\}$, $y_2 = \sup\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

Limit superior

Let (x_n) be a bounded sequence. Let $y_1 = \sup\{x_1, x_2, \dots\}$, $y_2 = \sup\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

By Monotone convergence theorem, (y_n) is convergent and converges to the infimum of $\{y_1, y_2, \dots\}$.

Limit superior

Let (x_n) be a bounded sequence. Let $y_1 = \sup\{x_1, x_2, \dots\}$, $y_2 = \sup\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

By Monotone convergence theorem, (y_n) is convergent and converges to the infimum of $\{y_1, y_2, \dots\}$.

The limit of the sequence (y_n) is called the limit superior of the sequence (x_n) , and is denoted by $\limsup x_n$.

Limit superior

Let (x_n) be a bounded sequence. Let $y_1 = \sup\{x_1, x_2, \dots\}$, $y_2 = \sup\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$y_n = \sup\{x_n, x_{n+1}, \dots\} = \sup_{k \geq n} x_k.$$

We have

$$y_1 \geq y_2 \geq y_3 \geq \dots.$$

By Monotone convergence theorem, (y_n) is convergent and converges to the infimum of $\{y_1, y_2, \dots\}$.

The limit of the sequence (y_n) is called the limit superior of the sequence (x_n) , and is denoted by $\limsup x_n$.

Thus,

$$\limsup x_n := \lim_{n \rightarrow \infty} y_n = \inf\{y_1, y_2, \dots\} = \inf_n \sup_{k \geq n} x_k.$$

Limit inferior

Similarly, let $z_1 = \inf\{x_1, x_2, \dots\}$, $z_2 = \inf\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

Limit inferior

Similarly, let $z_1 = \inf\{x_1, x_2, \dots\}$, $z_2 = \inf\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

Limit inferior

Similarly, let $z_1 = \inf\{x_1, x_2, \dots\}$, $z_2 = \inf\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

By Monotone convergence theorem, (z_n) is convergent and converges to the supremum of $\{z_1, z_2, \dots\}$.

Limit inferior

Similarly, let $z_1 = \inf\{x_1, x_2, \dots\}$, $z_2 = \inf\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

By Monotone convergence theorem, (z_n) is convergent and converges to the supremum of $\{z_1, z_2, \dots\}$.

The limit of the sequence (z_n) is called the limit inferior of the sequence (x_n) , and is denoted by $\liminf x_n$.

Limit inferior

Similarly, let $z_1 = \inf\{x_1, x_2, \dots\}$, $z_2 = \inf\{x_2, x_3, \dots\}$, and so on. That is, for $n \in \mathbb{N}$,

$$z_n = \inf\{x_n, x_{n+1}, \dots\} = \inf_{k \geq n} x_k.$$

We have

$$z_1 \leq z_2 \leq z_3 \leq \dots.$$

By Monotone convergence theorem, (z_n) is convergent and converges to the supremum of $\{z_1, z_2, \dots\}$.

The limit of the sequence (z_n) is called the limit inferior of the sequence (x_n) , and is denoted by $\liminf x_n$.

Thus,

$$\liminf x_n := \lim_{n \rightarrow \infty} z_n = \sup\{z_1, z_2, \dots\} = \sup_n \inf_{k \geq n} x_k.$$

Remark: Suppose that $|x_n| < M$ for $n \in \mathbb{N}$. Then $-M \leq z_n \leq y_n \leq M$ for all n . Hence,

$$-M \leq \liminf x_n \leq \limsup x_n \leq M.$$

Remark: Suppose that $|x_n| < M$ for $n \in \mathbb{N}$. Then $-M \leq z_n \leq y_n \leq M$ for all n . Hence,

$$-M \leq \liminf x_n \leq \limsup x_n \leq M.$$

Example: Consider the sequence (x_n) , where $x_n = (-1)^n$. Clearly, for any n , $y_n = \sup\{x_n, x_{n+1}, \dots\} = 1$ and $z_n = \inf\{x_n, x_{n+1}, \dots\} = -1$. Hence,

$$\limsup x_n = 1 \quad \text{and} \quad \liminf x_n = -1.$$

Remark: Suppose that $|x_n| < M$ for $n \in \mathbb{N}$. Then $-M \leq z_n \leq y_n \leq M$ for all n . Hence,

$$-M \leq \liminf x_n \leq \limsup x_n \leq M.$$

Example: Consider the sequence (x_n) , where $x_n = (-1)^n$. Clearly, for any n , $y_n = \sup\{x_n, x_{n+1}, \dots\} = 1$ and $z_n = \inf\{x_n, x_{n+1}, \dots\} = -1$. Hence,

$$\limsup x_n = 1 \quad \text{and} \quad \liminf x_n = -1.$$

Example: Consider the sequence (x_n) , where $x_n = \frac{1}{n}$.

Clearly, for any n , $y_n = \sup\{\frac{1}{k} : k \geq n\} = \frac{1}{n}$ and

$$z_n = \inf\{\frac{1}{k} : k \geq n\} = 0.$$

Hence, $\limsup x_n = \lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} \frac{1}{n} = 0$ and $\liminf x_n = 0$.

Theorem

Let (a_n) and (b_n) be two bounded sequences.

- ① $\liminf a_n \leq \limsup a_n$.
- ② If $a_n \leq b_n$ for all $n \in \mathbb{N}$, then $\limsup a_n \leq \limsup b_n$ and $\liminf a_n \leq \liminf b_n$.
- ③ $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$ and $\liminf(a_n + b_n) \geq \liminf a_n + \liminf b_n$.

Theorem

Let (a_n) and (b_n) be two bounded sequences.

- ① $\liminf a_n \leq \limsup a_n$.
- ② If $a_n \leq b_n$ for all $n \in \mathbb{N}$, then $\limsup a_n \leq \limsup b_n$ and $\liminf a_n \leq \liminf b_n$.
- ③ $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$ and $\liminf(a_n + b_n) \geq \liminf a_n + \liminf b_n$.

Theorem

If (a_n) is a convergent sequence, then

$$\liminf a_n = \lim_{n \rightarrow \infty} a_n = \limsup a_n.$$

Theorem

Let (a_n) and (b_n) be two bounded sequences.

- ① $\liminf a_n \leq \limsup a_n$.
- ② If $a_n \leq b_n$ for all $n \in \mathbb{N}$, then $\limsup a_n \leq \limsup b_n$ and $\liminf a_n \leq \liminf b_n$.
- ③ $\limsup(a_n + b_n) \leq \limsup a_n + \limsup b_n$ and $\liminf(a_n + b_n) \geq \liminf a_n + \liminf b_n$.

Theorem

If (a_n) is a convergent sequence, then

$$\liminf a_n = \lim_{n \rightarrow \infty} a_n = \limsup a_n.$$

Theorem

Let (a_n) be a bounded sequence. If $\limsup a_n = \liminf a_n$, then (a_n) is convergent and $\lim_{n \rightarrow \infty} a_n = \limsup a_n$.