

MA 101 (Mathematics-I)

Anupam Saikia and Rupam Barman
Department of Mathematics
IIT Guwahati

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an **interior point**. That is, there exists some $\delta > 0$ such that $(x_0 - \delta, x_0 + \delta) \subseteq D$.

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an **interior point**. That is, there exists some $\delta > 0$ such that $(x_0 - \delta, x_0 + \delta) \subseteq D$.

A function $f : D \rightarrow \mathbb{R}$ is said to be differentiable at x_0 if

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \quad \text{or, equivalently} \quad \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

exists in \mathbb{R} .

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an **interior point**. That is, there exists some $\delta > 0$ such that $(x_0 - \delta, x_0 + \delta) \subseteq D$.

A function $f : D \rightarrow \mathbb{R}$ is said to be differentiable at x_0 if

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \quad \text{or, equivalently} \quad \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

exists in \mathbb{R} .

If f is differentiable at x_0 , then the derivative of f at x_0 is

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}.$$

Introduction

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an **interior point**. That is, there exists some $\delta > 0$ such that $(x_0 - \delta, x_0 + \delta) \subseteq D$.

A function $f : D \rightarrow \mathbb{R}$ is said to be differentiable at x_0 if

$$\lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} \quad \text{or, equivalently} \quad \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

exists in \mathbb{R} .

If f is differentiable at x_0 , then the derivative of f at x_0 is

$$f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0} = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}.$$

$f : D \rightarrow \mathbb{R}$ is said to be differentiable if f is differentiable at each $x_0 \in D$.

Theorem: If $f : D \rightarrow \mathbb{R}$ is differentiable at $x_0 \in D$, then f is continuous at x_0 .

Theorem: If $f : D \rightarrow \mathbb{R}$ is differentiable at $x_0 \in D$, then f is continuous at x_0 .

Example

Theorem: If $f : D \rightarrow \mathbb{R}$ is differentiable at $x_0 \in D$, then f is continuous at x_0 .

Example

(a) $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = |x|$ is not differentiable at 0.

Theorem: If $f : D \rightarrow \mathbb{R}$ is differentiable at $x_0 \in D$, then f is continuous at x_0 .

Example

(a) $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = |x|$ is not differentiable at 0.

(b) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$

Then f is not differentiable at 0.

Theorem: If $f : D \rightarrow \mathbb{R}$ is differentiable at $x_0 \in D$, then f is continuous at x_0 .

Example

(a) $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = |x|$ is not differentiable at 0.

(b) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \begin{cases} x \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$

Then f is not differentiable at 0.

(c) $f(x) = \begin{cases} x^2 & \text{if } x \in \mathbb{Q}, \\ 0 & \text{if } x \in \mathbb{R} \setminus \mathbb{Q}. \end{cases}$ Then $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable only at 0 and $f'(0) = 0$.

Theorem (Rules for finding derivatives)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an interior point. Suppose $f, g : D \rightarrow \mathbb{R}$ are differentiable at x_0 . Then

Theorem (Rules for finding derivatives)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an interior point. Suppose $f, g : D \rightarrow \mathbb{R}$ are differentiable at x_0 . Then

- (a) If $\alpha \in \mathbb{R}$, then the function αf is differentiable at x_0 and $(\alpha f)'(x_0) = \alpha f'(x_0)$.

Theorem (Rules for finding derivatives)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an interior point. Suppose $f, g : D \rightarrow \mathbb{R}$ are differentiable at x_0 . Then

- (a) If $\alpha \in \mathbb{R}$, then the function αf is differentiable at x_0 and $(\alpha f)'(x_0) = \alpha f'(x_0)$.
- (b) The function $f + g$ is differentiable at x_0 and $(f + g)'(x_0) = f'(x_0) + g'(x_0)$.

Theorem (Rules for finding derivatives)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an interior point. Suppose $f, g : D \rightarrow \mathbb{R}$ are differentiable at x_0 . Then

- (a) If $\alpha \in \mathbb{R}$, then the function αf is differentiable at x_0 and $(\alpha f)'(x_0) = \alpha f'(x_0)$.
- (b) The function $f + g$ is differentiable at x_0 and $(f + g)'(x_0) = f'(x_0) + g'(x_0)$.
- (c) (Product rule) The function fg is differentiable at x_0 and $(fg)'(x_0) = f'(x_0)g(x_0) + f(x_0)g'(x_0)$.

Theorem (Rules for finding derivatives)

Let $D \subseteq \mathbb{R}$ and let $x_0 \in D$ be an interior point. Suppose $f, g : D \rightarrow \mathbb{R}$ are differentiable at x_0 . Then

- (a) If $\alpha \in \mathbb{R}$, then the function αf is differentiable at x_0 and $(\alpha f)'(x_0) = \alpha f'(x_0)$.
- (b) The function $f + g$ is differentiable at x_0 and $(f + g)'(x_0) = f'(x_0) + g'(x_0)$.
- (c) (Product rule) The function fg is differentiable at x_0 and $(fg)'(x_0) = f'(x_0)g(x_0) + f(x_0)g'(x_0)$.
- (d) (Quotient rule) If $g(x_0) \neq 0$, then the function f/g is differentiable at x_0 and

$$(f/g)'(x_0) = \frac{f'(x_0)g(x_0) - f(x_0)g'(x_0)}{(g(x_0))^2}.$$

Theorem (Chain rule for derivative)

Let $f : D \rightarrow \mathbb{R}$ and $g : E \rightarrow \mathbb{R}$. Let $f(D) \subseteq E$. Suppose that f is differentiable at x_0 and g is differentiable at $f(x_0)$. Then $g \circ f$ is differentiable at x_0 and $(g \circ f)'(x_0) = g'(f(x_0))f'(x_0)$.

Theorem (Chain rule for derivative)

Let $f : D \rightarrow \mathbb{R}$ and $g : E \rightarrow \mathbb{R}$. Let $f(D) \subseteq E$. Suppose that f is differentiable at x_0 and g is differentiable at $f(x_0)$. Then $g \circ f$ is differentiable at x_0 and $(g \circ f)'(x_0) = g'(f(x_0))f'(x_0)$.

Proof: We define a function $h : E \rightarrow \mathbb{R}$ by

$$h(y) = \begin{cases} \frac{g(y) - g(f(x_0))}{y - f(x_0)}, & y \neq f(x_0); \\ g'(f(x_0)), & y = f(x_0). \end{cases}$$

Theorem (Chain rule for derivative)

Let $f : D \rightarrow \mathbb{R}$ and $g : E \rightarrow \mathbb{R}$. Let $f(D) \subseteq E$. Suppose that f is differentiable at x_0 and g is differentiable at $f(x_0)$. Then $g \circ f$ is differentiable at x_0 and $(g \circ f)'(x_0) = g'(f(x_0))f'(x_0)$.

Proof: We define a function $h : E \rightarrow \mathbb{R}$ by

$$h(y) = \begin{cases} \frac{g(y) - g(f(x_0))}{y - f(x_0)}, & y \neq f(x_0); \\ g'(f(x_0)), & y = f(x_0). \end{cases}$$

We have $\lim_{y \rightarrow f(x_0)} h(y) = g'(f(x_0)) = h(f(x_0))$, and $g(y) - g(f(x_0)) = h(y) \times (y - f(x_0))$ for all $y \in E$.

Theorem (Chain rule for derivative)

Let $f : D \rightarrow \mathbb{R}$ and $g : E \rightarrow \mathbb{R}$. Let $f(D) \subseteq E$. Suppose that f is differentiable at x_0 and g is differentiable at $f(x_0)$. Then $g \circ f$ is differentiable at x_0 and $(g \circ f)'(x_0) = g'(f(x_0))f'(x_0)$.

Proof: We define a function $h : E \rightarrow \mathbb{R}$ by

$$h(y) = \begin{cases} \frac{g(y) - g(f(x_0))}{y - f(x_0)}, & y \neq f(x_0); \\ g'(f(x_0)), & y = f(x_0). \end{cases}$$

We have $\lim_{y \rightarrow f(x_0)} h(y) = g'(f(x_0)) = h(f(x_0))$, and $g(y) - g(f(x_0)) = h(y) \times (y - f(x_0))$ for all $y \in E$.

Hence, for $x \neq x_0$,

$$\frac{g(f(x)) - g(f(x_0))}{x - x_0} = h(f(x)) \frac{f(x) - f(x_0)}{x - x_0}$$

Example: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Then f is differentiable at 0. But $f' : \mathbb{R} \rightarrow \mathbb{R}$ is not continuous at 0.

Example: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Then f is differentiable at 0. But $f' : \mathbb{R} \rightarrow \mathbb{R}$ is not continuous at 0.

Clearly f is differentiable at all $x (\neq 0) \in \mathbb{R}$ and

$$f'(x) = 2x \sin \frac{1}{x} - \cos \frac{1}{x} \text{ for all } x (\neq 0) \in \mathbb{R}.$$

Example: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Then f is differentiable at 0. But $f' : \mathbb{R} \rightarrow \mathbb{R}$ is not continuous at 0.

Clearly f is differentiable at all $x (\neq 0) \in \mathbb{R}$ and

$$f'(x) = 2x \sin \frac{1}{x} - \cos \frac{1}{x} \text{ for all } x (\neq 0) \in \mathbb{R}.$$

For each $\varepsilon > 0$, choosing $\delta = \varepsilon > 0$, we find that

$$\left| \frac{f(x) - f(0)}{x - 0} \right| = \left| x \sin \frac{1}{x} \right| \leq |x| < \varepsilon \text{ for all } x \in \mathbb{R} \text{ satisfying } 0 < |x| < \delta. \text{ Hence, } f \text{ is differentiable at } 0 \text{ and } f'(0) = 0.$$

Example: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Then f is differentiable at 0. But $f' : \mathbb{R} \rightarrow \mathbb{R}$ is not continuous at 0.

Clearly f is differentiable at all $x (\neq 0) \in \mathbb{R}$ and

$$f'(x) = 2x \sin \frac{1}{x} - \cos \frac{1}{x} \text{ for all } x (\neq 0) \in \mathbb{R}.$$

For each $\varepsilon > 0$, choosing $\delta = \varepsilon > 0$, we find that

$$\left| \frac{f(x) - f(0)}{x - 0} \right| = \left| x \sin \frac{1}{x} \right| \leq |x| < \varepsilon \text{ for all } x \in \mathbb{R} \text{ satisfying } 0 < |x| < \delta. \text{ Hence, } f \text{ is differentiable at } 0 \text{ and } f'(0) = 0.$$

Since $\frac{1}{2n\pi} \rightarrow 0$ but $f'(\frac{1}{2n\pi}) \rightarrow -1 \neq f'(0)$, $f' : \mathbb{R} \rightarrow \mathbb{R}$ is not continuous at 0.

Definition: $f : D \rightarrow \mathbb{R}$ has a local maximum (resp. minimum) at $x_0 \in D$ if there exists $\delta > 0$ such that $f(x) \leq f(x_0)$ (resp. $f(x_0) \leq f(x)$) for all $x \in (x_0 - \delta, x_0 + \delta) \cap D$.

Definition: $f : D \rightarrow \mathbb{R}$ has a local maximum (resp. minimum) at $x_0 \in D$ if there exists $\delta > 0$ such that $f(x) \leq f(x_0)$ (resp. $f(x_0) \leq f(x)$) for all $x \in (x_0 - \delta, x_0 + \delta) \cap D$.

Theorem: If $f : D \rightarrow \mathbb{R}$ has a local maximum or local minimum at an interior point x_0 of D and if f is differentiable at x_0 , then $f'(x_0) = 0$.

Definition: $f : D \rightarrow \mathbb{R}$ has a local maximum (resp. minimum) at $x_0 \in D$ if there exists $\delta > 0$ such that $f(x) \leq f(x_0)$ (resp. $f(x_0) \leq f(x)$) for all $x \in (x_0 - \delta, x_0 + \delta) \cap D$.

Theorem: If $f : D \rightarrow \mathbb{R}$ has a local maximum or local minimum at an interior point x_0 of D and if f is differentiable at x_0 , then $f'(x_0) = 0$.

Rolle's Theorem: Let $f : [a, b] \rightarrow \mathbb{R}$. Suppose that

- (a) f is continuous on $[a, b]$.
- (b) f is differentiable on (a, b) .
- (c) $f(a) = f(b)$.

Then there exists $c \in (a, b)$ such that $f'(c) = 0$.

Definition: $f : D \rightarrow \mathbb{R}$ has a local maximum (resp. minimum) at $x_0 \in D$ if there exists $\delta > 0$ such that $f(x) \leq f(x_0)$ (resp. $f(x_0) \leq f(x)$) for all $x \in (x_0 - \delta, x_0 + \delta) \cap D$.

Theorem: If $f : D \rightarrow \mathbb{R}$ has a local maximum or local minimum at an interior point x_0 of D and if f is differentiable at x_0 , then $f'(x_0) = 0$.

Rolle's Theorem: Let $f : [a, b] \rightarrow \mathbb{R}$. Suppose that

- (a) f is continuous on $[a, b]$.
- (b) f is differentiable on (a, b) .
- (c) $f(a) = f(b)$.

Then there exists $c \in (a, b)$ such that $f'(c) = 0$.

Examples:

- a The equation $x^2 = x \sin x + \cos x$ has exactly two real roots.

Definition: $f : D \rightarrow \mathbb{R}$ has a local maximum (resp. minimum) at $x_0 \in D$ if there exists $\delta > 0$ such that $f(x) \leq f(x_0)$ (resp. $f(x_0) \leq f(x)$) for all $x \in (x_0 - \delta, x_0 + \delta) \cap D$.

Theorem: If $f : D \rightarrow \mathbb{R}$ has a local maximum or local minimum at an interior point x_0 of D and if f is differentiable at x_0 , then $f'(x_0) = 0$.

Rolle's Theorem: Let $f : [a, b] \rightarrow \mathbb{R}$. Suppose that

- (a) f is continuous on $[a, b]$.
- (b) f is differentiable on (a, b) .
- (c) $f(a) = f(b)$.

Then there exists $c \in (a, b)$ such that $f'(c) = 0$.

Examples:

- a The equation $x^2 = x \sin x + \cos x$ has exactly two real roots.
- b The equation $x^4 + 2x^2 - 6x + 2 = 0$ has exactly two real roots.

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Result: Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable. Then

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Result: Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable. Then

- a $f'(x) = 0$ for all $x \in I \Leftrightarrow f$ is constant on I .

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Result: Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable. Then

- a $f'(x) = 0$ for all $x \in I \Leftrightarrow f$ is constant on I .
- b $f'(x) \geq 0$ for all $x \in I \Leftrightarrow f$ is increasing on I .

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Result: Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable. Then

- a $f'(x) = 0$ for all $x \in I \Leftrightarrow f$ is constant on I .
- b $f'(x) \geq 0$ for all $x \in I \Leftrightarrow f$ is increasing on I .
- c $f'(x) \leq 0$ for all $x \in I \Leftrightarrow f$ is decreasing on I .

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Result: Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable. Then

- a $f'(x) = 0$ for all $x \in I \Leftrightarrow f$ is constant on I .
- b $f'(x) \geq 0$ for all $x \in I \Leftrightarrow f$ is increasing on I .
- c $f'(x) \leq 0$ for all $x \in I \Leftrightarrow f$ is decreasing on I .
- d $f'(x) > 0$ for all $x \in I \Rightarrow f$ is strictly increasing on I .

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Result: Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable. Then

- a $f'(x) = 0$ for all $x \in I \Leftrightarrow f$ is constant on I .
- b $f'(x) \geq 0$ for all $x \in I \Leftrightarrow f$ is increasing on I .
- c $f'(x) \leq 0$ for all $x \in I \Leftrightarrow f$ is decreasing on I .
- d $f'(x) > 0$ for all $x \in I \Rightarrow f$ is strictly increasing on I .
- e $f'(x) < 0$ for all $x \in I \Rightarrow f$ is strictly decreasing on I .

Theorem (Mean value theorem) If $f : [a, b] \rightarrow \mathbb{R}$ is continuous and if f is differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f(b) - f(a) = f'(c)(b - a)$.

Result: Let I be an interval and let $f : I \rightarrow \mathbb{R}$ be differentiable. Then

- a $f'(x) = 0$ for all $x \in I \Leftrightarrow f$ is constant on I .
- b $f'(x) \geq 0$ for all $x \in I \Leftrightarrow f$ is increasing on I .
- c $f'(x) \leq 0$ for all $x \in I \Leftrightarrow f$ is decreasing on I .
- d $f'(x) > 0$ for all $x \in I \Rightarrow f$ is strictly increasing on I .
- e $f'(x) < 0$ for all $x \in I \Rightarrow f$ is strictly decreasing on I .
- f $f'(x) \neq 0$ for all $x \in I \Rightarrow f$ is one-one on I .

Remark: Note that f is strictly increasing on I need not imply that $f'(x) > 0$ for all $x \in I$. For example, $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = x^3$ is strictly increasing but $f'(0) = 0$.

Remark: Note that f is strictly increasing on I need not imply that $f'(x) > 0$ for all $x \in I$. For example, $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = x^3$ is strictly increasing but $f'(0) = 0$.

Examples:

- a $\sin x \leq x$ for all $x \geq 0$.

Remark: Note that f is strictly increasing on I need not imply that $f'(x) > 0$ for all $x \in I$. For example, $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = x^3$ is strictly increasing but $f'(0) = 0$.

Examples:

- a $\sin x \leq x$ for all $x \geq 0$.
- b $\sin x \geq x - \frac{x^3}{6}$ for all $x \in [0, \frac{\pi}{2}]$.

Remark: Note that f is strictly increasing on I need not imply that $f'(x) > 0$ for all $x \in I$. For example, $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = x^3$ is strictly increasing but $f'(0) = 0$.

Examples:

- a $\sin x \leq x$ for all $x \geq 0$.
- b $\sin x \geq x - \frac{x^3}{6}$ for all $x \in [0, \frac{\pi}{2}]$.
- c If $f(x) = x^3 + x^2 - 5x + 3$ for all $x \in \mathbb{R}$, then f is one-one on $[1, 5]$ but not one-one on \mathbb{R} .

L'Hôpital's rules:

- (1) Let $f : (a, b) \rightarrow \mathbb{R}$ and $g : (a, b) \rightarrow \mathbb{R}$ be differentiable at $x_0 \in (a, b)$. Also, let $f(x_0) = g(x_0) = 0$ and $g'(x_0) \neq 0$.
Then $\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \frac{f'(x_0)}{g'(x_0)}$.

L'Hôpital's rules:

- (1) Let $f : (a, b) \rightarrow \mathbb{R}$ and $g : (a, b) \rightarrow \mathbb{R}$ be differentiable at $x_0 \in (a, b)$. Also, let $f(x_0) = g(x_0) = 0$ and $g'(x_0) \neq 0$.
Then $\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \frac{f'(x_0)}{g'(x_0)}$.
- (2) Let $f : (a, b) \rightarrow \mathbb{R}$ and $g : (a, b) \rightarrow \mathbb{R}$ be differentiable such that $\lim_{x \rightarrow a+} f(x) = \lim_{x \rightarrow a+} g(x) = 0$ and $g'(x) \neq 0$ for all $x \in (a, b)$. If $\lim_{x \rightarrow a+} \frac{f'(x)}{g'(x)} = \ell$, then $\lim_{x \rightarrow a+} \frac{f(x)}{g(x)} = \ell$.

L'Hôpital's rules:

- (1) Let $f : (a, b) \rightarrow \mathbb{R}$ and $g : (a, b) \rightarrow \mathbb{R}$ be differentiable at $x_0 \in (a, b)$. Also, let $f(x_0) = g(x_0) = 0$ and $g'(x_0) \neq 0$.
Then $\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \frac{f'(x_0)}{g'(x_0)}$.
- (2) Let $f : (a, b) \rightarrow \mathbb{R}$ and $g : (a, b) \rightarrow \mathbb{R}$ be differentiable such that $\lim_{x \rightarrow a+} f(x) = \lim_{x \rightarrow a+} g(x) = 0$ and $g'(x) \neq 0$ for all $x \in (a, b)$. If $\lim_{x \rightarrow a+} \frac{f'(x)}{g'(x)} = \ell$, then $\lim_{x \rightarrow a+} \frac{f(x)}{g(x)} = \ell$.

Examples: (a) $\lim_{x \rightarrow 0} \frac{\sqrt{1+x}-1}{x}$

(b) $\lim_{x \rightarrow \frac{\pi}{2}} \frac{1-\sin x}{1+\cos 2x}$

A very useful technique in the analysis of real functions is the approximation of functions by polynomials.

A very useful technique in the analysis of real functions is the approximation of functions by polynomials.

Theorem (Taylor's theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be such that f and its derivatives $f', f'', \dots, f^{(n)}$ are continuous on $[a, b]$ and that $f^{(n+1)}$ exists on (a, b) . If $x_0 \in [a, b]$, then for any $x \in [a, b]$ there exists a point c between x and x_0 such that

$$\begin{aligned} f(x) &= f(x_0) + f'(x_0)(x - x_0) + \frac{f''(x_0)}{2!}(x - x_0)^2 + \dots \\ &\quad \dots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n + \frac{f^{(n+1)}(c)}{(n+1)!}(x - x_0)^{n+1} \\ &= P_n(x) + \frac{f^{(n+1)}(c)}{(n+1)!}(x - x_0)^{n+1} \end{aligned}$$

Proof: Let x and x_0 be given and let J denote the closed interval with endpoints x_0 and x . Define F on J by

Proof: Let x and x_0 be given and let J denote the closed interval with endpoints x_0 and x . Define F on J by

$$F(t) := f(x) - f(t) - f'(t) \cdot (x - t) - \cdots - \frac{f^{(n)}(t)}{n!} (x - t)^n$$

$$\text{We have } F'(t) = -\frac{(x - t)^n}{n!} f^{(n+1)}(t).$$

Proof: Let x and x_0 be given and let J denote the closed interval with endpoints x_0 and x . Define F on J by

$$F(t) := f(x) - f(t) - f'(t) \cdot (x - t) - \cdots - \frac{f^{(n)}(t)}{n!} (x - t)^n$$

$$\text{We have } F'(t) = -\frac{(x - t)^n}{n!} f^{(n+1)}(t).$$

$$\text{Define } G(t) := F(t) - \left(\frac{x - t}{x - x_0} \right)^{n+1} F(x_0)$$

Proof: Let x and x_0 be given and let J denote the closed interval with endpoints x_0 and x . Define F on J by

$$F(t) := f(x) - f(t) - f'(t) \cdot (x - t) - \cdots - \frac{f^{(n)}(t)}{n!} (x - t)^n$$

$$\text{We have } F'(t) = -\frac{(x - t)^n}{n!} f^{(n+1)}(t).$$

$$\text{Define } G(t) := F(t) - \left(\frac{x - t}{x - x_0} \right)^{n+1} F(x_0)$$

$$\text{By Rolle's thm } 0 = G'(c) = F'(c) + (n + 1) \frac{(x - c)^n}{(x - x_0)^{n+1}} F(x_0).$$

Example: $1 + \frac{x}{2} - \frac{x^2}{8} \leq \sqrt{1+x} \leq 1 + \frac{x}{2}$ for all $x > 0$.

Example: $1 + \frac{x}{2} - \frac{x^2}{8} \leq \sqrt{1+x} \leq 1 + \frac{x}{2}$ for all $x > 0$.

Solution: Let $x > 0$ and let $f(t) = \sqrt{1+t}$ for all $x \in [0, x]$.

Example: $1 + \frac{x}{2} - \frac{x^2}{8} \leq \sqrt{1+x} \leq 1 + \frac{x}{2}$ for all $x > 0$.

Solution: Let $x > 0$ and let $f(t) = \sqrt{1+t}$ for all $x \in [0, x]$.

Then $f : [0, x] \rightarrow \mathbb{R}$ is twice differentiable and $f'(t) = \frac{1}{2\sqrt{1+t}}$,
 $f''(t) = -\frac{1}{4(1+t)^{3/2}}$ for all $t \in [0, x]$.

Example: $1 + \frac{x}{2} - \frac{x^2}{8} \leq \sqrt{1+x} \leq 1 + \frac{x}{2}$ for all $x > 0$.

Solution: Let $x > 0$ and let $f(t) = \sqrt{1+t}$ for all $x \in [0, x]$.

Then $f : [0, x] \rightarrow \mathbb{R}$ is twice differentiable and $f'(t) = \frac{1}{2\sqrt{1+t}}$,
 $f''(t) = -\frac{1}{4(1+t)^{3/2}}$ for all $t \in [0, x]$.

By Taylor's theorem, there exists $c \in (0, x)$ such that
 $f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(c) = 1 + \frac{x}{2} - \frac{x^2}{8} \cdot \frac{1}{(1+c)^{3/2}}.$

Example: $1 + \frac{x}{2} - \frac{x^2}{8} \leq \sqrt{1+x} \leq 1 + \frac{x}{2}$ for all $x > 0$.

Solution: Let $x > 0$ and let $f(t) = \sqrt{1+t}$ for all $x \in [0, x]$.

Then $f : [0, x] \rightarrow \mathbb{R}$ is twice differentiable and $f'(t) = \frac{1}{2\sqrt{1+t}}$,
 $f''(t) = -\frac{1}{4(1+t)^{3/2}}$ for all $t \in [0, x]$.

By Taylor's theorem, there exists $c \in (0, x)$ such that
 $f(x) = f(0) + xf'(0) + \frac{x^2}{2!}f''(c) = 1 + \frac{x}{2} - \frac{x^2}{8} \cdot \frac{1}{(1+c)^{3/2}}.$

Since $0 < \frac{1}{(1+c)^{3/2}} < 1$, we get $1 + \frac{x}{2} - \frac{x^2}{8} \leq \sqrt{1+x} \leq 1 + \frac{x}{2}.$

Theorem: Let $x_0 \in (a, b)$ and let $n \geq 2$. Also, let $f, f', \dots, f^{(n)}$ be continuous on (a, b) and $f'(x_0) = f''(x_0) = \dots = f^{(n-1)}(x_0) = 0$ but $f^{(n)}(x_0) \neq 0$.

Theorem: Let $x_0 \in (a, b)$ and let $n \geq 2$. Also, let $f, f', \dots, f^{(n)}$ be continuous on (a, b) and $f'(x_0) = f''(x_0) = \dots = f^{(n-1)}(x_0) = 0$ but $f^{(n)}(x_0) \neq 0$.

(a) If n is even and $f^{(n)}(x_0) < 0$, then f has a local maximum at x_0 .

Theorem: Let $x_0 \in (a, b)$ and let $n \geq 2$. Also, let $f, f', \dots, f^{(n)}$ be continuous on (a, b) and $f'(x_0) = f''(x_0) = \dots = f^{(n-1)}(x_0) = 0$ but $f^{(n)}(x_0) \neq 0$.

- (a) If n is even and $f^{(n)}(x_0) < 0$, then f has a local maximum at x_0 .
- (b) If n is even and $f^{(n)}(x_0) > 0$, then f has a local minimum at x_0 .

Theorem: Let $x_0 \in (a, b)$ and let $n \geq 2$. Also, let $f, f', \dots, f^{(n)}$ be continuous on (a, b) and $f'(x_0) = f''(x_0) = \dots = f^{(n-1)}(x_0) = 0$ but $f^{(n)}(x_0) \neq 0$.

- (a) If n is even and $f^{(n)}(x_0) < 0$, then f has a local maximum at x_0 .
- (b) If n is even and $f^{(n)}(x_0) > 0$, then f has a local minimum at x_0 .
- (c) If n is odd, then f has neither a local maximum nor a local minimum at x_0 .

Theorem: Let $x_0 \in (a, b)$ and let $n \geq 2$. Also, let $f, f', \dots, f^{(n)}$ be continuous on (a, b) and $f'(x_0) = f''(x_0) = \dots = f^{(n-1)}(x_0) = 0$ but $f^{(n)}(x_0) \neq 0$.

- (a) If n is even and $f^{(n)}(x_0) < 0$, then f has a local maximum at x_0 .
- (b) If n is even and $f^{(n)}(x_0) > 0$, then f has a local minimum at x_0 .
- (c) If n is odd, then f has neither a local maximum nor a local minimum at x_0 .

Example: Find local maximum and local minimum values of f , where $f(x) = x^5 - 5x^4 + 5x^3 + 12$ for all $x \in \mathbb{R}$.

Power series: A series of the form $\sum_{n=0}^{\infty} a_n(x - x_0)^n$, where x_0 , $a_n \in \mathbb{R}$ for $n = 0, 1, 2, \dots$ and $x \in \mathbb{R}$.

Power series: A series of the form $\sum_{n=0}^{\infty} a_n(x - x_0)^n$, where x_0 , $a_n \in \mathbb{R}$ for $n = 0, 1, 2, \dots$ and $x \in \mathbb{R}$.

It is sufficient to consider the series $\sum_{n=0}^{\infty} a_n x^n$.

Power series: A series of the form $\sum_{n=0}^{\infty} a_n(x - x_0)^n$, where x_0 , $a_n \in \mathbb{R}$ for $n = 0, 1, 2, \dots$ and $x \in \mathbb{R}$.

It is sufficient to consider the series $\sum_{n=0}^{\infty} a_n x^n$.

Convergence - Examples:

$$(a) \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (b) \sum_{n=0}^{\infty} n! x^n \quad (c) \sum_{n=0}^{\infty} x^n$$

Power series: A series of the form $\sum_{n=0}^{\infty} a_n(x - x_0)^n$, where x_0 , $a_n \in \mathbb{R}$ for $n = 0, 1, 2, \dots$ and $x \in \mathbb{R}$.

It is sufficient to consider the series $\sum_{n=0}^{\infty} a_n x^n$.

Convergence - Examples:

$$(a) \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (b) \sum_{n=0}^{\infty} n! x^n \quad (c) \sum_{n=0}^{\infty} x^n$$

Theorem:

- a If $\sum_{n=0}^{\infty} a_n x^n$ converges for $x = x_1 \neq 0$, then it converges absolutely for all $x \in \mathbb{R}$ satisfying $|x| < |x_1|$.
- b If $\sum_{n=0}^{\infty} a_n x^n$ diverges for $x = x_2$, then it diverges for all $x \in \mathbb{R}$ satisfying $|x| > |x_2|$.

Radius of convergence: For every power series $\sum_{n=0}^{\infty} a_n x^n$, there exists a unique R satisfying $0 \leq R \leq \infty$ such that the series converges absolutely if $|x| < R$ and diverges if $|x| > R$.

Radius of convergence: For every power series $\sum_{n=0}^{\infty} a_n x^n$, there exists a unique R satisfying $0 \leq R \leq \infty$ such that the series converges absolutely if $|x| < R$ and diverges if $|x| > R$.

Theorem

Consider the power series $\sum_{n=0}^{\infty} a_n x^n$. Let $\beta = \limsup \sqrt[n]{|a_n|}$ and $R = \frac{1}{\beta}$ (we define $R = 0$ if $\beta = \infty$ and $R = \infty$ if $\beta = 0$).

Then

- (a) $\sum_{n=0}^{\infty} a_n x^n$ converges for $|x| < R$
- (b) $\sum_{n=0}^{\infty} a_n x^n$ diverges for $|x| > R$.
- (c) No conclusion if $|x| = R$.

Theorem

Consider the power series $\sum_{n=0}^{\infty} a_n x^n$. Suppose $\beta = \lim \left| \frac{a_{n+1}}{a_n} \right|$ and $R = \frac{1}{\beta}$ (We define $R = 0$ if $\beta = \infty$ and $R = \infty$ if $\beta = 0$). Then

- (a) $\sum_{n=0}^{\infty} a_n x^n$ converges for $|x| < R$
- (b) $\sum_{n=0}^{\infty} a_n x^n$ diverges for $|x| > R$.
- (c) No conclusion if $|x| = R$.

Examples: (a) $\sum_{n=0}^{\infty} \frac{x^n}{n^2}$ (b) $\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n$

Examples: (a) $\sum_{n=0}^{\infty} \frac{x^n}{n^2}$ (b) $\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n$

Proof of (a) (Method-1): If $x = 0$, then the given series becomes $0 + 0 + \cdots$, which is clearly convergent. Let $x (\neq 0) \in \mathbb{R}$ and let $a_n = \frac{x^n}{n^2}$ for all $n \in \mathbb{N}$.

Examples: (a) $\sum_{n=0}^{\infty} \frac{x^n}{n^2}$ (b) $\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n$

Proof of (a) (Method-1): If $x = 0$, then the given series becomes $0 + 0 + \dots$, which is clearly convergent. Let $x (\neq 0) \in \mathbb{R}$ and let $a_n = \frac{x^n}{n^2}$ for all $n \in \mathbb{N}$.

Then $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = |x|$. Hence by ratio test, $\sum_{n=1}^{\infty} a_n$ is convergent (absolutely) if $|x| < 1$ and is not convergent if $|x| > 1$. Therefore the radius of convergence of the given power series is 1.

Examples: (a) $\sum_{n=0}^{\infty} \frac{x^n}{n^2}$ (b) $\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n$

Proof of (a) (Method-1): If $x = 0$, then the given series becomes $0 + 0 + \cdots$, which is clearly convergent. Let $x (\neq 0) \in \mathbb{R}$ and let $a_n = \frac{x^n}{n^2}$ for all $n \in \mathbb{N}$.

Then $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = |x|$. Hence by ratio test, $\sum_{n=1}^{\infty} a_n$ is convergent (absolutely) if $|x| < 1$ and is not convergent if $|x| > 1$. Therefore the radius of convergence of the given power series is 1.

Again, if $|x| = 1$, then $\sum_{n=1}^{\infty} |a_n| = \sum_{n=1}^{\infty} \frac{1}{n^2}$ is convergent and hence $\sum_{n=1}^{\infty} a_n$ is also convergent. Therefore the interval of convergence of the given power series is $[-1, 1]$.

Proof of (b) (Method-2): The given power series is

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n. \text{ Let } a_n = \frac{(-1)^n}{n \cdot 4^n} \text{ for all } n \in \mathbb{N}. \text{ Clearly,}$$
$$\beta = \lim \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{4}.$$

Proof of (b) (Method-2): The given power series is

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n. \text{ Let } a_n = \frac{(-1)^n}{n \cdot 4^n} \text{ for all } n \in \mathbb{N}. \text{ Clearly,}$$
$$\beta = \lim \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{4}.$$

Therefore the radius of convergence of the given power series is 4. Thus the given series is convergent (absolutely) if $|x-1| < 4$, that is, if $x \in (-3, 5)$ and is not convergent if $|x-1| > 4$, that is, if $x \in (-\infty, -3) \cup (5, \infty)$.

Proof of (b) (Method-2): The given power series is

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n. \text{ Let } a_n = \frac{(-1)^n}{n \cdot 4^n} \text{ for all } n \in \mathbb{N}. \text{ Clearly,}$$

$$\beta = \lim \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{4}.$$

Therefore the radius of convergence of the given power series is 4. Thus the given series is convergent (absolutely) if $|x-1| < 4$, that is, if $x \in (-3, 5)$ and is not convergent if $|x-1| > 4$, that is, if $x \in (-\infty, -3) \cup (5, \infty)$.

Again, if $x = -3$, then $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \frac{1}{n}$ is not convergent.

Proof of (b) (Method-2): The given power series is

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n \cdot 4^n} (x-1)^n. \text{ Let } a_n = \frac{(-1)^n}{n \cdot 4^n} \text{ for all } n \in \mathbb{N}. \text{ Clearly,}$$

$$\beta = \lim \left| \frac{a_{n+1}}{a_n} \right| = \frac{1}{4}.$$

Therefore the radius of convergence of the given power series is 4. Thus the given series is convergent (absolutely) if $|x-1| < 4$, that is, if $x \in (-3, 5)$ and is not convergent if $|x-1| > 4$, that is, if $x \in (-\infty, -3) \cup (5, \infty)$.

Again, if $x = -3$, then $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \frac{1}{n}$ is not convergent.

If $x = 5$, then $\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \frac{(-1)^n}{n}$ is convergent by Leibniz test.

Therefore the interval of convergence of the given power series is $(-3, 5]$.

Theorem (Term by term differentiation of power series)

A power series can be differentiated term by term within the interval of convergence. In fact, if $f(x) = \sum_{n=0}^{\infty} a_n x^n$ has radius of convergence R , then $f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$ for $|x| < R$.

Theorem (Term by term differentiation of power series)

A power series can be differentiated term by term within the interval of convergence. In fact, if $f(x) = \sum_{n=0}^{\infty} a_n x^n$ has radius

of convergence R , then $f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$ for $|x| < R$.

$$\limsup \sqrt[n]{|n a_n|} = \limsup \{ n^{\frac{1}{n}} \sqrt[n]{|a_n|} \} = \limsup \sqrt[n]{|a_n|} = \beta.$$

Theorem (Term by term differentiation of power series)

A power series can be differentiated term by term within the interval of convergence. In fact, if $f(x) = \sum_{n=0}^{\infty} a_n x^n$ has radius

of convergence R , then $f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$ for $|x| < R$.

$$\limsup \sqrt[n]{|n a_n|} = \limsup \{n^{\frac{1}{n}} \sqrt[n]{|a_n|}\} = \limsup \sqrt[n]{|a_n|} = \beta.$$

Hence, both the series have the same radius of convergence.

Theorem (Term by term differentiation of power series)

A power series can be differentiated term by term within the interval of convergence. In fact, if $f(x) = \sum_{n=0}^{\infty} a_n x^n$ has radius

of convergence R , then $f'(x) = \sum_{n=1}^{\infty} n a_n x^{n-1}$ for $|x| < R$.

$$\limsup \sqrt[n]{|n a_n|} = \limsup \{n^{\frac{1}{n}} \sqrt[n]{|a_n|}\} = \limsup \sqrt[n]{|a_n|} = \beta.$$

Hence, both the series have the same radius of convergence.

To prove that the series $\sum_{n=1}^{\infty} n a_n x^{n-1}$ converges to $f'(x)$ requires another concept called **uniform convergence** which is beyond the scope of this course.

If a function f has derivatives of all orders at a point $c \in \mathbb{R}$, then we can calculate the Taylor coefficients by

$$a_0 = f(c), a_n = \frac{f^{(n)}(c)}{n!} \text{ for all } n \in \mathbb{N}.$$

If a function f has derivatives of all orders at a point $c \in \mathbb{R}$, then we can calculate the Taylor coefficients by

$$a_0 = f(c), a_n = \frac{f^{(n)}(c)}{n!} \text{ for all } n \in \mathbb{N}.$$

In this way, we obtain a power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$.

If a function f has derivatives of all orders at a point $c \in \mathbb{R}$, then we can calculate the Taylor coefficients by

$$a_0 = f(c), a_n = \frac{f^{(n)}(c)}{n!} \text{ for all } n \in \mathbb{N}.$$

In this way, we obtain a power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$.

The power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$ converges to $f(x)$ for

$|x - c| < R$ if and only if the sequence $(R_n(x))$ of remainders converges to 0 for each x in $|x - c| < R$.

If a function f has derivatives of all orders at a point $c \in \mathbb{R}$, then we can calculate the Taylor coefficients by

$$a_0 = f(c), a_n = \frac{f^{(n)}(c)}{n!} \text{ for all } n \in \mathbb{N}.$$

In this way, we obtain a power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$.

The power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$ converges to $f(x)$ for $|x - c| < R$ if and only if the sequence $(R_n(x))$ of remainders converges to 0 for each x in $|x - c| < R$.

The power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$ is called the **Taylor series** of f at c .

If a function f has derivatives of all orders at a point $c \in \mathbb{R}$, then we can calculate the Taylor coefficients by

$$a_0 = f(c), a_n = \frac{f^{(n)}(c)}{n!} \text{ for all } n \in \mathbb{N}.$$

In this way, we obtain a power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$.

The power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$ converges to $f(x)$ for $|x - c| < R$ if and only if the sequence $(R_n(x))$ of remainders converges to 0 for each x in $|x - c| < R$.

The power series $\sum_{n=0}^{\infty} \frac{f^{(n)}(c)}{n!} (x - c)^n$ is called the **Taylor series** of f at c .

The Taylor series of a function f at $c = 0$ is known as **Maclaurin's series**.

Example: The Maclaurin series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n-1}}{(2n-1)!}$ of $\sin x$ converges to $\sin x$ for all $x \in \mathbb{R}$.

Example: The Maclaurin series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n-1}}{(2n-1)!}$ of $\sin x$ converges to $\sin x$ for all $x \in \mathbb{R}$.

If $f(x) = \sin x$ for all $x \in \mathbb{R}$, then $f : \mathbb{R} \rightarrow \mathbb{R}$ is infinitely differentiable and $f^{(2n-1)}(x) = (-1)^{n+1} \cos x$,
 $f^{(2n)}(x) = (-1)^n \sin x$ for all $x \in \mathbb{R}$ and for all $n \in \mathbb{N}$.

Example: The Maclaurin series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n-1}}{(2n-1)!}$ of $\sin x$ converges to $\sin x$ for all $x \in \mathbb{R}$.

If $f(x) = \sin x$ for all $x \in \mathbb{R}$, then $f : \mathbb{R} \rightarrow \mathbb{R}$ is infinitely differentiable and $f^{(2n-1)}(x) = (-1)^{n+1} \cos x$, $f^{(2n)}(x) = (-1)^n \sin x$ for all $x \in \mathbb{R}$ and for all $n \in \mathbb{N}$.

For $x = 0$, the Maclaurin series of $\sin x$ becomes $0 - 0 + 0 - \dots$, which clearly converges to $\sin 0 = 0$.

Example: The Maclaurin series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n-1}}{(2n-1)!}$ of $\sin x$ converges to $\sin x$ for all $x \in \mathbb{R}$.

If $f(x) = \sin x$ for all $x \in \mathbb{R}$, then $f : \mathbb{R} \rightarrow \mathbb{R}$ is infinitely differentiable and $f^{(2n-1)}(x) = (-1)^{n+1} \cos x$, $f^{(2n)}(x) = (-1)^n \sin x$ for all $x \in \mathbb{R}$ and for all $n \in \mathbb{N}$.

For $x = 0$, the Maclaurin series of $\sin x$ becomes $0 - 0 + 0 - \dots$, which clearly converges to $\sin 0 = 0$.

Let $x (\neq 0) \in \mathbb{R}$. The remainder term in the Taylor expansion of $\sin x$ about the point 0 is given by $R_n(x) = \frac{x^{n+1}}{(n+1)!} f^{(n+1)}(c_n)$, where c_n lies between 0 and x .

Example: The Maclaurin series $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^{2n-1}}{(2n-1)!}$ of $\sin x$ converges to $\sin x$ for all $x \in \mathbb{R}$.

If $f(x) = \sin x$ for all $x \in \mathbb{R}$, then $f : \mathbb{R} \rightarrow \mathbb{R}$ is infinitely differentiable and $f^{(2n-1)}(x) = (-1)^{n+1} \cos x$, $f^{(2n)}(x) = (-1)^n \sin x$ for all $x \in \mathbb{R}$ and for all $n \in \mathbb{N}$.

For $x = 0$, the Maclaurin series of $\sin x$ becomes $0 - 0 + 0 - \dots$, which clearly converges to $\sin 0 = 0$.

Let $x (\neq 0) \in \mathbb{R}$. The remainder term in the Taylor expansion of $\sin x$ about the point 0 is given by $R_n(x) = \frac{x^{n+1}}{(n+1)!} f^{(n+1)}(c_n)$, where c_n lies between 0 and x .

It follows that $\lim_{n \rightarrow \infty} R_n(x) = 0$.

Example: The Maclaurin series $1 + \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$ of $\cos x$ converges to $\cos x$ for all $x \in \mathbb{R}$.

Example: The Maclaurin series $1 + \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$ of $\cos x$ converges to $\cos x$ for all $x \in \mathbb{R}$.

Example: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Example: The Maclaurin series $1 + \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$ of $\cos x$ converges to $\cos x$ for all $x \in \mathbb{R}$.

Example: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Then f is differentiable any number of times, and that $f^{(n)}(0) = 0$ for every $n \in \mathbb{N}$. But the remainder term $R_n(c)$ does not converge to 0 for any $c \neq 0$.

Example: The Maclaurin series $1 + \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$ of $\cos x$ converges to $\cos x$ for all $x \in \mathbb{R}$.

Example: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$f(x) = \begin{cases} e^{-1/x^2} & \text{if } x \neq 0, \\ 0 & \text{if } x = 0. \end{cases}$$

Then f is differentiable any number of times, and that $f^{(n)}(0) = 0$ for every $n \in \mathbb{N}$. But the remainder term $R_n(c)$ does not converge to 0 for any $c \neq 0$.

Thus, an infinitely differentiable function may not have Taylor series representation.