

MAPPING OF GEOLOGICAL FRACTURES: A CNN APPROACH


AYOUB FATIHI AND ANINDITA SAMSU - INSTITUTE OF EARTH SCIENCES, UNIVERSITY OF LAUSANNE, 1015 LAUSANNE, SWITZERLAND

ayoub.fatihi@unil.ch

INTRODUCTION

- Natural fractures can create permeable fluid pathways that are important for geo-energy applications such as CO₂ sequestration and geothermal projects [1].
- Drone and LiDAR-based acquisition methods produce high-resolution 2D data, suitable for detailed fracture mapping [2].
- Manual and semi-automated fracture mapping is time-consuming and susceptible to interpreter bias [3].

Interactive results ▼




Unil
UNIL | Université de Lausanne




Can we use deep learning (Convolutional Neural Networks) to automate 2D fracture mapping for quick and accurate results?

Automatic: No manual intervention required;
Accurate: Results closely match real fractures and unaffected by interpreter bias.



Orthophoto of a section of Orregrund Island in Southwest Finland [3]

METHODS



MODELS:

All based on convolutions

- U-Net [4]
- DeepLabV3+ [5]
- PAN [6]

Post-processing

- Thin
- Smooth
- Vectorize

RESULTS AND DISCUSSION

Results

- The model generates more segments, with some being very short (noise)
- The model's trace length distribution has a lower cut-off value
- Rose diagrams have similar directional pattern

- Portion of the test dataset (OGI) [3]
- Manual mapping** (geologists' interpretations): Ground truth [7] and its respective fracture trace length distribution and rose diagram created using *fractopo* [8]
- Automated mapping:** Predictions of the PAN model and its respective fracture trace length distribution and rose diagram [8]

