REPUBLIC OF CAMEROON
Peace – Work – Fatherland
MINISTRY OF HIGHER EDUCATION
THE UNIVERSITY OF DOUALA
NATIONAL HIGHER POLYTECHNIC SCHOOL OF DOUALA

RÉPUBLIQUE DU CAMEROUN
Paix – Travail – Patrie
MINISTÈRE DE L'ENSEIGNEMENT SUPÉRIEUR
UNIVERSITÉ DE DOUALA
ÉCOLE NATIONALE SUPÉRIEURE POLYTECHNIQUE DE DOUALA

Laboratory of Computer Engineering, Data Science and Artificial Intelligence (LCEDSAI)

MASTER'S THESIS

Design of a Telemonitoring System for Patients with Chronic Diseases (Hypertension, Diabetes) Using Consultation Data, Home Measurements and Possibly Mobile Sensors

Presented by
Nsame Ayuni Rodrick

In partial fulfilment of the requirements for the degree of
Master of Science in Data Science and Artificial Intelligence

Supervisor
Dr. Malong Yannick

Co-supervisor
Dr. Noulapeu Ngaffo

Academic year 2025–2026

Douala, October 29, 2026

Declaration

I, **Nsame Ayuni Rodrick**, declare that this Master's thesis entitled *"Design of a Telemonitoring System for Patients with Chronic Diseases (Hypertension, Diabetes) Using Consultation Data, Home Measurements and Possibly Mobile Sensors"* is the result of my own research, carried out under the supervision of Dr. Malong Yannick and Dr. Noulapeu Ngaffo at the National Higher Polytechnic School of Douala (NHPSD), University of Douala, Cameroon.

I declare that, to the best of my knowledge, this work has not been submitted, in whole or in part, for any other degree, diploma or certification at any other institution.

I further declare that all sources of information, ideas and data used in this thesis have been acknowledged and referenced.

Signed: _____

Date: _____

Place: _____

Abstract

Hypertension and type 2 diabetes are common in Cameroon, and the follow-up of patients between consultations is fragmented. This thesis designs a telemonitoring system for patients with either or both conditions and evaluates its predictive component. The system comprises a patient application and a clinician view built with Flutter on Firebase, a guideline rule that grades every reading on four levels offline on the phone, a two-stage home triage in which a single elevated reading first prompts repeated measurements, and a server worker for escalation, reminders and notifications; it passes 219 automated tests. The predictive component forecasts the risk level of the next reading, so that the label, derived from official thresholds, never enters the inputs, and the guideline rule applied to the last reading (persistence) is the baseline. Eight model families were compared in a nested cross-validation grouped by hospital on the eICU demonstration database (4 182 haemodynamic and 1 356 glycaemic samples, 174 hospitals), and validated externally on 366 readings of 32 inpatients of a public hospital in Douala. On blood pressure, five of the eight families forecast significantly better than persistence; the deployed CatBoost model lowers the ranked probability score from 0.1384 to 0.1341 ($p = 0.002$), a gain of about 3%. On glucose, no family does so significantly (GRU-D 0.1275 against 0.1329, $p = 0.14$). Deep temporal models did not outperform tree ensembles. On the Douala patients, every model lost accuracy and under-predicted the alert; updating the intercept corrected the offset but not the slopes, and no model improved the alert decisions over the rule. Measures of the change in the patient's state discriminated weakly (AUROC at most 0.60). These results support a design in which the guideline rule is the floor and a model may only raise the alert level, and they call for revalidation on home data before any clinical use.

Keywords: telemonitoring; hypertension; type 2 diabetes; risk prediction; external validation; mobile health; Cameroon.

Résumé

L'hypertension artérielle et le diabète de type 2 sont fréquents au Cameroun, et le suivi des patients entre deux consultations est fragmenté. Ce mémoire conçoit un système de télésurveillance pour les patients atteints de l'une ou des deux affections et évalue sa composante prédictive. Le système comprend une application patient et une vue clinicien développées avec Flutter sur Firebase, une règle issue des recommandations qui classe chaque mesure en quatre niveaux, hors connexion, sur le téléphone, un triage à domicile en deux étapes dans lequel une mesure élevée isolée déclenche d'abord des mesures répétées, et un service serveur pour l'escalade, les rappels et les notifications ; il passe 219 tests automatisés. La composante prédictive prévoit le niveau de risque de la mesure suivante, de sorte que l'étiquette, dérivée de seuils officiels, n'entre jamais dans les variables d'entrée, et la règle appliquée à la dernière mesure (persistance) sert de référence. Huit familles de modèles ont été comparées par validation croisée imbriquée, groupée par hôpital, sur la base de démonstration eICU (4 182 échantillons hémodynamiques et 1 356 glycémiques, 174 hôpitaux), puis validées sur 366 mesures de 32 patients hospitalisés dans un hôpital public de Douala. Pour la pression artérielle, cinq familles sur huit prévoient significativement mieux que la persistance ; le modèle CatBoost retenu abaisse le score de probabilité ordonné (RPS) de 0,1384 à 0,1341 ($p = 0{,}002$), soit un gain d'environ 3 %. Pour la glycémie, aucune famille n'y parvient de façon significative (GRU-D 0,1275 contre 0,1329, $p = 0{,}14$). Les modèles temporels profonds n'ont pas surpassé les ensembles d'arbres. Sur les patients de Douala, tous les modèles ont perdu en exactitude et sous-estimé l'alerte ; la mise à jour de l'intercept a corrigé ce décalage mais pas les pentes, et aucun modèle n'a amélioré les décisions d'alerte par rapport à la règle. Les mesures de l'évolution de l'état du patient discriminent faiblement (AUROC au plus 0,60). Ces résultats confortent une conception dans laquelle la règle fixe le niveau minimal et un modèle ne peut que relever l'alerte, et ils imposent une revalidation sur des données recueillies à domicile avant tout usage clinique.

Mots-clés : télésurveillance ; hypertension artérielle ; diabète de type 2 ; prédiction du risque ; validation externe ; santé mobile ; Cameroun.

Contents

Declaration i

Abstract ii

Résumé iii

Contents iv

List of Acronyms viii

List of Figures ix

List of Tables xii

Introduction 1

1 Theoretical Foundations and Literature Review 5

1.1 Key Concepts and Conceptual Framework 5

1.2 Burden of Hypertension and Diabetes 6

1.2.1 The care cascade 6

1.2.2 The hypertension–diabetes comorbidity 7

1.2.3 Why one reading is not enough 8

1.3 Hypertension Care and Digital Health in Sub-Saharan Africa 9

1.3.1 Why the episodic model fails 9

1.3.2 Telemedicine evidence from Cameroon 9

1.3.3 Digital health landscape and barriers 10

1.4 Mobile Health and Clinical Evidence . . . 10
1.4.1 A randomised trial in the target population . . . 11
1.5 Devices, Sensors and Architectures . . . 12
1.5.1 The normative position on devices . . . 12
1.5.2 IoT architectures for remote monitoring . . . 12
1.6 Machine Learning for Clinical Risk Prediction . . . 12
1.7 Data Privacy and Ethics . . . 13
1.8 Critical Comparison of the Approaches . . . 14
1.9 Observed Gaps in the Literature . . . 14

2 Risk-Tiering Framework 16

2.1 Patient Series and Samples . . . 16
2.2 A Label from Official Thresholds . . . 17
2.3 The Forecasting Task . . . 18
2.4 The Decision Layer . . . 19
2.4.1 Three tiers and the rule floor . . . 19
2.4.2 The home triage . . . 19
2.5 Evolution of the Patient's State . . . 20

3 Methodology and System Implementation 22

3.1 Type of Research and Study Design . . . 22
3.2 Data Sources and Collection . . . 23
3.2.1 The three sources . . . 24
3.2.2 Sample size and the role of each dataset . . . 24
3.2.3 Harmonisation of the intensive-care records . . . 25
3.3 Data Curation and Preprocessing . . . 26
3.3.1 A two-phase pipeline . . . 26
3.3.2 What the readings say about the instruments . . . 26
3.3.3 Quality controls at entry . . . 27
3.3.4 Outliers: artefact or event? . . . 27
3.3.5 Missing data . . . 27
3.4 Features . . . 28
3.5 Candidate Models and Baselines . . . 29

3.6 Experimental Protocol . . . 29
3.7 Evaluation Metrics . . . 31
3.8 The Telemonitoring System . . . 32
3.8.1 Architecture . . . 32
3.8.2 Data model and rules . . . 32
3.8.3 Server functions . . . 33
3.8.4 Interfaces . . . 33
3.9 Implementation Details and Reproducibility . . . 34
3.10 Ethical Considerations . . . 35

4 Results and Discussion **37**

4.1 Data Collection and Treatment . . . 37
4.1.1 The Douala charts . . . 38
4.1.2 What the readings say about the instruments . . . 38
4.1.3 Labels . . . 39
4.1.4 The development data . . . 40
4.1.5 How often the level changes . . . 40
4.1.6 Sample size . . . 41
4.2 Benchmark of the Risk Models . . . 42
4.2.1 Learned models against the guideline rule . . . 42
4.2.2 Deep temporal against tabular models . . . 44
4.2.3 The deployed model and the stability of the choice . . . 44
4.2.4 What the haemodynamic model uses . . . 45
4.3 External Validation on the Douala Patients . . . 45
4.4 Evolution of the Patient's State . . . 48
4.5 System Verification . . . 48
4.6 Discussion . . . 49
4.6.1 Answers to the research questions . . . 49
4.6.2 What the data imply for the transfer . . . 50
4.6.3 Comparison with the literature . . . 51
4.6.4 Limitations . . . 51

General Conclusion and Perspectives **53**

References 56

A Data Dictionary 61

B Benchmark Configuration 63

C Record Schema 65

D Application Screens 67

E TRIPOD+AI Checklist 68

List of Acronyms

ADA American Diabetes Association
AI Artificial intelligence
AUC Area under the curve
AUROC Area under the ROC curve
BP Blood pressure
CI Confidence interval
CPU, GPU Central, graphics processing unit
DBP Diastolic blood pressure
ECG Electrocardiogram
eICU eICU Collaborative Research Database
ENSPD École Nationale Supérieure Polytechnique de Douala (NHPSD)
ESH European Society of Hypertension
GRU-D Gated recurrent unit with decay
HbA1c Glycated haemoglobin
HEARTS WHO technical package for cardiovascular care
IHSG International Hypoglycaemia Study Group
IoT Internet of Things
LASSO Least absolute shrinkage and selection operator
LSTM Long short-term memory
MAD Median absolute deviation
MAR, MCAR, MNAR Missing at random, completely at random, not at random
mHealth Mobile health
MQTT Message Queuing Telemetry Transport
NEWS2 National Early Warning Score 2
NHPSD National Higher Polytechnic School of Douala
PCA Principal component analysis
PPG Photoplethysmography
PRISMA Preferred Reporting Items for Systematic Reviews and Meta-Analyses
PROBAST Prediction model Risk Of Bias Assessment Tool
RCT Randomised controlled trial
ROC Receiver operating characteristic
RPS Ranked probability score
RQ Research question
SBP Systolic blood pressure
SD Standard deviation
SDK Software development kit
SHAP Shapley additive explanations
SMS Short message service
STEPS WHO STEPwise approach to surveillance
TPE Tree-structured Parzen estimator
TRIPOD Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis
WHO World Health Organization

List of Figures

1 Organisation of the thesis and where each research question is answered. 4

1.1 Conceptual framework of the thesis. 5

1.2 Estimates and 95% intervals of Table 1.4; the dashed line marks the pooled African estimate. The Central African interval spans 38 points. 8

1.3 Estimates of Table 1.8 with their 95% intervals. The only interval that includes no effect is that of the trials conducted in developing countries, none of which involved professional management. 11

2.1 Construction of one sample: the readings of the window, the embargo and the first reading after it. 16

2.2 Level of a single reading for systolic pressure and capillary glucose (Table 2.1). Diastolic pressure and pulse follow the same pattern with their own thresholds. 17

2.3 From a value forecast to level probabilities, for one channel. The forecast distribution is cut by the thresholds of Table 2.1; each area is the probability of a level. Within a condition, the channels are then combined by their worst level. 18

2.4 The decision layer. The model can raise the tier given by the guideline rule, never lower it; the cut-points 0.1 and 0.3 are provisional. 19

2.5 Path of a home reading through the two-stage triage. Only the mean of a verification profile, or a red reading, reaches a clinical decision. 20

2.6 The two windows compared for one channel. Each window is summarised by its mean; the four channel means form the state vectors **c** and **b**. 21

3.1 Study design. The same treatment, labels and samples apply to both datasets. . 23

3.2 From the paper records of the Douala hospital to the analysis data. No name leaves the hospital; the slot is morning or evening, the hour is not recorded. . . 24

3.3 The 366 readings of the Douala charts. One row per patient, one mark per reading, coloured by level (grey stable, blue moderate, light red high, red critical); open marks are readings without glucose. Patients are numbered by date of first reading. 25

3.4 Harmonisation of an intensive-care stay to the rhythm of the ward charts. At each slot, the reading nearest in time within two hours is kept for each channel (red); a slot with none is missing, as an unrecorded value on a chart would be. 26

3.5 The two-phase preprocessing pipeline. 26

3.6 Search and selection in one outer fold. The outer test hospitals are used once, after every choice has been made on the inner folds. 30

3.7 Nested cross-validation on the development data, repeated three times. 30

3.8 Architecture of the built system. Dashed: push notifications sent by the worker to the two applications. 32

3.9 Data model of the record in Cloud Firestore. Readings, events and messages are subcollections of the patient's document, so one security rule governs each of them. 33

3.10 Life of an alert. The phone warns the patient without a network; the record, the checks, the notifications and the escalation do not depend on the phone staying online. 34

4.1 Flow of the data from the two sources to the forecast samples. 37

4.2 Terminal digits of the Douala readings. The dashed line marks the uniform share of 0.1. 39

4.3 Levels at the forecast targets. Douala: 282 haemodynamic and 190 glycaemic targets; eICU demonstration subset: 4 182 and 1 356. 39

4.4 Distribution of the readings in the two datasets, with the thresholds of Table 2.1 on the axes. Douala: 356 systolic, 355 diastolic and 251 glucose readings; eICU: 6 577, 6 563 and 2 444. 40

4.5 Level of the target (columns) given the level of the anchor reading (rows), in percent of each row: S stable, M moderate, H high, C critical. The diagonal is what persistence predicts. Pairs where both readings carry the label: 282 and 4 182 haemodynamic, 182 and 894 glycaemic. 41

4.6 Parameters allowed by the criteria of Riley et al. with the effective sample size, against the anticipated C statistic. 41

4.7 Difference in RPS between each candidate and persistence in the 15 outer folds (dots) and its mean (red bar). Below the dashed line, the candidate forecasts better than the guideline rule applied to the last reading. 43

4.8 RPS in development (blue squares: mean of the 15 outer folds) and on the Douala charts (red: estimate and 95% patient-bootstrap interval). Lower is better. . . . 47

4.9 Decision curves on the Douala charts, probabilities as is. The net benefit counts true alerts minus false alarms weighted by the odds of the threshold. 47

4.10 AUROC of the measures of state evolution with 95% patient-level intervals. Dashed: chance. The measure is chosen on eICU only. 48

4.11 Precision against the share of readings that need attention, at a sensitivity of 0.90. The dashed line marks a precision of 80%. Dotted lines: the planning prevalence of 0.02 for a home deployment, and the prevalence of the haemodynamic alert at the targets of eICU (0.17) and Douala (0.23). 51

D.1 Screens of the application: (a) sign-in, shared by patients and clinicians; (b) enrolment with the clinician's code; (c) patient home with the level, the instruction and the verification profiles; (d) entry of the two readings of the protocol; (e) correction, a new reading that supersedes the old one; (f) history with level and corrections; (g) clinician queue, red before yellow; (h) nearby care, opened in the phone's maps. Rendered by the widget tests with fictitious patients; model names and probabilities are test values. 67

List of Tables

1.1 Sources of the review, by theme and in chronological order within each theme. . 6

1.2 Key concepts and the definitions adopted in this thesis. 7

1.3 Hypertension care cascade by WHO region, adults aged 30–79 years, 2019 [1]. 7

1.4 Sub-regional pooled prevalence of hypertension among people with diabetes in Africa [3]. 7

1.5 Most recent hypertension surveys in Central Africa (adapted from [2], Table 1). 9

1.6 The two Cameroonian evaluations of cardiovascular telemedicine. 10

1.7 Six dimensions of barriers to telemedicine in sub-Saharan Africa [11]. 10

1.8 Estimates of the effect of mobile health and telemonitoring on systolic pressure and glycaemic control. 11

1.9 How the reviewed machine learning studies were evaluated, against the design of this work. 13

1.10 Comparison of the approaches found in the literature. 14

1.11 Synthesis of the literature review. 15

2.1 Severity of a single reading, per channel. Pressure in mmHg, pulse in beats per minute, capillary glucose in mmol/L. 17

2.2 The three-level home triage of Mihevc et al. [8]. Pressure in mmHg, glucose in mmol/L. 20

2.3 Candidate measures of the change between current state and baseline, from the simplest. σ_j: between-patient SD of channel j; σ_{ij}: SD of patient i, shrunk towards the pooled within-patient SD; Σ: pooled within-patient covariance of changes. . 21

3.1 The three data sources and their role. 23

3.2 Quality controls applied at entry. 27

3.3 Preparation required by model family, fitted on the training fold. 28

3.4 Window length and the features it can support, at two readings a day. 28
3.5 Feature sets, fixed before any result. 29
3.6 Candidate families and baselines. 29
3.7 Admissible performance measures. 31
3.8 Server functions run by the worker. 33
3.9 Software and versions used for the analysis and the system. 35

4.1 The Douala charts after the quality controls of Table 3.2. 38
4.2 Levels of the 366 Douala readings under the earlier rule (rows) and the sourced rule (columns). 40
4.3 Effect of the slot tolerance on the eICU development samples. 41
4.4 Sample size against the criteria of Riley et al. Parameters for an anticipated C statistic of 0.70 to 0.80. 42
4.5 Outer-fold results, haemodynamic task: mean (SD) over 15 outer folds. ΔRPS: difference with persistence (negative is better); p: corrected resampled t test. Net benefit averaged over the thresholds 0.1 to 0.3. 42
4.6 Outer-fold results, glycaemic task: mean (SD) over 15 outer folds. ΔRPS: difference with persistence (negative is better); p: corrected resampled t test. Net benefit averaged over the thresholds 0.1 to 0.3. 43
4.7 Outer-fold results by subgroup, haemodynamic task: persistence against the deployed family (CatBoost). Each repeat pools its five outer folds; mean of the three repeats; n: targets per repeat. In 36 of the 49 hospitals with at least 30 targets, CatBoost has a lower RPS than persistence. 44
4.8 Outer-fold results by subgroup, glycaemic task: persistence against the deployed family (GRU-D). Each repeat pools its five outer folds; mean of the three repeats; n: targets per repeat. In 5 of the 7 hospitals with at least 30 targets, GRU-D has a lower RPS than persistence. 44
4.9 Stability of the choice of family, final search on all eICU: selection cost with equal weights (lower is better), and share of 1 000 random weightings (uniform Dirichlet) under which the family has the lowest cost. 45
4.10 Grouped permutation importance of the deployed model, haemodynamic task (CatBoost): increase in RPS when the features of one group are permuted together, mean (SD) of 20 permutations. 45
4.11 External validation on the Douala charts, haemodynamic task (282 targets, 32 patients). 95% intervals by patient bootstrap. Calibration intercept and slope as is, then after the intercept update. 46
4.12 External validation on the Douala charts, glycaemic task (190 targets, 24 patients). 95% intervals by patient bootstrap. Calibration intercept and slope as is, then after the intercept update. 46

4.13 Verification of the system. 49
4.14 Limitations, their consequences and the response of this work. 52

A.1 Columns of the Douala charts file. 61
A.2 Tables and columns of eICU used. 62

B.1 Search spaces and number of configurations. Ranges marked log are searched on a logarithmic scale. 63
B.2 Scripts of the research package and what they produce. 64

C.1 Fields of a reading, `users/{uid}/readings/{id}`. 65
C.2 Fields of an event, `users/{uid}/events/{id}`. 66

E.1 TRIPOD+AI items and where they are addressed. 68

Introduction

Hypertension is the leading modifiable risk factor for premature death worldwide. The number of adults aged 30–79 years living with hypertension doubled between 1990 and 2019 to reach 1.3 billion, about 78% of whom live in low- and middle-income countries, and elevated systolic blood pressure causes an estimated 10.8 million deaths each year [1]. Sub-Saharan Africa combines a high and rising prevalence with poor detection, treatment and control [2]: in the WHO African Region only 12% of hypertensive adults have their blood pressure controlled. Diabetes compounds the burden. In Africa, 58% of people living with diabetes are also hypertensive [3]; the two conditions share their risk factors, aggravate each other, and call for the same care: repeated measurement and treatment adjusted over decades.

The Follow-up Challenge

Care for hypertension and diabetes in Cameroon is episodic. Blood pressure and glucose are measured at consultations a few times a year, recorded on paper, and rarely assembled into a history that a clinician can read as a trend. Yet diagnosis and treatment adjustment are longitudinal acts: the WHO protocol requires measurements repeated over several visits and monthly follow-up until the target is reached [1, 4]. Two failures follow. Loss to follow-up affects half of patients or more in resource-limited settings, and therapeutic inertia leaves treatment unchanged while blood pressure stays above target [1]. In the African Region, 72% of the patients already diagnosed are not controlled: they are known to the health system and lost during follow-up.

The Potential of Telemonitoring and Prediction

Mobile telephony now reaches most of the population [5], and a controlled study in Cameroonian district hospitals has shown that remote clinical support is feasible in primary care [6]. Telemonitoring, the transmission of home measurements to a care team, lowers systolic pressure by 3 to 7 mmHg and glycated haemoglobin by 0.4 to 0.5 points, provided a clinician acts on the data [7, 8]. Where clinicians are scarce, the system must also decide which readings deserve their time. The only validated protocol for patients with both conditions sorts readings into three levels with fixed thresholds [8]: it reacts to the level of a reading, not to its trend, and uses no learned model. Machine learning could forecast deterioration from a patient's own series, but the reported results of the field are fragile. The same algorithm is reported as the best on one dataset [9] and as no better than chance on another [10], because the first study predicts a label already encoded in its inputs.

Research Problem and Motivation

This thesis addresses two problems.

1. **A practical problem.** No telemonitoring system evaluated in Central Africa combines consultation records, home measurements and a predictive model for patients with hypertension and diabetes, and orders the clinician's attention under intermittent connectivity and scarce specialist time [11, 12].
2. **A methodological problem.** The predictive results of the field rest on labels that leak into the features, improper metrics and validation on the training population. Whether a learned model forecasts deterioration better than the guideline rule, and whether a model developed elsewhere transfers to Cameroonian patients, has not been measured.

The data available to this work were not collected at home. The public eICU Collaborative Research Database, which records intensive-care stays in United States hospitals, is used to develop and compare the models. The observation charts of 32 patients hospitalised in Douala, with blood pressure, pulse and glucose recorded morning and evening, are used only to validate them externally. No home measurement or sensor data were collected: the system is built for home follow-up, but its predictive component is evaluated on hospital data, and its transfer to home data remains to be established.

Objectives and Research Questions

The general objective is to design a telemonitoring system for patients with hypertension and type 2 diabetes that supports the clinician through risk assessment and alert prioritisation, and to evaluate its predictive component without the biases of previous work. The specific objectives are:

- to derive the requirements of the system and the gaps of the field from the literature (Chapter 1);
- to formalise the risk-tiering problem: a sourced labelling rule, a forecasting task free of label leakage, and the combination of rule and model (Chapter 2);

- to build a documented pipeline of data collection and treatment, and a benchmark protocol that selects the model instead of presupposing it (Chapter 3);
- to compare the candidate models on public multi-centre data, validate them externally on Cameroonian patients, and build the system around the result (Chapter 4).

Four research questions follow:

RQ1. Can a telemonitoring system combining consultation records, home readings and risk prediction be built under the connectivity, workload and cost constraints of Cameroonian care?

RQ2. Do learned models forecast the risk level of the next reading better than the guideline rule, and does a deep temporal model outperform classical models?

RQ3. Does the advantage persist when the forecast is made 24 and 48 hours ahead?

RQ4. Do models developed on another health system transfer to Cameroonian patients?

Hypotheses

RQ1 is a design question, answered by the system and by the traceability of each of its elements to a requirement. RQ2 to RQ4 lead to four testable hypotheses, stated with the measures of Section 3.7:

H1. At least one learned model forecasts the level of the next reading with a lower ranked probability score than the guideline rule applied to the last reading (persistence).

H2. This advantage persists at lead times of 24 and 48 hours.

H3. The deep temporal models do not outperform the tabular models.

H4. Performance is lower on the Douala patients than on the development data, and updating the intercept on the local data recovers part of the calibration.

Contributions and Thesis Outline

This work contributes:

- **A data collection and treatment pipeline**: a labelling rule in which every threshold comes from an official source, quality controls that tell artefacts from events, an analysis of the instruments from the readings themselves, and the harmonisation of intensive-care records to the rhythm of ward charts.
- **A benchmark protocol**: candidates taken from published benchmarks rather than chosen in advance, nested validation grouped by hospital, proper scoring rules combined in a declared selection cost, and external validation on Cameroonian patients. The same discipline selects the measure that tracks the evolution of a patient's state.
- **A telemonitoring system**: a patient application and a clinician dashboard implementing the three-level home protocol, a local risk service sharing the code of the benchmark, and server functions for escalation, reminders and verdicts.




Figure 1 Organisation of the thesis and where each research question is answered.

Figure 1 summarises the organisation. Chapter 1 reviews the clinical and technical literature and identifies the gaps. Chapter 2 formalises the risk-tiering problem. Chapter 3 describes the data, their treatment, the benchmark protocol and the system. Chapter 4 reports the results and discusses them, and the Conclusion summarises the findings and the directions for future work. Every number of the results is produced by a script of the accompanying code, which can be rerun.

CHAPTER 1

Theoretical Foundations and Literature Review

This chapter establishes what is known about the burden of hypertension and diabetes and the failure of their care, the clinical evidence for telemonitoring, the instruments and architectures available, and the use of machine learning for clinical risk. It draws on nineteen peer-reviewed sources, each checked against PubMed, Crossref or the publisher's record, complemented by the guidelines of the World Health Organization (WHO) [1, 4, 13] and of the European Society of Hypertension (ESH) [14]. Each study is reported with its method, its main result and its limits, in six themes (Table 1.1); the chapter ends with a comparison of the approaches and the gaps that motivate this work.

1.1 Key Concepts and Conceptual Framework

The terms of Table 1.2 are used throughout the thesis with the meaning given here; several are used differently in the literature.

Four models structure the reasoning (Figure 1.1). The **care cascade** locates where an intervention acts: screening acts on diagnosis, follow-up on treatment and control, where most diagnosed patients in Africa are lost. The **measurement model** treats a reading as the usual pressure plus an error that repeated readings reduce, which justifies series over isolated values. The **layered architecture** describes a monitoring system as a chain from acquisition to processing and presentation [20, 25]. And **supervised learning for clinical prediction** fixes what a valid evaluation requires: a label outside the input window, proper measures, and a comparison with the guideline rule.




Figure 1.1 Conceptual framework of the thesis.

Table 1.1 Sources of the review, by theme and in chronological order within each theme.

Theme	Source	Design and setting
1. Burden of disease	Hinneh et al. [3]	systematic review and meta-analysis, Africa
	World Health Organization [1]	institutional global report
2. Care and digital health in sub-Saharan Africa	Kingue et al. [15]	interventional pilot, rural Cameroon
	Bediang et al. [6]	controlled multicentre study, Cameroon
	Dodoo, Al-Samarraie, and Alsswey [11]	review of 66 empirical studies, sub-Saharan Africa
	Gafane-Matemane et al. [2]	narrative review, sub-Saharan Africa
	Agbeyangi and Lukose [12]	systematic review, South Africa, Kenya, Nigeria
	Segun-Omosehin et al. [5]	perspective, sub-Saharan Africa
3. Mobile health and clinical evidence	Alessa et al. [16]	review and content analysis of applications, United Kingdom
	Mao et al. [7]	meta-analysis of 51 randomised trials
	Zhou et al. [17]	meta-analysis of randomised trials
	Bandeira et al. [18]	updated meta-analysis of randomised trials
	Mihevc et al. [8]	multicentre randomised trial, Slovenia
4. Sensors and IoT	Alhaddad et al. [19]	review of non-invasive glucose sensing
	Tai et al. [20]	IoT architecture and deep learning technique
	Efendi et al. [9]	IoT and cloud implementation with device validation
	Liu and Wang [21]	review of IoT sensing and machine learning
	Liu et al. [22]	smartphone photoplethysmography study
5. Machine learning	Chatrati et al. [23]	home monitoring system with machine learning
	Mendoza-Cano et al. [10]	cross-sectional predictive study, Mexico
6. Privacy and ethics	Alhammad et al. [24]	systematic review of patient perspectives

1.2 Burden of Hypertension and Diabetes

1.2.1 The care cascade

For population surveillance, the WHO defines hypertension as a systolic blood pressure (SBP) ≥ 140 mmHg, a diastolic blood pressure (DBP) ≥ 90 mmHg, or the use of antihypertensive medication [1]; the ESH uses the same threshold and divides hypertension into three grades [14]. The number of hypertensive adults rose from 650 million in 1990 to 1.3 billion in 2019, while the age-standardised prevalence barely changed, from 32% to 33%: population growth and ageing, not individual risk, drive the increase. Elevated systolic pressure is the leading risk factor for premature death among the 87 compared.

The WHO describes care as a cascade of nested stages measured on the same denominator (Table 1.3). The European Region has a higher prevalence than the African Region but a control rate more than twice as high: the African deficit is organisational rather than epidemiological. In the African Region, 57% of hypertensive adults are undiagnosed, 37% of the diagnosed are untreated and 56% of the treated are not controlled; among patients already diagnosed, 72% are not controlled. Screening acts on the first link of the chain; longitudinal follow-up acts on the second and third.

Table 1.2 Key concepts and the definitions adopted in this thesis.

Concept	Definition adopted
Hypertension	systolic pressure ≥ 140 mmHg, diastolic pressure ≥ 90 mmHg, or antihypertensive treatment [1]; ESH grades from 140/90, 160/100 and 180/110 mmHg [14]; home threshold 135/85 mmHg
Type 2 diabetes	chronic hyperglycaemia due to insulin resistance and relative insulin deficiency; its follow-up requires a biological measurement (capillary or venous glucose, glycated haemoglobin)
Care cascade	nested stages of care—hypertensive, diagnosed, treated, controlled—measured on the same population [1]
Loss to follow-up	interruption of the planned follow-up of a patient already in care
Therapeutic inertia	failure to intensify treatment while blood pressure remains above target
Telemonitoring	transmission of physiological data from the patient's home to a care team, which provides clinical feedback
Mobile health (mHealth)	health practices supported by mobile devices: applications, text messages, connected devices
Self-measurement	measurement of blood pressure or capillary glucose by the patient at home, with a validated device and a standard protocol
Internet of Things (IoT)	connected devices that acquire and transmit data without human intervention
Clinical decision support	system that turns patient data into alerts or recommendations for a clinician, who keeps the decision
Supervised learning	estimation of a function mapping input features to a known label; here, a forecast of a future clinical state from a past window of readings
Label leakage	situation in which the label can be recovered from the inputs, so that measured performance reflects a rule rather than a prediction
Triage	ordering of readings or patients by priority for clinical review

Table 1.3 Hypertension care cascade by WHO region, adults aged 30–79 years, 2019 [1].

Indicator	African Region	World	Americas	Europe
Age-standardised prevalence	36%	33%	35%	37%
Diagnosis coverage	43%	54%	70%	66%
Treatment coverage	27%	42%	60%	53%
Effective treatment (control)	12%	21%	36%	26%

1.2.2 The hypertension–diabetes comorbidity

Diabetes stiffens small vessels, which raises peripheral resistance and blood pressure, and hypertension accelerates the vascular damage of diabetes. Hinneh et al. [3] pooled 41 cross-sectional studies (144 433 participants) following PRISMA: 58.1% of people with diabetes in Africa are also hypertensive (95% CI 52.0–63.2), and 60.8% of those with type 2 diabetes. The risk rises with age, excess weight, years lived with diabetes, urban residence and male sex. The sub-regional estimates place Central Africa highest (Table 1.4 and Figure 1.2).

Table 1.4 Sub-regional pooled prevalence of hypertension among people with diabetes in Africa [3].

Sub-region	Pooled prevalence	95% CI	Studies
Central Africa	77.6%	53.0–91.4	2
Southern Africa	69.1%	59.8–77.1	10
Eastern Africa	53.0%	45.8–59.1	15
Western Africa	51.5%	41.8–61.1	10




Figure 1.2 Estimates and 95% intervals of Table 1.4; the dashed line marks the pooled African estimate. The Central African interval spans 38 points.

The Central African value must be read with its interval. It rests on two small Cameroonian hospital series (109 and 210 patients), while three other Central African studies listed in the article (1 832 patients, prevalences of 35.0–50.4%) do not enter the subgroup; heterogeneity is extreme ($I^2 \approx 98\%$), and the restriction to English-language publications under-represents francophone Central Africa. It cannot carry the case for this work on its own. That case rests on the continental estimate, on the mechanism, and on the WHO position that uncontrolled hypertension is among the risk factors most strongly associated with premature death in people with diabetes [1]. More generally, the prevalences quoted in this chapter differ in population, threshold, number of measurement visits and recruitment; none is nationally representative for Cameroon, so the nature of each estimate is stated where it is used.

1.2.3 Why one reading is not enough

The WHO diagnostic protocol is longitudinal by construction: hypertension is diagnosed only if the threshold is exceeded at two visits [1]. A reading of patient i at time t can be written $b_{i,t} = \mu_i + \varepsilon_{i,t}$, where μ_i is the patient's usual pressure, the clinically relevant quantity, and $\varepsilon_{i,t}$ gathers circadian variation, activity, the measurement setting and instrument error. With n readings far enough apart for their errors to be independent,

$$\operatorname{Var}(\hat{\mu}_i) = \sigma_\varepsilon^2 / n. \tag{1.1}$$

A consultation provides two to four readings a year; home follow-up provides several dozen, which estimate μ_i far more precisely and also estimate the patient's own variability σ_ε. Follow-up must moreover be frequent: Recommendation 7 of the WHO guideline sets monthly follow-up after a treatment change until the target is reached, then every three to six months [4]. With 56% of treated patients uncontrolled, this means about eight contacts per patient per year, where an episodic model delivers one or two.

1.3 Hypertension Care and Digital Health in Sub-Saharan Africa

1.3.1 Why the episodic model fails

Gafane-Matemane et al. [2] synthesise the hypertension literature of sub-Saharan Africa from 2017 to 2023: a high burden with poor detection and control, driven by a westernised lifestyle, and obstacles to care that are individual (literacy, cost, distance) and systemic (medicines, staffing). Their inventory of national surveys is the most recent for Central Africa (Table 1.5): prevalence ranges from 13.5% to 42.1%, reflecting the diversity of instruments as much as real differences, and three countries have no data. In Cameroon, the May Measurement Month campaign of 2019 found a prevalence of 20.8%, with only 11.2% of hypertensive participants controlled.

Table 1.5 Most recent hypertension surveys in Central Africa (adapted from [2], Table 1).

Country	Instrument	Year	*n*	Prev.	Aware.	Treat.	Control
Burundi	hospital screening	2018	353	16.7%	–	–	–
Cameroon	May Measurement Month	2019	30 167	20.8%	29.9%	24.0%	11.2%
Central African Rep.	WHO STEPS	2017	3 265	42.1%	30.2%	–	–
Comoros	national survey	2019	2 602	38.4%	47.9%	45.4%	30.2%
DR Congo	May Measurement Month	2019	29 857	25.5%	33.1%	23.2%	11.9%
Congo (Rep.)	May Measurement Month	2019	3 157	33.5%	42.6%	37.3%	23.3%
Gabon	retrospective study	2013–15	2 104	13.5%	–	–	–
Chad, Eq. Guinea, São Tomé	no data	–	–	–	–	–	–

The WHO documents the mechanisms [1]: loss to follow-up often affects half of patients or more, so screening is useful only when linked to follow-up; therapeutic inertia makes the last link the weakest; lifestyle change alone lowers systolic pressure by only about 1.3 mmHg, so nearly every patient needs medication and its timely adjustment; consultations often last less than five minutes; and paper records are lost or unavailable during the visit.

1.3.2 Telemedicine evidence from Cameroon

Two controlled evaluations of cardiovascular telemedicine have been conducted in Cameroonian primary care (Table 1.6). In TELEMED-CAM, Kingue et al. [15] supported thirty rural health centres from Yaoundé: among 268 participants followed for 24 weeks, attendance at scheduled visits was higher in the intervention arm at every grade of hypertension, but the benefit on control was confined to stage III. Bediang et al. [6] equipped two district hospitals with an electrocardiograph and a tele-expertise platform: access to an electrocardiogram with a cardiologist's opinion reached 98.9% against 33.3%, and the expert modified the local diagnosis in 46.2% of cases. The specialist's opinion, however, took 10.3 days on average against a protocol maximum of 72 hours, no faster than in the control centres: the bottleneck was expert availability, not transmission.

Read together, the two studies support three conclusions. Retention is the most robust result, observed by different mechanisms eight years apart; an effect on blood pressure control is not established in Cameroon; and the value added lies in the clinical decision rather than in a tool

Table 1.6 The two Cameroonian evaluations of cardiovascular telemedicine.

	Kingue et al. [15]	Bediang et al. [6]
Function	telemanagement of treatment	tele-expertise for diagnosis
Design	partly randomised by centre, 268 patients	matched centres, not randomised, 171 patients
Principal result	control improved in stage III only	ECG with expertise, 98.9% vs 33.3%
Retention	visit adherence higher at every grade	52 control patients not seen at 3 months
Main weakness	clustering ignored; effect of a package	non-comparable groups; differential attrition

placed in the patient's hands. Both show that systematic expert review of every case does not scale where specialists are scarce.

1.3.3 Digital health landscape and barriers

Dodoo, Al-Samarraie, and Alsswey [11] reviewed 66 empirical studies of telemedicine in sub-Saharan Africa and grouped the barriers into six dimensions (Table 1.7).

Table 1.7 Six dimensions of barriers to telemedicine in sub-Saharan Africa [11].

Dimension	Definition given by the authors
Technological	information systems and connectivity
Organisational	stakeholder engagement, sustainability of systems
Individual	users' experience, knowledge and awareness
Financial	financial resources and operating cost
Cultural	social constraints and traditional beliefs
Legal and regulatory	policies, legislation, confidentiality and privacy protection

Most projects lost momentum once external funding ended, and health professionals facing a heavy workload perceived telemedicine as an additional task. Agbeyangi and Lukose [12] compared adoption in South Africa, Kenya and Nigeria over 53 studies; Central Africa is absent from the comparison. Segun-Omosehin et al. [5] report that more than 80% of the population holds a mobile subscription, while about 550 million people lack reliable electricity and internet access and up to 18% of countries had no registered cardiologist in 2020; artificial intelligence remains largely unvalidated on African populations. The population is reachable by mobile networks and the specialist is not: a system that forwards every reading reproduces the scarcity it should relieve, whereas one that sorts readings and orders the clinician's attention addresses it.

1.4 Mobile Health and Clinical Evidence

Alessa et al. [16] analysed the 186 hypertension applications of the two main stores in 2018: most offered a single function, none of the thirty analysed in depth had been evaluated scientifically, and they contained a median of nine behaviour change techniques. Mao et al. [7] pooled 51 randomised trials (13 054 participants with diabetes or hypertension): mobile health lowered systolic pressure by 2.99 mmHg and glycated haemoglobin (HbA1c) by 0.39 points. Combined with professional management, it lowered systolic pressure by 6.17 mmHg; alone, by a non-significant 2.16 mmHg; and in developing countries, whose trials used no professional management, it had no effect on systolic pressure. Only one of the 51 trials took place in sub-Saharan Africa. Zhou

et al. [17] found −5.78 mmHg over 16 trials of mobile applications, larger with at least eleven behaviour change techniques, and Bandeira et al. [18] identify telemonitoring with clinician feedback as the most effective modality.

1.4.1 A randomised trial in the target population

Mihevc et al. [8] ran the trial most relevant to this work: 128 patients aged 65 or more with both hypertension and type 2 diabetes, in three Slovenian primary-care centres. The intervention group measured blood pressure and glucose at home with Bluetooth devices transmitting to a cloud platform, and a telemedicine centre reviewed the values according to a three-level protocol (green, yellow, red) adopted in this work (Section 2.4.2). After twelve months, systolic pressure was 6.9 mmHg lower than in the control group and HbA1c 0.5 points lower. The authors credit the effect to contact with the care team and treatment adapted on reliable measurements, not to behaviour change. Seven of the 64 intervention patients (10.9%) withdrew in the first month because they could not use the equipment, and a third of the systolic effect was lost between six and twelve months.

The clinical evidence (Table 1.8) therefore supports a modest but clinically relevant effect, conditional on professional oversight. The evidence from sub-Saharan Africa is almost absent, and the absence of effect in developing countries is plausibly a design gap—no clinician at the end of the chain—rather than a context effect. Telemonitoring must be built as decision support linked to a clinician, not as a self-tracking tool.

Table 1.8 Estimates of the effect of mobile health and telemonitoring on systolic pressure and glycaemic control.

Source	Design	Effect on SBP (95% CI)	HbA1c
Mao et al. [7], overall	30 RCTs on SBP	−2.99 mmHg (−4.19 to −1.80)	−0.39
with professional management	subgroup	−6.17 mmHg (−8.83 to −3.50)	–
developing countries	subgroup	+0.25 mmHg (−3.10 to 3.59), n.s.	−0.52
Zhou et al. [17]	16 RCTs	−5.78 mmHg (−7.97 to −3.59)	–
Mihevc et al. [8]	RCT, 128 patients	−6.9 mmHg (−11 to −2.7)	−0.5




Figure 1.3 Estimates of Table 1.8 with their 95% intervals. The only interval that includes no effect is that of the trials conducted in developing countries, none of which involved professional management.

1.5 Devices, Sensors and Architectures

1.5.1 The normative position on devices

The WHO recommends independently validated automated upper-arm cuffs and does not recommend cuffless devices for managing hypertension, because their accuracy and their ability to track change are unresolved; only about one automated device in five on the market has evidence of validation [1]. Non-invasive glucose sensing by electrocardiography, bioimpedance or photoplethysmography (PPG) shows promise for trends and hypoglycaemia but not for clinical-grade measurement [19], and blood pressure from a smartphone camera correlates with reference devices on 127 participants but remains below cuff accuracy [22]. Blood pressure must therefore come from a validated cuff and glucose from a glucometer; PPG is at most a source of heart rate.

1.5.2 IoT architectures for remote monitoring

Tai et al. [20] embed a deep learning technique for chronic disease monitoring in a three-phase architecture—acquisition, processing, visualisation—evaluated in controlled conditions. Efendi et al. [9] built an Internet of Things (IoT) system for elderly people: an ESP32 microcontroller reads an optical heart-rate and oxygen sensor and a temperature sensor, sends the values over MQTT to a Firebase database with offline persistence, and displays them in a mobile application. Against a digital oximeter and thermometer, six volunteers measured for ten minutes gave a mean absolute percentage error of 0.90%. The system measures neither blood pressure nor glucose. Mucchi et al. [25] decompose remote monitoring into body, environment, edge and cloud layers and tested their system with ten nurses on simulated patients; Yazdani et al. [26] add a priority-ordered clinician queue and a configurable checkup period, without implementing them; Liu and Wang [21] survey IoT sensing with machine learning and describe the defects of electronic records: duplicates, unit errors and values carried over. The architectures converge on acquisition, transmission and storage, processing and presentation; none has been validated on patients in routine care in a low-resource setting.

1.6 Machine Learning for Clinical Risk Prediction

Chatrati et al. [23] proposed a home system that analyses blood pressure and glucose readings with rules for hypertension and a classifier for diabetes; it classifies single readings rather than series. Liu and Wang [21] report accuracies of 95 to 100% for the dominant algorithm families, whereas the only recurrent network applied to a genuine clinical prediction, the readmission of lupus patients, reaches an area under the curve of 0.70. The classifier comparison of Efendi et al. [9] shows why such figures must be read with care. XGBoost reached an accuracy of 0.973, but the label *stable* is defined by the same normal ranges used to discretise the three input features, so the label is a deterministic function of the inputs:

$$y = 1 \iff (x_1 = 1) \land (x_2 = 1) \land (x_3 = 1), \tag{1.2}$$

and the perfectly balanced classes (2 250 against 2 250) show that the dataset was constructed rather than observed. The models recover a rule instead of predicting a state.

Mendoza-Cano et al. [10] provide the counter-example of a genuine prediction: uncontrolled blood pressure among 1 308 hypertensive adults of a nationally representative Mexican survey,

predicted from seven predictors selected by LASSO. Random forest reached an area under the curve of 0.75, logistic regression 0.61 and XGBoost 0.54, indistinguishable from chance; the best model detects only 51% of uncontrolled cases. The same algorithm is thus the best in one study and the worst in the other. Reported performance is governed by how the task is constructed—label integrity and informativeness of the features—far more than by the algorithm; across clinical prediction studies, machine learning shows no systematic benefit over logistic regression [27]. Accuracy is moreover an improper measure when the positive class is rare: with prevalence π, a model that never alerts reaches $1 - \pi$. Proper scoring rules, which a model cannot improve by reporting other probabilities than its own [28], and calibration [29] are the measures this literature lacks. Table 1.9 sets the studies side by side with the design adopted here.

Table 1.9 How the reviewed machine learning studies were evaluated, against the design of this work.

Study	Data	Predicted	Reported	Limit for this problem
Chatrati et al. [23]	home BP and glucose readings	state of the current reading	rules and a classifier	single readings, not a series
Tai et al. [20]	monitoring architecture	chronic disease status	controlled conditions	no patient validation
Efendi et al. [9]	constructed set, 2 250 against 2 250	“stable”, defined from the input ranges	XGBoost accuracy 0.973	label is a function of the inputs
Mendoza-Cano et al. [10]	1 308 hypertensive adults, national survey	uncontrolled pressure	AUC 0.75 (forest), 0.54 (XGBoost)	cross-sectional; sensitivity 0.51
Liu and Wang [21]	review of IoT studies	various	accuracy 95–100%; AUC 0.70 for readmission	accuracy on unclear tasks
This work	eICU, 174 hospitals; Douala, 32 inpatients	level of the next reading	proper scores against persistence	evaluated on hospital data

Tracking how a patient changes is a related problem. Patient-similarity research uses many distances without a common benchmark [30, 31]; the largest independent comparison of distance measures for series, nine measures on 38 data sets, separates lock-step measures, which compare aligned elements, from elastic ones [32]; and statistical control charts, which compare a patient’s readings with their own baseline, are promising in hypertension and diabetes but little evaluated [33].

1.7 Data Privacy and Ethics

Alhammad et al. [24] reviewed 33 studies of patients’ views on the privacy of mobile health data: concern about data is a consistent barrier to adoption, and encryption, secure authentication and transparency improve trust. Most commercial applications lack a privacy policy or share data with third parties [16], the regulatory dimension is one of the six barriers of the region [11], and the reference implementation of Efendi et al. [9] stores its data in a cloud region outside the country of use. Data protection is therefore a design constraint of this work (Section 3.10).

1.8 Critical Comparison of the Approaches

The studies reviewed follow five approaches (Table 1.10). None combines what the problem requires: clinical validity, professional oversight, longitudinal data, prediction and suitability for low-resource settings.

Table 1.10 Comparison of the approaches found in the literature.

Approach	Studies	Strengths	Weaknesses
Self-management applications	Alessa et al. [16] and Zhou et al. [17]	low cost, behaviour change techniques	little clinical validation; weak effect without a professional
Rule-based telemonitoring with a clinician	Mihevc et al. [8] and Kingue et al. [15]	validated effect on BP and HbA1c; simple, transparent	fixed thresholds, reacts to level not trend, no learning; costly expert time
Tele-expertise between professionals	Bediang et al. [6]	changes clinical decisions; feasible in Cameroon	no home data; specialist delay unchanged
IoT platforms with machine learning	Efendi et al. [9], Tai et al. [20], and Liu and Wang [21]	complete technical chain; offline storage	no BP or glucose sensors; tiny or controlled validation; label leakage
Predictive models on clinical data	Mendoza-Cano et al. [10] and Chatrati et al. [23]	interpretable risk predictors	cross-sectional or single readings; modest sensitivity; no deployment

Three trends emerge from the chronology. Telemonitoring moves from self-management towards systems linked to a clinician, which the most recent evidence identifies as the condition of effectiveness [7, 8, 18]. Prediction moves from rules towards machine learning, with interpretability tools such as SHAP becoming standard [10, 22]. And sensing moves towards cuffless and smartphone-based measurement [19, 22], which the WHO does not accept for clinical decisions [1]. Two points remain controversial: whether mobile health works in Africa, since the effect disappears in developing countries when no professional is involved [7]; and which algorithm is best, since XGBoost is reported as the best [9] and as the worst [10]—a contradiction explained by how the tasks were constructed rather than by the algorithm (Section 2.3).

1.9 Observed Gaps in the Literature

Even with recent progress, five gaps limit the follow-up of patients with hypertension and diabetes in Central Africa:

- **Blind spots.** Central Africa is almost absent from the evidence: one of 51 trials in sub-Saharan Africa [7], comparative reviews limited to South Africa, Kenya and Nigeria [12], and no system combining consultation records, home readings and prediction for both conditions.
- **A rule without memory.** The only validated home protocol for both conditions [8] uses fixed thresholds identical for all patients and reacts to the level of a reading, not to its trend.
- **Fragile predictive evidence.** Machine learning studies of the field suffer from label leakage, improper metrics and validation on the training population [9, 10]; none forecasts a future

state from a patient's own series of blood pressure and glucose.

- **No transfer measured.** No model trained elsewhere has been validated on Cameroonian patients; early-warning scores derived elsewhere lose discrimination in African wards [34, 35].
- **Clinician time.** Systematic expert review does not scale [6, 15], yet no design sizes the alert volume or orders the clinician's work.

Table 1.11 summarises what the literature establishes and the consequence for this work. The present work adopts the three-level protocol of Mihevc et al. [8], the layered architecture of Mucchi et al. [25], offline-first storage as in Efendi et al. [9] and the interpretable modelling practice of Mendoza-Cano et al. [10]; it departs from them by forecasting from each patient's series, by keeping the guideline rule as a floor that the model can raise but never lower, and by validating externally on Cameroonian patients.

Table 1.11 Synthesis of the literature review.

Theme	Established	Not established	Consequence for this work
Burden	cascade fails at follow-up; 72% of diagnosed not controlled; 58% of people with diabetes hypertensive	reliable Cameroonian estimates	longitudinal follow-up; joint blood pressure and glucose
Context	loss to follow-up, inertia, paper records; specialist scarcity	effect of telemedicine on control in Cameroon	offline-first; triage of clinician attention
Clinical evidence	modest effect, conditional on a clinician	effect in sub-Saharan Africa	clinician-in-the-loop decision support
Devices	validated cuff and glucometer required	validated cuffless measurement	manual or Bluetooth entry from validated devices
Machine learning	task construction drives performance	forecasting from individual series	leakage-free task; proper scores; benchmark
Privacy	data concerns hinder adoption	evidence from African patients	anonymisation; declared data residency

Key points of Chapter 1

- Care for both conditions fails at follow-up: in the African Region, 72% of the hypertensive adults already diagnosed are not controlled.
- Telemonitoring lowers systolic pressure by 3 to 7 mmHg when a clinician acts on the data; without one, no effect was shown in developing countries.
- The only validated protocol for both conditions triages single readings with fixed thresholds; it reacts to a level, not to a trend.
- Reported machine learning performance depends on how the task is built: the same algorithm is the best in one study and no better than chance in another.
- Central Africa is almost absent from the evidence, and no learned model has been validated on Cameroonian patients.

CHAPTER 2

Risk-Tiering Framework

This chapter formalises the problem the system solves: turning a patient's irregular series of readings into a level of attention for the clinician. It defines the series and the samples, a label whose every threshold comes from an official source, a forecasting task in which the label cannot leak into the inputs, the decision layer that combines the guideline rule with a model, and a measure of how a patient's state evolves. Chapter 3 instantiates each element on the data.

2.1 Patient Series and Samples

A patient i is observed through four channels $c \in \{\text{SBP}, \text{DBP}, \text{pulse}, \text{glucose}\}$, each sampled on its own schedule: a reading is a pair $(t, x_{i,t}^{(c)})$, and a channel may be missing at any time. The series is not resampled onto a grid: interpolating sparse home readings would invent values and understate variability, one of the quantities of interest.

A sample anchored at time t pairs a window of past readings with a label taken outside it:

$$\mathcal{S}_{i,t} = \Big(\{x_{i,s} : t - L < s \le t\},\ y_{i,t'} \Big), \qquad t' = \min\{s > t + \varepsilon\}, \tag{2.1}$$

where L is the window length and $\varepsilon \ge 0$ an embargo that sets the lead time: the target is the first reading after $t + \varepsilon$ (Figure 2.1). Windows are defined in time rather than in number of readings, because every clinical reference is expressed in time—the seven-day profile, the monthly follow-up—and five readings may span four days for one patient and seven months for another. All the samples of a patient are correlated: the patient, not the sample, is the independent unit.




Figure 2.1 Construction of one sample: the readings of the window, the embargo and the first reading after it.

2.2 A Label from Official Thresholds

The label states how much clinical attention a reading calls for, on four ordered levels: 0 stable, 1 moderate, 2 high, 3 critical. Every threshold comes from an official source, low values count as well as high ones, and a missing channel never counts as normal (Table 2.1). Three choices need a justification. The ESH grades classify a patient's usual pressure, not one reading; they are used here as severity bands of a single reading, the repetition being carried by the window. For pressure and pulse, the National Early Warning Score 2 (NEWS2) is used only through its "red score", the single-parameter value that calls on its own for urgent review [36]: its intermediate bands have meaning only inside the aggregate score. And no accepted cut-off exists for severe hyperglycaemia; the 300 mg/dL of the official hospital-harm measure [37] is a convention, so a sensitivity analysis caps hyperglycaemia at the high level. Boundaries are stored in the unit of their source and converted at comparison, so that no rounding moves a reading across a level; the table is versioned, and every result records the version it was produced with.

Table 2.1 Severity of a single reading, per channel. Pressure in mmHg, pulse in beats per minute, capillary glucose in mmol/L.

Level	Haemodynamic channels	Glucose
3 critical	SBP ≥ 180 or DBP ≥ 110, grade 3 [14]; SBP ≤ 90; pulse ≤ 40 or ≥ 131 [36]	< 3.0, level 2 hypoglycaemia [38]; > 16.7 (300 mg/dL) [37]
2 high	SBP 160–179 or DBP 100–109, grade 2	3.0 to < 3.9, level 1 hypoglycaemia; ≥ 10.0 (180 mg/dL) [39]
1 moderate	SBP 140–159 or DBP 90–99, grade 1	> 7.8 (140 mg/dL) [39]
0 stable	otherwise	otherwise

Within a condition the worst channel governs, and one label is computed per condition:

$$y_t^{\mathrm{H}} = \max\left(s_{\mathrm{SBP},t}, s_{\mathrm{DBP},t}, s_{\mathrm{pulse},t}\right), \qquad y_t^{\mathrm{G}} = s_{\mathrm{glu},t} \text{ if glucose is measured at } t,$$
$$y_t = \max\left(y_t^{\mathrm{H}}, y_t^{\mathrm{G}}\right), \tag{2.2}$$

where s is the level of a channel and the maximum runs over the labels that exist. A single label would read a missing glucose as a normal one; since glucose is ordered for some patients and not for others, the error would be systematic. The *alert* of a condition is a level of high or critical. Figure 2.2 draws the bands of the two channels that carry most of the alerts; both are U-shaped, with a critical level at each end.




Figure 2.2 Level of a single reading for systolic pressure and capillary glucose (Table 2.1). Diastolic pressure and pulse follow the same pattern with their own thresholds.

2.3 The Forecasting Task

If the label is a rule applied to the current reading, $y_t = h(x_t)$, a model that sees x_t only recovers the rule—the circularity of Section 1.6. The legitimate task forecasts a future state from a past window,

$$\hat{y}_{t'} = g\big(\{x_{i,s}\}_{t-L<s\le t}\big), \qquad t' > t + \varepsilon, \tag{2.3}$$

in which the label lies outside the input by construction. Its performance is lower, and that lower figure is the honest one. The comparison that matters is with *persistence*, the guideline rule applied to the last reading of each channel: physiological series are strongly autocorrelated, so carrying the current level forward is a strong baseline, and the contribution of a model is its margin over it.

Two formulations reach the same output, the probability $P(y_{t'} = k)$ of each level k, and neither is assumed better. The **direct ordinal classification** predicts the level of each condition, with a loss that respects the order of the levels. The **value forecast** predicts the value $\hat{x}^{(c)}$ of each channel at the target; with $\hat{\sigma}_c$ the root mean squared error of out-of-fold forecasts, the level probabilities follow by integrating the forecast distribution over the intervals of Table 2.1,

$$P(s_c = k) = \sum_{[a,b)\,:\,\text{level } k} \left[\Phi\left(\tfrac{b-\hat{x}^{(c)}}{\hat{\sigma}_c}\right) - \Phi\left(\tfrac{a-\hat{x}^{(c)}}{\hat{\sigma}_c}\right)\right], \qquad P(y \le k) = \prod_c P(s_c \le k), \tag{2.4}$$

the product assuming that the channel errors are independent given the inputs (Figure 2.3). The value forecast gives a training signal in clinical units on every sample and survives a revision of the guidelines without retraining; the classification needs no assumption on the channels. The alert probability is $p = P(y = 2) + P(y = 3)$.




Figure 2.3 From a value forecast to level probabilities, for one channel. The forecast distribution is cut by the thresholds of Table 2.1; each area is the probability of a level. Within a condition, the channels are then combined by their worst level.

2.4 The Decision Layer

2.4.1 Three tiers and the rule floor

The clinician needs a decision, not a probability: green (record), yellow (more measurements), red (review now). The model's alert probability is mapped to a tier by two cut-points $p_{\text{yellow}} < p_{\text{red}}$. Each cut-point is a decision threshold: alerting at p_t means accepting $(1 - p_t)/p_t$ false alarms per true alert. The yellow tier costs the patient extra readings and the clinician nothing, so its threshold is low; the red tier costs clinician time. Until the partner physician states the costs of errors, $p_{\text{yellow}} = 0.1$ and $p_{\text{red}} = 0.3$ are used, which accept nine and about two false alarms per true alert. The model may raise the tier set by the guideline rule, but never lower it:

$$\text{tier}_{\text{final}} = \max\Big(\text{tier}_{\text{rule}}(x_{\text{last}}),\ \text{tier}_{\text{model}}(p)\Big), \qquad \text{green} \prec \text{yellow} \prec \text{red}. \tag{2.5}$$

A patient at 185/110 mmHg is red whatever the model says, and if the model fails the system falls back to the standard of care rather than to nothing (Figure 2.4).




Figure 2.4 The decision layer. The model can raise the tier given by the guideline rule, never lower it; the cut-points 0.1 and 0.3 are provisional.

2.4.2 The home triage

The label of Table 2.1 uses in-hospital thresholds, those of the data actually collected. A home deployment substitutes the out-of-office thresholds of the only validated protocol for patients with both conditions [8] (Table 2.2). The protocol is *two-stage*: a yellow reading leads to a verification profile—seven days of blood pressure, two readings morning and evening with the second retained, or a one-day profile of six glucose readings—and only the mean of the profile reaches a clinical decision. Noise is thus separated from trend by increasing the number of readings rather than by lowering the threshold. The source leaves five points open, which this work decides: the more severe level applies when systolic and diastolic pressures disagree; a glucose of unknown meal status is held to the after-meal targets; the pulse keeps the NEWS2 red score; a seven-day profile is concluded only with at least six retained readings on three days, the minimum of the ESH practice guidelines [40], and is otherwise reported as incomplete, never as on target; and a glucose profile is done within two days of the trigger. Figure 2.5 shows the resulting path of a reading.




Figure 2.5 Path of a home reading through the two-stage triage. Only the mean of a verification profile, or a red reading, reaches a clinical decision.

Table 2.2 The three-level home triage of Mihevc et al. [8]. Pressure in mmHg, glucose in mmol/L.

Level	Blood pressure	Glucose	Action
Green	110–134 / 60–84	4–7 before a meal; 7–10 after	no change
Yellow	135–174 / 85–104	7.1–10 before; 10.1–14 after	verification profile
Red	≥ 175 / ≥ 105, or < 110 / < 60	< 4; > 10 before; > 14 after	teleconsultation
Profile mean	≥ 135/85 or < 110/60	> 7 before or > 10 after	teleconsultation

2.5 Evolution of the Patient's State

The level answers whether a reading calls for attention; the clinician also needs to know whether the patient has changed. The state of a patient over a window is the vector of the mean of each channel, $\mathbf{s} = (\bar{x}_{\mathrm{SBP}}, \bar{x}_{\mathrm{DBP}}, \bar{x}_{\mathrm{pulse}}, \bar{x}_{\mathrm{glucose}})$. The current state $\mathbf{c}$ covers $(t - 24\,\mathrm{h}, t]$ and the baseline $\mathbf{b}$ the three days before, $(t - 96\,\mathrm{h}, t - 24\,\mathrm{h}]$; evolution is a distance $d(\mathbf{c}, \mathbf{b})$ computed on the channels present in both (Figure 2.6).

Cosine similarity compares directions, not sizes: if every channel rises by half, $\mathbf{c} = 1.5\,\mathbf{b}$ and the cosine distance is zero, so a patient whose pressures and pulse rise together appears unchanged. On raw values it is also dominated by the channel with the largest unit, and its behaviour depends on how the vectors are scaled [41]. No measure is therefore assumed: the candidates (Table 2.3) are the lock-step measures of the benchmark of Wang et al. [32], plus two families for monitoring




Figure 2.6 The two windows compared for one channel. Each window is summarised by its mean; the four channel means form the state vectors **c** and **b**.

an individual: scaling each change by the patient's own variability, the idea of the reference change value of laboratory medicine [42], and the Mahalanobis distance [43], the statistic of multivariate control charts for individual observations [44]. Elastic measures such as dynamic time warping align sequences; a state is a vector of four means, so they do not apply.

Table 2.3 Candidate measures of the change between current state and baseline, from the simplest. σ_j: between-patient SD of channel j; σ_{ij}: SD of patient i, shrunk towards the pooled within-patient SD; Σ: pooled within-patient covariance of changes.

Measure	Definition	Property
Cosine	$1 - \mathbf{c} \cdot \mathbf{b}/(\|\mathbf{c}\|\|\mathbf{b}\|)$	blind to a common change of size
Euclidean, raw	$\|\mathbf{c} - \mathbf{b}\|$	dominated by the largest unit
Euclidean, cohort scale	$\|(\mathbf{c} - \mathbf{b})/\boldsymbol{\sigma}\|$	one scale for all patients
Cosine, centred	cosine of the deviations from the cohort mean	direction of the deviation
Euclidean, patient scale	$\|(\mathbf{c} - \mathbf{b})/\boldsymbol{\sigma}_i\|$	change against own variability
Manhattan, patient scale	$\sum_j \|c_j - b_j\|/\sigma_{ij}$	less weight to one large change
Chebyshev, patient scale	$\max_j \|c_j - b_j\|/\sigma_{ij}$	the largest single change
Mahalanobis	$\sqrt{(\mathbf{c} - \mathbf{b})^\top \Sigma^{-1} (\mathbf{c} - \mathbf{b})}$	correlated channels counted once

Key points of Chapter 2

- One label per condition, on four ordered levels; every threshold comes from an official source, and a missing channel never counts as normal.
- The task forecasts the level of the next reading from a past window, so the label cannot leak into the inputs; persistence is the baseline to beat.
- Ordinal classification and value forecasting give the same output and are compared, not chosen in advance.
- The model may raise the tier of the guideline rule, never lower it; at home, a yellow reading leads to a verification profile.
- Eight measures of state evolution are candidates; the benchmark of Chapter 3 selects one.

CHAPTER 3

Methodology and System Implementation

This chapter instantiates the framework of Chapter 2. It describes the data and how they were collected, their treatment, the features, the candidate models, the experimental protocol and the metrics, then the telemonitoring system built around them and the implementation details that make every result reproducible.

3.1 Type of Research and Study Design

The research is applied and follows a design-science logic: requirements are derived from the literature (Chapter 1), the system is built, and its predictive component is evaluated experimentally on retrospective data under a protocol fixed in advance. The approach is quantitative; the labels, however, are clinical judgements, so the partner physician validates the labelling rule and relabels a sample of readings. The data are secondary: no patient was recruited and no measurement was taken for this work. Figure 3.1 shows how the datasets enter the analysis: models are developed and compared on public multi-centre data only, and the Douala charts are used once, for external validation.




Figure 3.1 Study design. The same treatment, labels and samples apply to both datasets.

Table 3.1 The three data sources and their role.

Source	Content	Collection	Role
eICU Collaborative Research Database [45, 46]	intensive-care stays in United States hospitals, 2014–2015; open demonstration subset of 2 520 stays of 1 841 patients in 186 hospitals	electronic records of routine care, de-identified	development and comparison of the models
Douala inpatient charts	32 patients, 366 readings, 8 June to 30 August 2026; pressure, pulse and capillary glucose, morning and evening	paper observation charts, transcribed by hand without names	external validation only
Douala consultation records	58 outpatients, 81 readings, 49 with a single reading	paper consultation records, transcribed without names	description of the fragmentation of follow-up

3.2 Data Sources and Collection

3.2.1 The three sources

Three sources are used, each for a single purpose (Table 3.1). The research was carried out at the Laboratory of Computer Engineering, Data Science and Artificial Intelligence of the NHPSD and, for the Douala records, at the Laquintinie Hospital of Douala, a public hospital, under the supervision of an endocrinologist of the hospital (Section 3.10).

The Douala charts were transcribed one row per reading. The transcription records whether a reading was taken in the morning or the evening, but not its hour nor the admission diagnosis. In the wards, pressure and pulse follow the nursing schedule, whereas glucose is ordered according to the clinical picture. The consultation records, with a single reading for most patients, document the fragmentation of follow-up but cannot support longitudinal modelling; no home measurement could be collected, so the method is written for irregular home series and applied to the twice-daily charts. Figure 3.2 traces the path from the paper record to the analysis data, and Figure 3.3 shows what the charts contain, patient by patient.




Figure 3.2 From the paper records of the Douala hospital to the analysis data. No name leaves the hospital; the slot is morning or evening, the hour is not recorded.

Values are stored in one canonical unit per channel, with plausible ranges of 60–260 mmHg for SBP, 30–160 for DBP, 30–220 beats per minute for pulse and 1.0–35.0 mmol/L for glucose. Glucose is the riskiest field, because three conventions coexist and Cameroonian practice uses g/L: 1 mmol/L = 18.0156 mg/dL from the molar mass of glucose. The exact factor matters: with the rounded value of 18.0, a reading of 1.80 g/L falls below the 180 mg/dL threshold of Table 2.1.

3.2.2 Sample size and the role of each dataset

Whether the Douala charts suffice to develop a model is answered with the criteria of Riley et al. [47] and Riley et al. [48]: the sample must keep the optimism of the model small (shrinkage $S \geq 0.9$), keep the apparent and adjusted R^2 within 0.05, and estimate the overall risk within ± 0.05. The first criterion gives

$$n \geq \frac{p}{(S-1)\,\ln\!\left(1 - R^2_{\mathrm{CS}}/S\right)} \tag{3.1}$$

for p candidate parameters and an anticipated Cox–Snell R^2_{CS} [49]. The criteria are computed with a port of the reference implementation [50], checked against its published example. The samples of one patient are correlated, so the effective sample size divides their number by the design effect $1 + (\bar{m} - 1)\rho$ [51], with $\bar{m}$ the mean number of samples per patient and ρ the intraclass correlation of the alert.




Figure 3.3 The 366 readings of the Douala charts. One row per patient, one mark per reading, coloured by level (grey stable, blue moderate, light red high, red critical); open marks are readings without glucose. Patients are numbered by date of first reading.

The charts fall far below these criteria (Section 4.1). The candidate models are therefore developed and compared on eICU, which covers many hospitals: the outer folds can be grouped by hospital, which estimates how a model transfers between sites before it meets the Douala patients [52]. The Douala charts serve for external validation only, a transfer that no study identified in the region has measured for a learned model; early-warning scores derived elsewhere lose discrimination in African wards and require local validation and recalibration [34, 35]. With about 64 haemodynamic and 112 glycaemic alert targets, below the 100 events and 100 non-events recommended for external validation [53, 54], its estimates are reported with patient-level intervals and read as exploratory.

3.2.3 Harmonisation of the intensive-care records

In eICU, every adult stay with hypertension or diabetes recorded in the diagnoses or the past history is eligible. Intensive-care monitoring is far denser than a ward chart, so each stay is resampled to the rhythm of the charts: two slots a day, at 08:00 and 20:00, each taking the non-invasive cuff pressure, the pulse and the bedside glucose nearest to the slot within two hours (Figure 3.4). The slot hours are an assumption, since the charts record only morning and evening, and the tolerance is varied in a sensitivity analysis. Both datasets then share one sample definition, and the remaining differences—intensive care against the ward, the United States against Cameroon—are the shift that external validation measures.




Figure 3.4 Harmonisation of an intensive-care stay to the rhythm of the ward charts. At each slot, the reading nearest in time within two hours is kept for each channel (red); a slot with none is missing, as an unrecorded value on a chart would be.

3.3 Data Curation and Preprocessing

3.3.1 A two-phase pipeline

This is where label leakage and biased imputation are either prevented or built in. Every step that learns a parameter from the data must be fitted on the training fold only, so the treatment is split into two phases separated by the train–test split (Figure 3.5) and implemented as a single pipeline object. A step placed on the wrong side of the split causes no error: it silently produces an optimistic estimate that fails in deployment.




Figure 3.5 The two-phase preprocessing pipeline.

3.3.2 What the readings say about the instruments

The charts do not name the devices, so their validation cannot be checked; two analyses of the readings bound what this uncertainty implies. **Terminal digits**: a manual aneroid or mercury scale is graduated in steps of 2 mmHg, so manual readings end in even digits with an excess of zeros, whereas an electronic display shows any integer [55]; a χ^2 test of uniformity of the last digit tells the two apart. **Label uncertainty**: a validated cuff may err by up to 5 mmHg on average [56], and a glucose meter by 15 mg/dL below 100 mg/dL and 15% above [57]; shifting every reading by these tolerances and counting the readings whose level changes measures how much of the disagreement between a model and the label is noise that no model can remove.

3.3.3 Quality controls at entry

The controls of Table 3.2 learn nothing from the data, so they are applied to all records before the train–test split. Flagged records are kept: a run of identical values is information about how the data were produced. Impossible values are the exception, and are removed from features and labels.

Table 3.2 Quality controls applied at entry.

Control	Rule	On violation
Plausibility	value within the plausible range of its channel	value removed
Coherence	SBP > DBP and SBP − DBP ≥ 10 mmHg	pair removed: one value is wrong, which cannot be known
Duplicate	same patient, time and values	one kept, the rest flagged
Carried forward	three or more identical consecutive values	flagged
Unit	magnitude consistent with another glucose convention	flagged
Extreme	robust $\lvert z\rvert > 3.5$ (below)	flagged; “contradicted” if the next reading is not extreme

3.3.4 Outliers: artefact or event?

In a risk-detection system the outlier is often the event of interest: a systolic reading of 185 mmHg is a statistical outlier for a patient whose median is 138, and also a red alert. An automatic cleaning step would delete the positive class. The dividing line is therefore physiological plausibility, not statistical rarity. A statistical threshold is used for flagging only, with the median absolute deviation (MAD), which is not inflated by the outliers it detects, shrunk towards the cohort value because a personal MAD from ten readings is unstable:

$$z_{i,t} = \frac{x_{i,t} - \operatorname{median}(x_i)}{1.4826\,\widehat{\mathrm{MAD}}_i}, \qquad \widehat{\mathrm{MAD}}_i = w_i\,\mathrm{MAD}_i + (1 - w_i)\,\mathrm{MAD}_{\mathrm{pooled}}, \quad w_i = \frac{n_i}{n_i + n_0}. \tag{3.2}$$

With $n_0 = 20$, a patient with ten readings takes one third of the estimate from their own data; as the series grows, the estimate becomes personal. A reading is flagged when $\lvert z_{i,t}\rvert > 3.5$ [58]; flags serve clinician review and never enter the features.

3.3.5 Missing data

Imputation is valid when data are missing completely at random or at random given observed values, and invalid when missingness depends on the unobserved value. Glucose in the wards is ordered according to the clinical picture, and at home discouraging readings may be omitted: in both cases absence is informative. Imputing a normal value would teach the model that absence predicts stability, whereas it predicts deterioration—a sign error that more data cannot correct. Absence is therefore represented as information: the number of glucose readings in the window and the time since the last one are features. Tree ensembles route missing values at each split. The recurrent model GRU-D [59] receives the mask of what was measured and decays the last value towards the training mean as the gap grows,

$$\tilde{x}_t^{(c)} = m_t^{(c)} x_t^{(c)} + \left(1 - m_t^{(c)}\right)\left[\gamma_t^{(c)}\, x_{\mathrm{last}}^{(c)} + \left(1 - \gamma_t^{(c)}\right)\bar{x}^{(c)}\right], \qquad \gamma_t^{(c)} = \exp\!\left(-\kappa_c\, \Delta t_{\mathrm{last}}^{(c)}\right), \tag{3.3}$$

with a decay rate per channel, since a three-day-old pressure says more about today than a three-day-old glucose. The proportional-odds model, which cannot accept missing inputs, receives the training median with a missingness indicator. The features keep their clinical units, since the label is defined on absolute thresholds; inputs are standardised with training statistics for the linear and neural models (Table 3.3).

Table 3.3 Preparation required by model family, fitted on the training fold.

Model	Scaling	Missing values
Proportional-odds logistic regression	yes	training median with missingness indicators
Random forest, CatBoost	no	handled natively by the splits
GRU-D, masked LSTM	yes	mask and decay towards the training mean

3.4 Features

The tabular models receive descriptors of the window, computed per channel on its readings only: last value, mean, standard deviation, slope, number of readings and time since the last one, plus pulse pressure and the age and sex of the patient, the only covariates recorded in both datasets. The slope is the least-squares coefficient on the real reading times, which handles gaps without resampling, and is computed only from at least three readings spanning half the window; a feature the window cannot support is missing, not zero. A window of n readings cannot carry more than n independent quantities—with two readings, the standard deviation, the slope and the successive difference are the same number—so the feature sets are kept small and fixed before any result (Table 3.5). No signed margin to a threshold is computed: for a fixed threshold it is an affine function of the last value and adds no information. The sequence models receive the readings of the window with their mask.

The window length is a hyperparameter, searched over 24, 48, 72 and 96 hours, and what each length can carry follows from the number of readings it holds (Table 3.4). Because the length is searched, the set of forecast targets must not depend on it, or each length would be evaluated on different targets: a reading is an anchor when the shortest window, 24 hours, holds at least two pressure readings, and longer windows add history to the same anchors. Overlapping windows multiply the rows but not the information, so the patient is the independent unit: all the windows of a patient fall on the same side of every split, and intervals are computed at the patient level.

Table 3.4 Window length and the features it can support, at two readings a day.

L (h)	Readings	Features supported
24	2	level only
48	4	level and slope; variability marginal
72	6	level, slope and variability
96	8	level, slope and variability

Table 3.5 Feature sets, fixed before any result.

Set	p	Features
Primary	8	last SBP, slope of SBP, last DBP, last pulse, last glucose, slope of glucose, number of glucose readings, age
Extended	16	primary, plus SD of SBP and glucose, slopes of DBP and pulse, pulse pressure, time since the last glucose, mean SBP, sex

3.5 Candidate Models and Baselines

No model is chosen in advance: the candidates are the families that published benchmarks of comparable problems put forward, and the benchmark of this work selects among them (Table 3.6). On clinical prediction tasks, machine learning has shown no systematic benefit over logistic regression [27]; on tabular data, tree ensembles generally match or beat deep networks [60], and in the most recent living benchmark CatBoost is the best gradient-boosting model when tuned [61]. Only deep temporal models read the data as a sequence, which hypotheses H1 and H3 compare. Four families are left out for stated reasons: a single tree is dominated by its ensembles; support vector machines give poorly calibrated probabilities; XGBoost ranks below CatBoost in the same benchmark; and the foundation model TabPFN is limited to 10 000 training samples [62], which the development data exceed.

Table 3.6 Candidate families and baselines.

Family	Candidate	Why it is in the pool
Baselines	prior (level frequencies); persistence (rule on the last reading, with training transition probabilities)	the margin over persistence is the contribution of a model
Linear	proportional-odds logistic regression [63], L2 penalty; the same on principal components	no systematic loss against machine learning [27]; order of the levels in the model
Bagging	random forest [64]	reference of the reviewed literature [10]
Boosting	CatBoost [65]	best tuned gradient boosting [61]
Value forecast	ridge or CatBoost regression per channel, levels by integration (Section 2.3)	the second formulation
Deep temporal	GRU-D [59]; masked LSTM [66]	hypothesis H3

3.6 Experimental Protocol

All constraints combine into a nested cross-validation on the development data (Figure 3.7). Five outer folds estimate performance and are grouped by hospital, so that every test fold contains hospitals the model has never seen; four inner folds, grouped by patient, select the window and the hyperparameters. The search uses the tree-structured Parzen estimator of Optuna [67, 68] over a space declared before any run, split into structural hyperparameters (window, depth, units, components) and training hyperparameters (learning rate, penalties, dropout, class weights); each configuration is scored by its ranked probability score on pooled inner-validation predictions, and the configuration kept is the one with the lowest selection cost (Section 3.7). The number of configurations per candidate is fixed in advance (Appendix B), since the optimism

of a selection grows with it [69]. Across families, the lowest inner selection cost gives the *selected* model of the fold, and its outer result estimates the whole procedure. The procedure is repeated three times with different fold assignments, and two models are compared with the corrected resampled t test, which accounts for the overlap between training sets [70]. Figure 3.6 details what happens inside one outer fold.




Figure 3.6 Search and selection in one outer fold. The outer test hospitals are used once, after every choice has been made on the inner folds.




Figure 3.7 Nested cross-validation on the development data, repeated three times.

Each selected configuration is then refitted on all the development data and applied once to the Douala charts; nothing is tuned on them, and their intervals come from a bootstrap that resamples patients [53]. The prevalence of the alert differs between intensive care and Douala wards, which shifts calibration even for a model that transfers well, so performance is reported as is, and again after updating the intercept alone on the Douala data [71]: the first is the test of transportability (H4), the second shows what a local recalibration recovers. A classifier trained with class weights w_k overstates each level in proportion to its weight; its probabilities are corrected exactly before use, $p_k = (p'_k/w_k)/\sum_j p'_j/w_j$. H2 is tested by repeating the benchmark with embargoes of 12 and 36 hours, for targets 24 and 48 hours ahead.

3.7 Evaluation Metrics

The measures are chosen by explicit criteria [72]: a clear focus, and, for overall accuracy, a proper scoring rule [28]. Accuracy, the F1 score and recall at a fixed threshold fail these criteria. Table 3.7 lists the admissible measures; the positive class is the alert, and every measure carries a patient-level confidence interval. Performance is also reported by sex and by age band (below and above 65 years), and across the hospitals of the outer folds.

Table 3.7 Admissible performance measures.

Focus	Measure	Why
Overall, four levels	ranked probability score (RPS) [73]	proper for ordered levels: an error across several levels costs more than an adjacent one
Overall, alert	Brier score, log loss	proper
Discrimination	area under the ROC curve (AUROC) of the alert	independent of any threshold
Calibration	intercept and slope [29]	a probability shown to a clinician must mean what it says
Clinical utility	net benefit, decision curve [74]	true alerts weighed against false alarms at the clinical threshold

In the inner folds, configurations are compared on one number, the **selection cost**. Four terms—the RPS, the AUROC, the calibration error $|a|+|\ln b|$ for intercept a and slope b, and the net benefit averaged over thresholds of 0.1 to 0.3—are standardised across the configurations compared (minus the median, divided by the scaled MAD, oriented so that lower is better) and combined:

$$C = \sum_m w_m \, z_m, \qquad \sum_m w_m = 1. \tag{3.4}$$

The weights express how much each focus matters clinically, so they are a value judgement, not a parameter: tuning them on the data that judge the models would let any weighting favour a chosen model. They are equal in the provisional analysis, to be replaced by weights derived from the costs of errors stated by the partner physician, and a sensitivity analysis draws them at random to report how often the selected configuration remains first. The outer folds report every measure separately, never the composite alone. Interpretability is reported only for a model shown to beat persistence: grouped permutation importance, which permutes the features of one channel together because systolic and diastolic pressures are correlated [75], and exact tree SHAP values for CatBoost [76].

The measures of state evolution (Table 2.3) are compared on the same samples: a measure is useful if its distance ranks higher the samples in which the level of the next reading differs from that of the baseline state (primary outcome), or becomes an alert (secondary outcome). Each is evaluated by its AUROC with a patient-level interval, its scales are estimated on eICU only, and the measure retained is the best on the primary outcome, or the simplest candidate whose interval contains the best value.

3.8 The Telemonitoring System

3.8.1 Architecture

The system has four parts (Figure 3.8). A cross-platform **application** (Flutter) serves the patient and the clinician. A managed document database (**Cloud Firestore**) holds the record and synchronises the applications, with a local cache that turns a reading entered offline into a delayed one. A local **risk service** (FastAPI) scores a patient's recent readings with the model selected by the benchmark, using the same code as the research pipeline, so that the features seen in deployment cannot drift from those seen in training. And a local **worker** runs the server functions: measurement check, escalation, notification, the non-transmission sweep and the escalation of events left unreviewed. Hosting the service and the worker online is outside the scope of this work; without them, the application still applies the home triage on the phone.




Figure 3.8 Architecture of the built system. Dashed: push notifications sent by the worker to the two applications.

3.8.2 Data model and rules

Each patient's readings, events and messages are subcollections of their profile, so that one security rule governs them and the readings are read in time order with one query. A reading holds one measurement session: the measurement time stated by the patient and the arrival time set by the server, the retained values, the first reading of a protocol pair kept for audit, the context of a glucose reading, and the quality flag set by the worker. The record is append-only: the rules refuse any change or deletion of a reading, and a correction is a new reading that names the one it supersedes. Each write is validated on the server (known fields, plausible ranges, coherent pressure pair, measurement time at most thirty days before arrival). An event stores what raised it (rule, model, verification profile, or a silence longer than the patient's checkup period), the triggering values, the threshold and model versions, and the clinician's verdict and action; it is closed once, and its identifier is derived from what raised it, so the same alert raised by the phone and by the worker is one document (Figure 3.9).




Figure 3.9 Data model of the record in Cloud Firestore. Readings, events and messages are subcollections of the patient's document, so one security rule governs each of them.

3.8.3 Server functions

Table 3.8 lists the functions of the worker. It reads and writes the database through the Firebase Admin SDK at a fixed interval; readings and events older than a day found at its first run are marked without being notified, so that a backlog does not flood the queue. The delays are configuration values, provisional like the tier cut-points.

Table 3.8 Server functions run by the worker.

Function	Fires on	Action
Measurement check	new reading	quality controls of Section 3.3.3; quality flag
Escalation	yellow or red home level	event added to the clinician's queue
Scoring	latest reading	rule and model, same code as the risk service; event if the model raises the tier
Notification	new event	red: patient and clinician; yellow: patient, prompt for the verification profile
Non-transmission	no reading for longer than the checkup period	event; daily reminder to the patient
Queue age	event open 4 h (red) or 3 days (yellow)	escalated to the clinician's named alternate
Verdict	event closed	the patient is told it was reviewed

Figure 3.10 follows one alert through these functions, from the patient's reading to the clinician's verdict.

3.8.4 Interfaces

The **patient** enters the two pressure readings of the protocol, of which the second is retained, and may date a reading taken earlier; the glucose unit is fixed to mmol/L, a value outside its plausible range is refused with a message naming the unit, and the measurement context is required for glucose because the target depends on it. A reading at the red level brings up a local warning at once, whatever the connectivity. The home screen shows the level of the latest readings with the one instruction the patient may receive—contact the health centre, or seek care today—never a dose and never a probability; the verification profile to carry out and its progress; the fourteen-day trend with the target band; the evolution of each channel; and the




Figure 3.10 Life of an alert. The phone warns the patient without a network; the record, the checks, the notifications and the escalation do not depend on the phone staying online.

outcome of the clinician's reviews. The **clinician** works from a queue of unreviewed events, red before yellow and then oldest first, each row showing the source of the alert, the triggering value, the fourteen-day trend and the versions, and closes an event with a verdict (true alert, false alert, uncertain) and the action taken in two or three taps; the verdicts are the clinically adjudicated labels that escape the circularity of rule-derived labels. A health assistant answers general questions in French or English under four guardrails (no medication advice, no diagnosis, no false reassurance, emergency symptoms sent to hospital); it receives no identifier, and its guardrails are tested on a fixed set of 22 adversarial questions. Appendix D shows the screens.

3.9 Implementation Details and Reproducibility

Environment. The analysis runs on a laptop (AMD Ryzen 5 PRO 3500U, four cores, no GPU) with the software of Table 3.9; the exact versions of every package are fixed in a lock file. The tree SHAP values of CatBoost are computed by CatBoost itself.

Reproducibility. The labelling rule, the home triage, the quality controls, the sample-size criteria and the measures are implemented once and checked by automated tests (171 in the research package, 48 in the application), including the parity of the levels computed on the phone with the Python reference on every threshold boundary. Thresholds, the selection cost, search spaces and the delays of the worker are kept in versioned configuration files, and exact package versions in a lock file. Every benchmark run is saved in a folder named after a hash of everything that determines its results, with its configuration, a copy of the code, every trial, every prediction and the fitted models, so that each figure is redrawn from saved outputs and an interrupted run resumes where it stopped. The seed is fixed. Code and configurations are versioned; the Douala data are not published, and the eICU data are available from PhysioNet under its licence.

Table 3.9 Software and versions used for the analysis and the system.

Role	Software	Version
Language of the analysis	Python	3.10.11
Data handling	pandas; NumPy	2.3.3; 2.2.6
Proportional-odds model, statistical tests	SciPy	1.15.3
Random forest, principal components, folds	scikit-learn	1.7.2
Gradient boosting, tree SHAP values	CatBoost	1.2.10
GRU-D and masked LSTM	PyTorch (CPU)	2.12.0
Hyperparameter search	Optuna	5.0.0
Tests of the research package	pytest	9.1.1
Risk service	FastAPI; Uvicorn	0.149.0; 0.54.0
Worker (server functions)	Firebase Admin SDK for Python	7.7.0
Application	Flutter; Dart	3.44.2; 3.12.2
Record and notifications	Cloud Firestore; Firebase Cloud Messaging plugins	6.7.1; 16.7.0

3.10 Ethical Considerations

The eICU data are de-identified and distributed by PhysioNet under its terms [45, 77]. The Douala charts are a secondary use of routine records, transcribed without names under the research authorisation of the Laquintinie Hospital of Douala (No. 185/AR/MINSANTE/DHL/CM of 25 September 2026, for the period from 28 September to 4 November 2026). The authorisation places the work under the supervision of an endocrinologist of the hospital and under the hospital's code of ethics and professional conduct, and requires that any publication preserve the interests of the hospital and of its staff and that a copy be deposited at its documentation centre. A prospective study with home measurements, which involves patients directly, would in addition require the opinion of a research ethics committee. Removing names is not enough to anonymise a small cohort: before any analysis file is shared, ages are grouped in five-year bands and calendar dates replaced by the time elapsed since each patient's first reading, which is all the analysis uses. Personal data in Cameroon are governed by Law No. 2024/017 on the protection of personal data [78], which makes any transfer abroad conditional on the authorisation of a supervisory authority, and the security of electronic communications by Law No. 2010/012 [79]; the region of the database is therefore chosen before the first write. The health assistant sends conditions, medications and readings to a provider outside Cameroon, which requires the patient's informed consent.

Key points of Chapter 3

- Models are developed and compared on eICU data from 174 hospitals, and applied once to the 32 Douala inpatients; nothing is tuned on the Douala data.
- Quality controls flag rather than delete: only impossible values are removed, because an outlier may be the event the system must detect.
- Absence of a measurement is information; every step that learns from the data is fitted inside the training fold.
- Nested cross-validation grouped by hospital and patient, a declared search budget and a declared selection cost replace the choice of a model in advance.
- The system—application, append-only record, risk service and worker—applies the home triage on the phone, without a network.

CHAPTER 4

Results and Discussion

This chapter reports what the data, the benchmark and the system produced, in the order of Chapter 3: the collected data and their treatment, the benchmark of the risk models and their external validation, the comparison of the measures of state evolution, and the verification of the system. It then answers the research questions, compares the results with the literature and states the limitations.

4.1 Data Collection and Treatment

Figure 4.1 traces both datasets from their source to the forecast samples on which the models are developed and validated; the subsections below describe each step.




Figure 4.1 Flow of the data from the two sources to the forecast samples.

4.1.1 The Douala charts

The charts give 366 readings of 32 inpatients between 8 June and 30 August 2026, a median of 11.5 readings per patient (range 5–18). The rhythm is that of the nursing round: two readings on 177 of the 189 patient-days, one on the other 12, and no gap longer than one day. Pressure and pulse are present at almost every reading, glucose at two thirds of them, and 8 patients have no glucose at all (Table 4.1).

Table 4.1 The Douala charts after the quality controls of Table 3.2.

	SBP	DBP	Pulse	Glucose
Readings present (of 366)	97.3%	97.0%	96.7%	68.6%
Out of plausible range	0	1	0	0
Incoherent pressure pair	1		–	–
Carried forward (≥ 3 identical)	0	6	9	0
Extreme, robust $\lvert z\rvert > 3.5$	4	3	0	0
of which contradicted by the next reading	4	3	–	–
Unit suspect, duplicate	0	0	0	0

Three values were removed: a diastolic pressure outside its range, and both values of an incoherent pressure pair on another reading; everything else is kept and flagged. The seven extreme pressures are all isolated, each followed by a reading that is not extreme. They are therefore either transcription errors or transient events, which the readings alone cannot distinguish; they stay in the data, which is the reason for keeping outliers stated in Section 3.3.4. The flagging depends little on the shrinkage constant: with $n_0 = 10$, 20 and 40, the number of extreme readings is 5, 7 and 9. The 15 carried-forward values concern the diastolic pressure and the pulse, never the systolic pressure or glucose.

4.1.2 What the readings say about the instruments

The terminal digits separate the channels clearly (Figure 4.2). Systolic and diastolic values do not depart from a uniform last digit (χ^2 test, $p = 0.74$ and 0.26): 50% and 46% end in an odd digit, and zeros account for 10% and 13%, where a manual scale graduated in steps of 2 mmHg would give no odd digit and an excess of zeros. The pressures were therefore read from an electronic display. The pulse behaves differently: 84% of the values are even and 69% end in 0, 4 or 8 ($p < 10^{-50}$), consistent with values read off the graphical pulse curve of the chart, whose grid sets the resolution. This resolution does not affect the label, whose pulse thresholds lie far from it: the four readings at or above 131 beats per minute range from 135 to 144.

The device tolerance has a larger effect. Shifting every reading by the tolerance of a validated device can change the level of 27% of the systolic readings, 39% of the diastolic readings and 56% of the glucose readings: these readings lie close enough to a threshold for the measurement error alone to move them across it. On them, a disagreement between a model and the label is not necessarily a model error, and no model can be expected to remove it.




Figure 4.2 Terminal digits of the Douala readings. The dashed line marks the uniform share of 0.1.

4.1.3 Labels

Figure 4.3 gives the levels at the forecast targets, the unit at which the models are evaluated. In Douala the alert concerns 22.7% of the 282 haemodynamic targets (64 alerts) and 58.9% of the 190 glycaemic targets (112 alerts); critical hyperglycaemia alone accounts for a quarter of the glycaemic targets.




Figure 4.3 Levels at the forecast targets. Douala: 282 haemodynamic and 190 glycaemic targets; eICU demonstration subset: 4 182 and 1 356.

The sourced rule against the earlier rule. The first version of this work used an unsourced rule (ESH grades, glucose cut-points of 8, 11 and 14 mmol/L, no low thresholds). On the 366 readings as transcribed, the two rules agree on 84% of the readings (quadratically weighted $\kappa = 0.81$), and 58 readings change level (Table 4.2). The changes have two sources. Low values now count: 18 readings rated stable by the earlier rule are critical, 15 of them with a systolic pressure of 90 mmHg or less and 4 with a pulse of 131 or more. And the glucose bands moved to the official thresholds: the 22 readings between 14.0 and 16.7 mmol/L go from critical to high, and 11 readings of 10 mmol/L or more go from moderate to high. Capping hyperglycaemia at the high level (Section 2.2) has a large effect: the critical glycaemic readings fall from 65 to 2.

The relabelling by the partner physician, the secondary endpoint, is sized by simulation on the observed levels: 100 readings give a 95% interval of half-width 0.10 for a weighted kappa near 0.74, and 150 readings a half-width of 0.09 for a kappa near 0.63.

Table 4.2 Levels of the 366 Douala readings under the earlier rule (rows) and the sourced rule (columns).

	stable	moderate	high	critical	total
stable	106	2	0	18	126
moderate	0	57	15	1	73
high	0	0	63	0	63
critical	0	0	22	82	104
total	106	59	100	101	366

4.1.4 The development data

The eICU demonstration subset holds 2 520 stays of 1 841 patients in 186 hospitals. Of these, 1 437 stays of 1 141 patients in 184 hospitals are eligible (hypertension 1 200, diabetes 684, both 447). Harmonisation to the two daily slots gives 7 269 readings in 1 348 stays, then 4 189 forecast samples of 817 stays, 700 patients and 174 hospitals. The rhythm is now that of the charts, but the content is not: glucose is present in 34% of the harmonised readings against 69% in Douala, and critical hyperglycaemia concerns 2.8% of the glycaemic targets against 26.3% (Figure 4.3). The slot tolerance matters for glucose and little for the pressures (Table 4.3).

The values differ as well (Figure 4.4). Systolic pressure has almost the same distribution in both datasets: median 124 mmHg (interquartile range 108–145) in Douala and 122 (107–141) in intensive care. Diastolic pressure does not: median 82 mmHg (70–94) against 64 (55–74), a shift that the haemodynamic label inherits through its diastolic thresholds. Glucose is higher in the wards, median 11.1 mmol/L (7.2–16.6) against 7.9 (6.3–10.2), with a long tail above the critical threshold that the intensive-care readings rarely reach.




Figure 4.4 Distribution of the readings in the two datasets, with the thresholds of Table 2.1 on the axes. Douala: 356 systolic, 355 diastolic and 251 glucose readings; eICU: 6 577, 6 563 and 2 444.

4.1.5 How often the level changes

Persistence, the reference every model must beat, predicts that the next reading has the level of the current one. Figure 4.5 shows how often this holds. For the haemodynamic label the level is unchanged in 56% of the Douala pairs and 59% of the eICU pairs, and a reading below the alert becomes an alert in 13% of the pairs in both datasets. An alert, however, persists more often in Douala (54%) than in intensive care (34%). The glycaemic label behaves differently in the wards: the level is unchanged in only 30% of the Douala pairs, against 55% in eICU, and a reading below the alert is followed by an alert in 43% of the cases. Glucose in the Douala wards

moves between levels from one reading to the next, which makes the glycaemic forecast harder there than in the development data, whatever the model.




Figure 4.5 Level of the target (columns) given the level of the anchor reading (rows), in percent of each row: S stable, M moderate, H high, C critical. The diagonal is what persistence predicts. Pairs where both readings carry the label: 282 and 4 182 haemodynamic, 182 and 894 glycaemic.

Table 4.3 Effect of the slot tolerance on the eICU development samples.

Tolerance around 08:00 and 20:00	± 1 h	± 2 h (main)	± 3 h
Glucose readings kept	1 350	2 444	3 331
Haemodynamic samples	4 018	4 182	4 249
Glycaemic samples	645	1 356	1 983

4.1.6 Sample size

Table 4.4 applies the criteria of Section 3.2.2 to both datasets. The samples of a patient are strongly correlated (intraclass correlation 0.13 to 0.32), so the effective sample sizes are three to five times smaller than the number of samples. The Douala charts cannot support a model: even the overall risk cannot be estimated to ±0.05 (270 and 372 samples needed, against 72 and 53 effective), and the budget is at most one parameter. In the development data, the haemodynamic task is well within the criteria. The glycaemic task is not: at a C statistic of 0.75, it supports about four parameters, and eight would need about 520 patients where the demonstration subset has 312. The full eICU database, about eighty times larger, removes this constraint. Figure 4.6 shows how the budget depends on the discrimination anticipated: at no plausible value do the Douala charts support the eight features of the primary set.




Figure 4.6 Parameters allowed by the criteria of Riley et al. with the effective sample size, against the anticipated C statistic.

Table 4.4 Sample size against the criteria of Riley et al. Parameters for an anticipated C statistic of 0.70 to 0.80.

	Douala		eICU demonstration	
	haem.	**glyc.**	**haem.**	**glyc.**
Samples (patients)	282 (32)	190 (24)	4 182 (699)	1 356 (312)
Prevalence of the alert	0.23	0.59	0.17	0.29
Intraclass correlation; design effect	0.32; 3.9	0.31; 3.6	0.13; 3.2	0.30; 5.4
Effective sample size	72	53	1 292	251
Samples needed for the overall risk	270	372	215	318
Parameters allowed, effective sample	0–1	0–1	10–25	2–7
Patients needed for 8 parameters ($C = 0.75$)	≈ 216	≈ 168	≈ 325	≈ 518

4.2 Benchmark of the Risk Models

Both runs are complete: five outer folds grouped by hospital, repeated three times, so 15 outer estimates per candidate, with the search budgets of Appendix B. Each repeat predicts 4 182 haemodynamic and 1 356 glycaemic targets of the development data once. Tables 4.5 and 4.6 give every measure for every candidate, the baselines first; the row *Selected (procedure)* is the outer result of the model chosen in each fold by the inner selection cost, the estimate of the whole procedure. Figure 4.7 shows the quantity on which H1 rests, the difference in RPS with persistence in each outer fold.

Table 4.5 Outer-fold results, haemodynamic task: mean (SD) over 15 outer folds. ΔRPS: difference with persistence (negative is better); p: corrected resampled t test. Net benefit averaged over the thresholds 0.1 to 0.3.

Model	**RPS**	**ΔRPS**	p	**Brier**	**AUROC**	**Int.**	**Slope**	**NB**
Prior	0.1504 (0.0090)	+0.0119	< 0.001	0.140	0.500 (0.00)	−0.01	0.72	0.025
Persistence	0.1384 (0.0082)	–	–	0.133	0.648 (0.03)	−0.01	0.99	0.038
Ordinal logistic	0.1391 (0.0087)	+0.0007	0.727	0.134	0.616 (0.04)	0.07	0.67	0.036
Logistic on PCA	0.1437 (0.0103)	+0.0052	0.034	0.137	0.595 (0.03)	0.05	0.84	0.030
Random forest	0.1339 (0.0074)	−0.0045	0.004	0.130	0.678 (0.03)	0.02	1.02	0.043
CatBoost	0.1341 (0.0077)	−0.0043	0.002	0.131	0.668 (0.03)	0.02	1.02	0.042
Value forecast (ridge)	0.1342 (0.0081)	−0.0042	0.004	0.131	0.677 (0.04)	−0.02	0.92	0.045
Value forecast (CatBoost)	0.1349 (0.0074)	−0.0036	0.003	0.131	0.667 (0.04)	−0.03	1.03	0.042
GRU-D	0.1348 (0.0089)	−0.0036	0.121	0.131	0.658 (0.03)	0.07	0.77	0.043
Masked LSTM	0.1339 (0.0083)	−0.0046	0.049	0.129	0.668 (0.03)	0.08	0.81	0.044
Selected (procedure)	0.1345 (0.0079)	−0.0039	0.035	0.131	0.665 (0.03)	0.01	0.88	0.042

4.2.1 Learned models against the guideline rule

Haemodynamic task. Persistence is a strong baseline: the guideline rule on the last reading, with the transition probabilities of the training data, has an RPS of 0.1384, against 0.1504 for the prior. Five families do better, with a difference that the corrected resampled t test finds significant: random forest (−0.0045, $p = 0.004$), CatBoost (−0.0043, $p = 0.002$), the ridge and CatBoost value forecasts (−0.0042 and −0.0036) and the masked LSTM (−0.0046, $p = 0.049$). The procedure as a whole improves on persistence by 0.0039 ($p = 0.035$). The gain is small, about 3% of the RPS of persistence, but consistent: CatBoost has a lower RPS than persistence

Table 4.6 Outer-fold results, glycaemic task: mean (SD) over 15 outer folds. ΔRPS: difference with persistence (negative is better); p: corrected resampled t test. Net benefit averaged over the thresholds 0.1 to 0.3.

Model	RPS	ΔRPS	p	Brier	AUROC	Int.	Slope	NB
Prior	0.1615 (0.0064)	+0.0285	< 0.001	0.208	0.500 (0.00)	−0.01	0.44	0.101
Persistence	0.1329 (0.0112)	–	–	0.172	0.739 (0.05)	−0.03	1.03	0.161
Ordinal logistic	0.1326 (0.0092)	−0.0004	0.908	0.173	0.755 (0.04)	0.02	0.94	0.160
Logistic on PCA	0.1375 (0.0089)	+0.0046	0.257	0.178	0.731 (0.05)	0.02	0.98	0.152
Random forest	0.1301 (0.0098)	−0.0028	0.199	0.169	0.760 (0.04)	0.02	1.00	0.159
CatBoost	0.1303 (0.0109)	−0.0027	0.245	0.169	0.762 (0.05)	0.02	1.08	0.161
Value forecast (ridge)	0.1432 (0.0088)	+0.0103	0.048	0.186	0.756 (0.05)	−0.58	1.14	0.136
Value forecast (CatBoost)	0.1435 (0.0092)	+0.0106	0.049	0.185	0.761 (0.05)	−0.57	1.40	0.135
GRU-D	0.1275 (0.0102)	−0.0055	0.138	0.166	0.780 (0.04)	0.07	1.07	0.167
Masked LSTM	0.1296 (0.0113)	−0.0033	0.341	0.169	0.768 (0.05)	0.05	0.97	0.164
Selected (procedure)	0.1278 (0.0110)	−0.0051	0.157	0.166	0.777 (0.05)	0.07	1.03	0.165

in 36 of the 49 hospitals that contribute at least 30 targets, and the margin is the same for women and men and below and above 65 years (Table 4.7). The two linear models do not beat persistence; the logistic regression on principal components is significantly worse (+0.0052, $p = 0.034$).

Glycaemic task. No family beats persistence significantly. GRU-D has the lowest RPS (0.1275 against 0.1329, $\Delta = -0.0055$, $p = 0.138$) and the best discrimination of the alert (AUROC 0.780 against 0.739), and the procedure improves on persistence by 0.0051 ($p = 0.157$). With only 15 outer estimates of a task that falls short of the sample-size criteria (Table 4.4), a difference of this size cannot be distinguished from the variation between folds. The two value forecasts are worse than persistence (+0.010, $p = 0.048$ and 0.049): they over-predict the alert in the development data, with a calibration intercept of −0.58 and −0.57, against values close to zero for the classifiers. GRU-D improves on persistence in 5 of the 7 hospitals with at least 30 targets, and in every subgroup (Table 4.8).




Figure 4.7 Difference in RPS between each candidate and persistence in the 15 outer folds (dots) and its mean (red bar). Below the dashed line, the candidate forecasts better than the guideline rule applied to the last reading.

Table 4.7 Outer-fold results by subgroup, haemodynamic task: persistence against the deployed family (CatBoost). Each repeat pools its five outer folds; mean of the three repeats; *n*: targets per repeat. In 36 of the 49 hospitals with at least 30 targets, CatBoost has a lower RPS than persistence.

		RPS			AUROC	
Subgroup	*n*	**Persist.**	**CatBoost**	Δ	**Persist.**	**CatBoost**
All	4182	0.1380	0.1337	−0.0043	0.634	0.666
Sex F	1690	0.1491	0.1444	−0.0047	0.644	0.682
Sex M	2492	0.1305	0.1264	−0.0041	0.624	0.651
Age under 65	1283	0.1456	0.1411	−0.0045	0.673	0.701
Age 65 and over	2899	0.1347	0.1304	−0.0042	0.613	0.646

Table 4.8 Outer-fold results by subgroup, glycaemic task: persistence against the deployed family (GRU-D). Each repeat pools its five outer folds; mean of the three repeats; *n*: targets per repeat. In 5 of the 7 hospitals with at least 30 targets, GRU-D has a lower RPS than persistence.

		RPS			AUROC	
Subgroup	*n*	**Persist.**	**GRU-D**	Δ	**Persist.**	**GRU-D**
All	1356	0.1323	0.1271	−0.0052	0.724	0.778
Sex F	491	0.1399	0.1314	−0.0085	0.691	0.759
Sex M	865	0.1280	0.1247	−0.0033	0.744	0.790
Age under 65	393	0.1205	0.1185	−0.0020	0.735	0.789
Age 65 and over	963	0.1371	0.1307	−0.0064	0.717	0.771

4.2.2 Deep temporal against tabular models

On the haemodynamic task the deep models bring nothing over the tree ensembles: the masked LSTM and the random forest have the same RPS (difference 0.0000, $p = 0.99$), and GRU-D is slightly worse than both ensembles (+0.0009, $p = 0.58$). On the glycaemic task GRU-D is ahead of the random forest and CatBoost by 0.0027 and 0.0028, but not significantly ($p = 0.41$ and 0.40); it is significantly better only than the ordinal logistic regression (−0.0051, $p = 0.040$). The deep models therefore do not outperform the tabular ones on the evidence of these data, which is what H3 predicted, with the reservation that on glucose the selection procedure prefers them consistently.

4.2.3 The deployed model and the stability of the choice

In each outer fold the inner selection cost chose one family. For the haemodynamic task the choice varied: random forest in 7 folds, ridge value forecast in 5, masked LSTM in 2, CatBoost in 1. For the glycaemic task it was stable: GRU-D in 12 folds, masked LSTM in 3. Applied once more to all development data, the procedure chooses **CatBoost** for the haemodynamic task (24-hour window, depth 4, 479 trees, balanced class weights) and **GRU-D** for the glycaemic task; these are the models exported to the risk service. Table 4.9 tests the dependence of this choice on the provisional equal weights of the selection cost. GRU-D remains first under 86% of random weightings. CatBoost is first under 52% and the random forest under 29%: on blood pressure the choice is between two tree ensembles whose costs differ by 0.02, and either would serve.

Table 4.9 Stability of the choice of family, final search on all eICU: selection cost with equal weights (lower is better), and share of 1 000 random weightings (uniform Dirichlet) under which the family has the lowest cost.

	Haemodynamic		Glycaemic	
Family	**Cost**	**First**	**Cost**	**First**
CatBoost	−0.59	52%	−0.59	0%
Random forest	−0.57	29%	−0.42	2%
Masked LSTM	−0.25	6%	−0.36	0%
Value forecast (ridge)	−0.21	0%	2.80	0%
Value forecast (CatBoost)	−0.13	6%	3.84	0%
GRU-D	0.11	9%	−0.88	86%
Ordinal logistic	5.32	0%	−0.15	12%
Logistic on PCA	7.03	0%	0.68	0%

4.2.4 What the haemodynamic model uses

Interpretability is reported only for a model shown to beat persistence, so for the haemodynamic CatBoost alone. Permuting the pressure features together raises its RPS by 0.035 on eICU and by 0.050 on the Douala samples; the pulse, age and sex, and glucose each contribute less than a tenth of that (Table 4.10). The tree SHAP values say the same at the level of single features: the mean absolute contribution of the last systolic pressure is about six times that of the next feature, the last pulse, followed by the last diastolic pressure and age, while the slopes of systolic pressure and glucose contribute nothing. The model is, in effect, a refined persistence: it reads the last pressure as the rule does, and corrects it with the pulse and the patient's age. This explains why its margin over persistence is real but small.

Table 4.10 Grouped permutation importance of the deployed model, haemodynamic task (CatBoost): increase in RPS when the features of one group are permuted together, mean (SD) of 20 permutations.

Group	**eICU**	**Douala**
Pressure (systolic, diastolic, pulse pressure)	0.0350 (0.0013)	0.0499 (0.0054)
Pulse	0.0045 (0.0003)	0.0037 (0.0011)
Age and sex	0.0029 (0.0002)	−0.0019 (0.0009)
Glucose	0.0022 (0.0002)	0.0007 (0.0014)

Lead time. The benchmark was run for the next reading only. The runs with embargoes of 12 and 36 hours that H2 requires were not made, so H2 is not tested.

4.3 External Validation on the Douala Patients

The models refitted on all development data were applied once to the Douala charts: 282 haemodynamic targets of 32 patients and 190 glycaemic targets of 24 patients (Tables 4.11 and 4.12). With so few patients, the 95% intervals of the patient bootstrap overlap for every pair of models, and the comparisons below are descriptive.

Accuracy and discrimination. Every model, persistence included, has a higher RPS on Douala than in development (Figure 4.8), as the differences of prevalence and of glucose levels predicted (Section 4.1). The ordering relative to persistence holds: the deployed CatBoost has an RPS of

Table 4.11 External validation on the Douala charts, haemodynamic task (282 targets, 32 patients). 95% intervals by patient bootstrap. Calibration intercept and slope as is, then after the intercept update.

			As is		Updated	
Model	**RPS**	**AUROC**	**Int.**	**Slope**	**Int.**	**Slope**
Prior	0.1786 [0.1351, 0.2306]	0.500 [0.50, 0.50]	0.37	0.55	0.00	0.60
Persistence	0.1531 [0.1148, 0.1946]	0.772 [0.70, 0.84]	0.25	1.81	−0.00	1.81
Ordinal logistic	0.1550 [0.1146, 0.2005]	0.668 [0.48, 0.83]	0.41	0.97	0.00	0.97
Logistic on PCA	0.1579 [0.1203, 0.2023]	0.664 [0.48, 0.83]	0.17	0.95	−0.00	0.95
Random forest	0.1450 [0.1109, 0.1825]	0.783 [0.68, 0.86]	0.24	1.80	0.00	1.80
CatBoost	0.1467 [0.1104, 0.1874]	0.759 [0.65, 0.84]	0.40	1.49	0.00	1.49
Value forecast (ridge)	0.1385 [0.1096, 0.1686]	0.796 [0.70, 0.86]	0.04	1.63	0.00	1.63
Value forecast (CatBoost)	0.1390 [0.1120, 0.1669]	0.779 [0.67, 0.85]	0.03	1.80	0.00	1.80
GRU-D	0.1402 [0.1042, 0.1799]	0.766 [0.63, 0.86]	0.33	1.18	0.00	1.18
Masked LSTM	0.1405 [0.1087, 0.1752]	0.787 [0.69, 0.86]	0.20	1.54	0.00	1.54

Table 4.12 External validation on the Douala charts, glycaemic task (190 targets, 24 patients). 95% intervals by patient bootstrap. Calibration intercept and slope as is, then after the intercept update.

			As is		Updated	
Model	**RPS**	**AUROC**	**Int.**	**Slope**	**Int.**	**Slope**
Prior	0.2673 [0.2283, 0.3040]	0.500 [0.50, 0.50]	1.24	−0.18	−0.00	0.12
Persistence	0.2282 [0.1993, 0.2536]	0.661 [0.59, 0.73]	0.78	0.43	−0.00	0.43
Ordinal logistic	0.2411 [0.2057, 0.2744]	0.638 [0.56, 0.71]	0.78	0.31	0.00	0.31
Logistic on PCA	0.2442 [0.2079, 0.2787]	0.644 [0.57, 0.72]	0.71	0.28	0.00	0.28
Random forest	0.2422 [0.2100, 0.2714]	0.614 [0.54, 0.68]	0.83	0.30	−0.00	0.30
CatBoost	0.2393 [0.2092, 0.2663]	0.611 [0.54, 0.68]	0.89	0.34	0.00	0.34
Value forecast (ridge)	0.2140 [0.1876, 0.2406]	0.693 [0.62, 0.76]	0.05	0.38	0.00	0.38
Value forecast (CatBoost)	0.2156 [0.1902, 0.2395]	0.652 [0.57, 0.72]	0.08	0.43	0.00	0.43
GRU-D	0.2100 [0.1773, 0.2446]	0.716 [0.61, 0.79]	0.85	0.50	−0.00	0.50
Masked LSTM	0.2049 [0.1715, 0.2405]	0.711 [0.60, 0.79]	0.74	0.51	0.00	0.51

0.1467 against 0.1531 for persistence, and GRU-D 0.2100 against 0.2282, that is 4% and 8% lower, margins at least as large as in development. Discrimination does not fall uniformly. The AUROC of the haemodynamic alert is higher on Douala than in development for every model, persistence included (CatBoost 0.759 against 0.668, persistence 0.772 against 0.648), so the Douala alerts are easier to rank than those of intensive care; the glycaemic AUROC of GRU-D falls from 0.780 to 0.716.

Calibration. The calibration intercepts are positive for every classifier, from 0.17 to 0.41 on blood pressure and from 0.71 to 0.89 on glucose: the models under-predict the alert on Douala, as expected from a prevalence of 23% against 17%, and 59% against 29%. Updating the intercept on the local data removes this offset by construction and lowers the Brier score (CatBoost 0.148 to 0.142, GRU-D 0.237 to 0.221), but it leaves the slopes unchanged, and they are far from one: above one on blood pressure (1.49 for CatBoost, predictions too cautious), 0.3 to 0.5 on glucose (predictions too extreme). One family behaves differently: the value forecasts have intercepts of 0.03 to 0.08 in both tasks, close to zero before any update, and the lowest Douala RPS on blood pressure (0.1385 for the ridge forecast). This is consistent with how the formulation works: a value forecast passes the higher local pressures and glucose levels into its probabilities through




Figure 4.8 RPS in development (blue squares: mean of the 15 outer folds) and on the Douala charts (red: estimate and 95% patient-bootstrap interval). Lower is better.

the values themselves, whereas a classifier learns the base rate of intensive care. The formulation that was worse in development transfers its calibration best.

Clinical utility. The decision curves (Figure 4.9) do not show the margin in RPS. On blood pressure, the deployed CatBoost does not exceed persistence at any of the thresholds 0.1, 0.2 and 0.3: at 0.2, its net benefit is 0.107 against 0.120 (interval 0.030–0.207 against 0.042–0.220), and 0.088 after the intercept update. On glucose, GRU-D exceeds persistence at 0.2 and 0.3 (0.434 against 0.405, 0.395 against 0.330), but with an alert in 59% of the targets, raising an alert for every target does better still (0.487 and 0.414). On these few patients, no deployed model would have improved the alert decisions over the rule.




Figure 4.9 Decision curves on the Douala charts, probabilities as is. The net benefit counts true alerts minus false alarms weighted by the odds of the threshold.

4.4 Evolution of the Patient's State

The eight measures of Table 2.3 were compared on 3 392 development samples of 551 patients, in half of which the level of the next reading differs from that of the baseline state, and checked on 253 Douala samples of 32 patients (Figure 4.10). All the measures are weak. On the primary outcome, the development AUROC ranges from 0.52 for the centred cosine to 0.60 for the Mahalanobis distance; the measures that scale each change by the patient's own variability come within 0.005 of it. By the rule declared in advance—the simplest candidate whose interval contains the best value—the measure retained is the **raw Euclidean distance** (AUROC 0.576, interval 0.552–0.601). The cosine measures are the weakest, as the analysis of Section 2.5 anticipated: the plain cosine is at chance on Douala for the primary outcome (0.49) and below it for a new alert (0.37). On the Douala check the measures scaled by the patient's variability do best for a new alert (0.58–0.62), but the intervals of 24 patients are too wide to reverse a choice made on the development data, and Douala is not used to choose.




Figure 4.10 AUROC of the measures of state evolution with 95% patient-level intervals. Dashed: chance. The measure is chosen on eICU only.

The distance between the current state and the baseline therefore says little about the level of the next reading. It is kept in the risk card as a description of how far the patient has moved from their own recent values, not as a trigger of alerts.

4.5 System Verification

The system was verified by automated tests and by rendering its screens; it has not been used by patients or clinicians (Table 4.13). The test that matters most is the parity of the levels: the home triage is computed on the phone, in Dart, so that it works offline, and in Python by the worker and the risk service. A fixture generated by the Python reference—500 readings, the values on either side of every threshold and 11 verification profiles—is replayed by the application's tests, which must find the same level and the same profile status for every case.

Table 4.13 Verification of the system.

Component	Verification	Result
Research package	171 automated tests: labelling rule, quality controls, windows, metrics, sample size, home triage, state measures, risk service, worker	all pass
Sample-size port	published example of the reference implementation (623, 662 and 221 samples for the three criteria)	reproduced exactly
Triage parity	500 readings, threshold boundaries and 11 profiles, Python reference against the application	identical levels and statuses
Application	48 tests (record, events, queue order, assistant instruction, state evolution, parity); static analysis	all pass; no issue
Screens	28 renderings: light and dark themes, text enlarged to 130%	no overflow
Security rules	append-only readings, field and range validation, events and verdicts by role	written; deployment and emulator test pending
Health assistant	fixed set of 22 adversarial questions against the four guardrails	set defined; run on a phone pending

4.6 Discussion

4.6.1 Answers to the research questions

RQ1 asked whether the system can be built under the constraints of Cameroonian care. It was built, and each constraint is met by a design element rather than by an assumption. Intermittent connectivity is handled by the offline cache and by computing the home triage on the phone, so that a red reading brings up its warning without a network. Scarce clinician time is protected by the two-stage protocol, in which a yellow reading first asks the patient for a verification profile and only the mean of the profile reaches a clinical decision, and by a queue that puts red before yellow and the oldest first, each event closed in two or three taps. No proprietary device is needed, since any validated cuff or meter can be read by the patient. What remains open is whether patients and clinicians use it as intended, which only a pilot can measure.

RQ2 asked whether learned models forecast the level of the next reading better than the guideline rule, and whether a deep temporal model does better than classical ones. For blood pressure, yes to the first part: five of the eight families, and the selection procedure as a whole, have a significantly lower RPS than persistence, in most hospitals and in every subgroup, so **H1 is supported for the haemodynamic task**. The gain is about 3%, and the importance analysis shows why it is no larger: the model reads the last systolic pressure as the rule does and corrects it with little else. For glucose, no family beats persistence significantly, so **H1 is not supported for the glycaemic task**; the best model, GRU-D, is ahead by a margin of the same size, and the task falls short of the sample-size criteria, so the question is open rather than answered in the negative. On the second part, the deep models equal the tree ensembles on blood pressure and are ahead on glucose without a significant difference: **H3 is supported**, the deep models do not outperform the tabular ones, although the procedure prefers GRU-D for glucose in 12 of 15 folds and under 86% of the weightings of the selection cost.

RQ3, on lead times of 24 and 48 hours, is not answered: the benchmark was run for the next reading only, and **H2 is not tested**.

RQ4 asked whether models developed on another health system transfer to Cameroonian patients. Partly. The RPS is higher on Douala for every model, and the calibration intercepts show the expected under-prediction of the alert, so **H4 is supported as to the loss of performance and the calibration offset**; updating the intercept removes the offset and lowers the Brier score, but leaves the slopes far from one, so it recovers only part of the calibration. Against this, the deployed models keep or widen their margin in RPS over persistence, the haemodynamic alerts are ranked better than in development, and the value forecasts arrive almost calibrated in the large without any update. What does not transfer is clinical utility: on 32 patients, no deployed model improves the alert decisions over the rule. The predictive component should therefore run as the system was designed to run it, on top of the rule floor, and be revalidated on home data before it is allowed to raise any tier.

4.6.2 What the data imply for the transfer

The harmonisation made the two datasets alike in rhythm, not in content. Three differences remain, and each predicts a specific effect. The prevalence of the alert differs (17% against 23% for the haemodynamic task, 29% against 59% for the glycaemic task), which shifts calibration-in-the-large even for a model that ranks patients well; this is what the intercept update of H4 addresses. Critical hyperglycaemia is ten times more frequent in the Douala wards than in intensive care, so a model developed on eICU has seen few examples of the level that matters most for the glycaemic task. And glucose is measured twice as often in Douala, so the features that count glucose readings and the time since the last one have a different meaning in the two settings: in intensive care a long gap is the norm, in the ward it is not. The label uncertainty sets the expectations for all models: with more than half of the glucose readings within the device tolerance of a threshold, a glycaemic forecast will disagree with the label on many readings whatever the model.

4.6.3 Comparison with the literature

The study closest to this work reports an accuracy of 0.973 for XGBoost on blood pressure and glucose readings [9]; on similar data, Mendoza-Cano et al. [10] find the same algorithm no better than chance. The difference is explained by the task, not by the algorithm: when the label is a threshold rule applied to the reading the model sees, the model recovers the rule (Section 2.3). The present work avoids this by forecasting the next reading, so the guideline rule applied to the last reading—persistence—is the reference any model must beat, and accuracy is not reported because it rewards the majority level. Its results are therefore not comparable with those figures, and are not meant to be. The reviewed telemonitoring protocol [8] reports clinical effect, not predictive performance; the present system implements it as the rule floor, so a model can only add to it.

The arithmetic of alert volume. Whatever the benchmark shows, the prevalence of the alert at home will be far lower than in the inpatient data. With π the share of readings that need attention, a classifier of sensitivity Se and specificity Sp has a precision

$$\mathrm{PPV} = \frac{\pi\,\mathrm{Se}}{\pi\,\mathrm{Se} + (1-\pi)(1-\mathrm{Sp})}. \tag{4.1}$$

These formulas reproduce the precisions reported by Mendoza-Cano et al. [10] at their prevalence of 0.303 (0.92 against 0.94 for the random forest). At a planning prevalence of 0.02, a sensitivity of 0.90 and a specificity of 0.98 give a precision of 0.48 and 19.6 false alerts per 1 000 readings; a precision of 80% requires a specificity above 0.995. The two-stage home protocol answers this arithmetic: a single yellow reading is queued behind the red ones and leads first to more measurements, so the clinician's decision rests on the mean of a profile rather than on one value, and only red readings call for immediate review. Figure 4.11 shows how far the inpatient data used here are from that regime: the same classifier that is precise at the prevalence of the wards loses most of its precision at home.




Figure 4.11 Precision against the share of readings that need attention, at a sensitivity of 0.90. The dashed line marks a precision of 80%. Dotted lines: the planning prevalence of 0.02 for a home deployment, and the prevalence of the haemodynamic alert at the targets of eICU (0.17) and Douala (0.23).

4.6.4 Limitations

Table 4.14 states each limitation with its consequence and with what this work does about it.

Table 4.14 Limitations, their consequences and the response of this work.

Limitation	Consequence	Response
32 local patients	local estimates have wide intervals	development on eICU, validation only in Douala; patient-level intervals; read as exploratory
Demonstration subset of eICU	glycaemic task below the sample-size criteria (Table 4.4)	primary set of 8 features; benchmark to be rerun on the full database
Inpatients, not home users	prevalence and dynamics differ from home	home triage with out-of-office thresholds; model revalidated on home data before use
Labels derived from thresholds	a model may learn the rule; no clinical adjudication yet	forecasting task; physician relabelling sized at 100–150 readings; clinician verdicts in deployment
Unknown devices, hand transcription	up to 56% of readings near a threshold within device tolerance	devices inferred from digits; uncertainty reported; isolated extremes flagged
Diagnoses not transcribed	Douala cohort may include patients with neither condition	eligibility verified in eICU; Douala used for validation only
Slot hours assumed	glycaemic samples depend on the tolerance	sensitivity analysis at ± 1 and ± 3 h
Provisional choices	selection-cost weights and tier cut-points are not yet the physician's	declared in configuration; weight sensitivity; to be replaced by stated costs of errors
No use by patients	no usability or effectiveness result	verified by tests; prospective pilot as future work

Key points of Chapter 4

- The Douala pressures were taken with electronic devices and the pulse read from the chart curve; the device tolerance alone can change the level of 27 to 56% of the readings.
- The sourced labelling rule changes the level of 58 of the 366 readings compared with the earlier rule, mostly through low values and the glucose bands.
- eICU and Douala share the rhythm of two readings a day but not the content: diastolic pressure, glucose, its frequency and the prevalence of the alert differ.
- The level of the next glucose reading equals the current one in only 30% of the Douala pairs, so the glycaemic forecast is harder there than in the development data.
- The Douala charts cannot support model development; the demonstration subset of eICU supports the haemodynamic task, not fully the glycaemic one.
- On blood pressure, five of eight families beat persistence significantly, by about 3% of the RPS (H1 supported); on glucose, none does (H1 not supported). Deep models do not outperform tree ensembles (H3 supported); H2 was not tested.
- The deployed models are CatBoost for blood pressure and GRU-D for glucose. On Douala their RPS is higher than in development and their alerts are under-predicted (H4); the intercept update corrects the offset, not the slopes, and no model improves the alert decisions over the rule.
- The measures of state evolution are weak (AUROC 0.52 to 0.60); the raw Euclidean distance is retained by the declared rule, as a description for the clinician, not as an alert.
- The system passes 219 automated tests, including the parity of the triage on the phone with the Python reference; it has not yet been used by patients.

General Conclusion and Perspectives

Summary of the Work

This thesis set out to design a telemonitoring system for patients with hypertension and type 2 diabetes in Cameroon, and to evaluate its predictive component without the biases of previous work. The literature showed that telemonitoring improves blood pressure and glycated haemoglobin when a clinician acts on the data, that the only validated protocol for patients with both conditions triages single readings with fixed thresholds, and that the predictive results of the field rest on labels that leak into the features and on validation in the training population (Chapter 1). The risk-tiering problem was therefore formalised as a forecast of the next reading, labelled by official thresholds, combined with the guideline rule so that a model can raise an alert but never lower it (Chapter 2). A pipeline of data collection and treatment and a benchmark protocol were built to select the model rather than presuppose it, and the system was built around them (Chapter 3).

Principal Findings

The data work produced four results. First, the Douala charts were taken with electronic devices for the pressures, while the pulse was read off the chart curve; the readings themselves showed this, since the devices are not named. Second, the device tolerance alone can change the level of a quarter to more than half of the readings, which bounds what any model can achieve against a threshold label. Third, the sourced labelling rule changes the level of 58 of the 366 readings compared with the earlier unsourced rule, chiefly because low pressures now count and the glucose bands follow the official thresholds. Fourth, the 32 local patients are far below the size needed to develop a model, and the development data differ from them in prevalence and in the frequency of critical hyperglycaemia, which predicts a loss of calibration that a local intercept update should partly recover.

The benchmark answered the predictive questions with a mixed result. On blood pressure, learned models forecast the level of the next reading better than the guideline rule applied to the last reading: five of eight families, and the selection procedure as a whole, improve on persistence significantly, in most hospitals and in every subgroup, but by about 3% of the ranked probability score, and the deployed CatBoost relies mostly on the last systolic pressure. On glucose, no model beats persistence significantly, on a task that the demonstration data cannot fully support. Deep temporal models did not outperform tree ensembles, although GRU-D was the procedure's consistent choice for glucose. Applied to the Douala patients, every model lost accuracy and under-predicted the alert; an intercept update corrected the offset but not the slopes, the value forecasts arrived nearly calibrated without it, and no model improved the alert decisions over the rule. The measures of the evolution of the patient's state carry little information about the next level, and the one retained is shown to the clinician as a description, not used as an alert. The lead times of 24 and 48 hours were not tested.

These results justify the design choice made before any of them was known: the guideline rule is the floor, and a model can only add to it.

The system answers the first research question: a patient application and a clinician dashboard that implement the three-level home protocol offline, a record that cannot be altered, a queue that orders the clinician's attention, and server functions for escalation, reminders and verdicts, verified by 219 automated tests, among them the parity of the triage computed on the phone with the Python reference.

Future Directions

- **Full development data.** Rerunning the benchmark on the full eICU database, available under a data use agreement, removes the sample-size constraint of the glycaemic task.
- **Clinical input.** The relabelling of 100 to 150 readings by the partner physician, and the costs of errors that fix the weights of the selection cost and the tier cut-points, replace the provisional choices.
- **Home data.** A prospective pilot in Douala would collect the home readings for which the system is designed, revalidate the model on them, and measure adherence, usability and the clinician's workload; the clinician verdicts it records become labels that do not derive from the thresholds.
- **Acquisition.** Readings from Bluetooth devices would remove transcription errors, and a low-cost sensor node (ESP32 with pulse oximetry and temperature) could add covariates; neither was built in this work.
- **Deployment.** Hosting the risk service and the worker, an SMS fallback for patients without data, and a database region chosen under Law No. 2024/017 are required before any use with patients.

References

[1] World Health Organization, *Global report on hypertension: The race against a silent killer*. Geneva: World Health Organization, 2023. [Online]. Available: https://www.who.int/publications/i/item/9789240081062

[2] L. F. Gafane-Matemane et al., "Hypertension in sub-Saharan Africa: The current profile, recent advances, gaps, and priorities," *Journal of Human Hypertension*, vol. 39, no. 2, pp. 95–110, 2024. DOI: 10.1038/s41371-024-00913-6

[3] T. Hinneh et al., "Regional prevalence of hypertension among people diagnosed with diabetes in Africa, a systematic review and meta-analysis," *PLOS Global Public Health*, vol. 3, no. 12, 2023, Art. no. e0001931. DOI: 10.1371/journal.pgph.0001931

[4] World Health Organization, *Guideline for the pharmacological treatment of hypertension in adults*. Geneva: World Health Organization, 2021.

[5] O. A. Segun-Omosehin et al., "Evaluating the digital health technology landscape in sub-Saharan Africa and its implications for cardiovascular health," *npj Cardiovascular Health*, vol. 2, 2025, Art. no. 20. DOI: 10.1038/s44325-025-00055-9

[6] G. Bediang et al., "Evaluation of the effectiveness of telemedicine in the management of cardiovascular diseases in primary health care in Cameroon: An interventional study," *Studies in Health Technology and Informatics*, vol. 281, pp. 615–619, 2021. DOI: 10.3233/SHTI210244

[7] Y. Mao et al., "Impact and efficacy of mobile health intervention in the management of diabetes and hypertension: A systematic review and meta-analysis," *BMJ Open Diabetes Research & Care*, vol. 8, no. 1, 2020, Art. no. e001225. DOI: 10.1136/bmjdrc-2020-001225

[8] M. Mihevc et al., "Impact of 12-month mHealth home telemonitoring on clinical outcomes in older individuals with hypertension and type 2 diabetes: Multicenter randomized controlled trial," *JMIR mHealth and uHealth*, vol. 13, 2025, Art. no. e59733. DOI: 10.2196/59733

[9] A. Efendi et al., "IoT-based elderly health monitoring system using Firebase cloud computing," *Health Science Reports*, vol. 8, no. 3, 2025, Art. no. e70498. DOI: 10.1002/hsr2.70498

[10] O. Mendoza-Cano et al., "Data-driven identification of key predictors of uncontrolled hypertension: A cross-sectional study," *PLOS ONE*, vol. 20, no. 9, 2025, Art. no. e0331565. DOI: 10.1371/journal.pone.0331565

[11] J. E. Dodoo, H. Al-Samarraie, and A. Alsswey, "The development of telemedicine programs in Sub-Saharan Africa: Progress and associated challenges," *Health and Technology*, vol. 12, no. 1, pp. 33–46, 2022. DOI: 10.1007/s12553-021-00626-7

[12] A. O. Agbeyangi and J. M. Lukose, "Telemedicine adoption and prospects in Sub-Sahara Africa: A systematic review with a focus on South Africa, Kenya, and Nigeria," *Healthcare*, vol. 13, no. 7, 2025, Art. no. 762. DOI: 10.3390/healthcare13070762

[13] World Health Organization, *HEARTS technical package for cardiovascular disease management in primary health care: Evidence-based treatment protocols*. Geneva: World Health Organization, 2018.

[14] G. Mancia, R. Kreutz, M. Brunström, et al., "2023 ESH guidelines for the management of arterial hypertension," *Journal of Hypertension*, vol. 41, no. 12, pp. 1874–2071, 2023. DOI: 10.1097/HJH.0000000000003480

[15] S. Kingue et al., "Efficiency of an intervention package for arterial hypertension comprising telemanagement in a Cameroonian rural setting: The TELEMED-CAM study," *Pan African Medical Journal*, vol. 15, 2013, Art. no. 153.

[16] T. Alessa et al., "Smartphone apps to support self-management of hypertension: Review and content analysis," *JMIR mHealth and uHealth*, vol. 7, no. 5, 2019, Art. no. e13645. DOI: 10.2196/13645

[17] Y. Zhou et al., "Behavior change techniques used in self-management interventions based on mHealth apps for adults with hypertension: Systematic review and meta-analysis of randomized controlled trials," *Journal of Medical Internet Research*, vol. 26, 2024, Art. no. e54978. DOI: 10.2196/54978

[18] A. C. N. Bandeira et al., "Effect of m-health-based interventions on blood pressure: An updated systematic review with meta-analysis," *Telemedicine and e-Health*, vol. 30, no. 4, pp. 987–1001, 2024. DOI: 10.1089/tmj.2023.0545

[19] A. Y. Alhaddad et al., "Sense and learn: Recent advances in wearable sensing and machine learning for blood glucose monitoring and trend-detection," *Frontiers in Bioengineering and Biotechnology*, vol. 10, 2022, Art. no. 876672. DOI: 10.3389/fbioe.2022.876672

[20] C. Tai et al., "Internet of things-enabled intelligent systems for remote chronic disease monitoring," *Transactions on Emerging Telecommunications Technologies*, vol. 35, no. 4, 2024, Art. no. e4919. DOI: 10.1002/ett.4919

[21] Y. Liu and B. Wang, "Advanced applications in chronic disease monitoring using IoT mobile sensing device data, machine learning algorithms and frame theory: A systematic review," *Frontiers in Public Health*, vol. 13, 2025, Art. no. 1510456. DOI: 10.3389/fpubh.2025.1510456

[22] I. S. Liu et al., "A finger on the pulse of cardiovascular health: Estimating blood pressure with smartphone photoplethysmography-based pulse waveform analysis," *Biomedical Engineering Online*, vol. 24, no. 1, 2025, Art. no. 36. DOI: 10.1186/s12938-025-01365-w

[23] S. P. Chatrati et al., "Smart home health monitoring system for predicting type 2 diabetes and hypertension," *Journal of King Saud University – Computer and Information Sciences*, vol. 34, no. 3, pp. 862–870, 2022. DOI: 10.1016/j.jksuci.2020.01.010

[24] N. Alhammad et al., "Patients' perspectives on the data confidentiality, privacy, and security of mHealth apps: Systematic review," *Journal of Medical Internet Research*, vol. 26, 2024, Art. no. e50715. DOI: 10.2196/50715

[25] L. Mucchi et al., "Tele-monitoring system for chronic diseases management: Requirements and architecture," *International Journal of Environmental Research and Public Health*, vol. 18, no. 14, 2021, Art. no. 7459. DOI: 10.3390/ijerph18147459

[26] A. Yazdani et al., "Scalable architecture for telemonitoring chronic diseases in order to support the CDSSs in a common platform," *Acta Informatica Medica*, vol. 26, no. 3, pp. 195–200, 2018. DOI: 10.5455/aim.2018.26.195-200

[27] E. Christodoulou et al., "A systematic review shows no performance benefit of machine learning over logistic regression for clinical prediction models," *Journal of Clinical Epidemiology*, vol. 110, pp. 12–22, 2019. DOI: 10.1016/j.jclinepi.2019.02.004

[28] T. Gneiting and A. E. Raftery, "Strictly proper scoring rules, prediction, and estimation," *Journal of the American Statistical Association*, vol. 102, no. 477, pp. 359–378, 2007. DOI: 10.1198/016214506000001437

[29] B. Van Calster et al., "Calibration: The Achilles heel of predictive analytics," *BMC Medicine*, vol. 17, p. 230, 2019. DOI: 10.1186/s12916-019-1466-7

[30] E. Parimbelli et al., "Patient similarity for precision medicine: A systematic review," *Journal of Biomedical Informatics*, vol. 83, pp. 87–96, 2018. DOI: 10.1016/j.jbi.2018.06.001

[31] A. Sharafoddini, J. A. Dubin, and J. Lee, "Patient similarity in prediction models based on health data: A scoping review," *JMIR Medical Informatics*, vol. 5, no. 1, e7, 2017. DOI: 10.2196/medinform.6730

[32] X. Wang et al., "Experimental comparison of representation methods and distance measures for time series data," *Data Mining and Knowledge Discovery*, vol. 26, no. 2, pp. 275–309, 2013. DOI: 10.1007/s10618-012-0250-5

[33] R. Tennant et al., "Monitoring patients using control charts: A systematic review," *International Journal for Quality in Health Care*, vol. 19, no. 4, pp. 187–194, 2007. DOI: 10.1093/intqhc/mzm015

[34] C. C. Moore, R. Hazard, K. J. Saulters, et al., "Derivation and validation of a universal vital assessment (UVA) score: A tool for predicting mortality in adult hospitalised patients in sub-Saharan Africa," *BMJ Global Health*, vol. 2, no. 2, 2017, Art. no. e000344. DOI: 10.1136/bmjgh-2017-000344

[35] E. Tomás et al., "Effectiveness of early warning scores in predicting in-hospital mortality and adverse events in adult general-ward patients in low- and middle-income countries: A systematic review," *African Journal of Emergency Medicine*, vol. 16, no. 1, 2026, Art. no. 100943. DOI: 10.1016/j.afjem.2026.100943

[36] Royal College of Physicians, "National early warning score (NEWS) 2: Standardising the assessment of acute-illness severity in the NHS," Royal College of Physicians, London, Updated report of a working party, 2017. [Online]. Available: https://www.rcp.ac.uk/improving-care/resources/national-early-warning-score-news-2/

[37] Centers for Medicare & Medicaid Services. "Hospital harm – severe hyperglycemia (CMS871v5), electronic clinical quality measure specification," eCQI Resource Center, Accessed: Oct. 19, 2026. [Online]. Available: https://ecqi.healthit.gov/sites/default/files/ecqm/measures/CMS871-v5.1.000-QDM.html

[38] International Hypoglycaemia Study Group, "Glucose concentrations of less than 3.0 mmol/l (54 mg/dl) should be reported in clinical trials: A joint position statement of the American Diabetes Association and the European Association for the Study of Diabetes," *Diabetes Care*, vol. 40, no. 1, pp. 155–157, 2017. DOI: 10.2337/dc16-2215

[39] American Diabetes Association Professional Practice Committee, "16. Diabetes care in the hospital: Standards of Care in Diabetes—2026," *Diabetes Care*, vol. 49, no. Supplement 1, 2026. DOI: 10.2337/dc26-S016

[40] G. S. Stergiou et al., "2021 European Society of Hypertension practice guidelines for office and out-of-office blood pressure measurement," *Journal of Hypertension*, vol. 39, no. 7, pp. 1293–1302, 2021. DOI: 10.1097/HJH.0000000000002843

[41] H. Steck, C. Ekanadham, and N. Kallus. "Is cosine-similarity of embeddings really about similarity?" arXiv: 2403.05440 [cs.IR].

[42] C. G. Fraser and E. K. Harris, "Generation and application of data on biological variation in clinical chemistry," *Critical Reviews in Clinical Laboratory Sciences*, vol. 27, no. 5, pp. 409–437, 1989. DOI: 10.3109/10408368909106595

[43] P. C. Mahalanobis, "On the generalised distance in statistics," *Proceedings of the National Institute of Sciences of India*, vol. 2, no. 1, pp. 49–55, 1936.

[44] N. D. Tracy, J. C. Young, and R. L. Mason, "Multivariate control charts for individual observations," *Journal of Quality Technology*, vol. 24, no. 2, pp. 88–95, 1992. DOI: 10.1080/00224065.1992.12015232

[45] T. J. Pollard et al., "The eICU collaborative research database, a freely available multi-center database for critical care research," *Scientific Data*, vol. 5, p. 180 178, 2018. DOI: 10.1038/sdata.2018.178

[46] A. Johnson et al. "eICU collaborative research database demo (version 2.0.1)," PhysioNet.

[47] R. D. Riley et al., "Minimum sample size for developing a multivariable prediction model: Part II – binary and time-to-event outcomes," *Statistics in Medicine*, vol. 38, no. 7, pp. 1276–1296, 2019. DOI: 10.1002/sim.7992

[48] R. D. Riley et al., "Calculating the sample size required for developing a clinical prediction model," *BMJ*, vol. 368, p. m441, 2020. DOI: 10.1136/bmj.m441

[49] R. D. Riley, B. Van Calster, and G. S. Collins, "A note on estimating the Cox-Snell R^2 from a reported C statistic (AUROC) to inform sample size calculations for developing a prediction model with a binary outcome," *Statistics in Medicine*, vol. 40, no. 4, pp. 859–864, 2021. DOI: 10.1002/sim.8806

[50] J. Ensor, *Pmsampsize: Sample size for development of a prediction model*, version 1.1.3, 2023. [Online]. Available: https://CRAN.R-project.org/package=pmsampsize

[51] L. Kish, *Survey Sampling*. New York: Wiley, 1965.

[52] E. W. Steyerberg and F. E. Harrell, "Prediction models need appropriate internal, internal–external, and external validation," *Journal of Clinical Epidemiology*, vol. 69, pp. 245–247, 2016. DOI: 10.1016/j.jclinepi.2015.04.005

[53] G. S. Collins, E. O. Ogundimu, and D. G. Altman, "Sample size considerations for the external validation of a multivariable prognostic model: A resampling study," *Statistics in Medicine*, vol. 35, no. 2, pp. 214–226, 2016. DOI: 10.1002/sim.6787

[54] R. D. Riley et al., "Minimum sample size for external validation of a clinical prediction model with a binary outcome," *Statistics in Medicine*, vol. 40, no. 19, pp. 4230–4251, 2021. DOI: 10.1002/sim.9025

[55] T. E. Kottke et al., "Association of an automated blood pressure measurement quality improvement program with terminal digit preference and recorded mean blood pressure in 11 clinics," *JAMA Network Open*, vol. 5, no. 8, e2229098, 2022. DOI: 10.1001/jamanetworkopen.2022.29098

[56] G. S. Stergiou et al., "A universal standard for the validation of blood pressure measuring devices: Association for the Advancement of Medical Instrumentation/European Society of Hypertension/International Organization for Standardization (AAMI/ESH/ISO) collaboration statement," *Journal of Hypertension*, vol. 36, no. 3, pp. 472–478, 2018. DOI: 10.1097/HJH.0000000000001634

[57] *In vitro diagnostic test systems – requirements for blood-glucose monitoring systems for self-testing in managing diabetes mellitus*, ISO 15197:2013, Geneva: International Organization for Standardization, 2013.

[58] B. Iglewicz and D. C. Hoaglin, *How to Detect and Handle Outliers* (ASQC Basic References in Quality Control 16). Milwaukee, WI: ASQC Quality Press, 1993.

[59] Z. Che et al., "Recurrent neural networks for multivariate time series with missing values," *Scientific Reports*, vol. 8, p. 6085, 2018. DOI: 10.1038/s41598-018-24271-9

[60] L. Grinsztajn, E. Oyallon, and G. Varoquaux, "Why do tree-based models still outperform deep learning on typical tabular data?" In *Advances in Neural Information Processing Systems 35, Datasets and Benchmarks Track*, 2022. arXiv: 2207.08815.

[61] N. Erickson et al. "TabArena: A living benchmark for machine learning on tabular data." version 4. arXiv: 2506.16791.

[62] N. Hollmann et al., "Accurate predictions on small data with a tabular foundation model," *Nature*, vol. 637, no. 8045, pp. 319–326, 2025. DOI: 10.1038/s41586-024-08328-6

[63] P. McCullagh, "Regression models for ordinal data," *Journal of the Royal Statistical Society: Series B (Methodological)*, vol. 42, no. 2, pp. 109–127, 1980. DOI: 10.1111/j.2517-6161.1980.tb01109.x

[64] L. Breiman, "Random forests," *Machine Learning*, vol. 45, no. 1, pp. 5–32, 2001. DOI: 10.1023/A:1010933404324

[65] L. Prokhorenkova et al., "CatBoost: Unbiased boosting with categorical features," in *Advances in Neural Information Processing Systems 31*, 2018. arXiv: 1706.09516.

[66] S. Hochreiter and J. Schmidhuber, "Long short-term memory," *Neural Computation*, vol. 9, no. 8, pp. 1735–1780, 1997. DOI: 10.1162/neco.1997.9.8.1735

[67] J. Bergstra et al., "Algorithms for hyper-parameter optimization," in *Advances in Neural Information Processing Systems 24*, 2011, pp. 2546–2554.

[68] T. Akiba et al., "Optuna: A next-generation hyperparameter optimization framework," in *Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, 2019, pp. 2623–2631. DOI: 10.1145/3292500.3330701

[69] G. C. Cawley and N. L. C. Talbot, "On over-fitting in model selection and subsequent selection bias in performance evaluation," *Journal of Machine Learning Research*, vol. 11, no. 70, pp. 2079–2107, 2010. [Online]. Available: https://www.jmlr.org/papers/v11/cawley10a.html

[70] C. Nadeau and Y. Bengio, "Inference for the generalization error," *Machine Learning*, vol. 52, no. 3, pp. 239–281, 2003. DOI: 10.1023/A:1024068626366

[71] K. J. M. Janssen et al., "Updating methods improved the performance of a clinical prediction model in new patients," *Journal of Clinical Epidemiology*, vol. 61, no. 1, pp. 76–86, 2008. DOI: 10.1016/j.jclinepi.2007.04.018

[72] B. Van Calster et al., "Evaluation of performance measures in predictive artificial intelligence models to support medical decisions: Overview and guidance," *The Lancet Digital Health*, vol. 7, no. 12, p. 100 916, 2025.

[73] E. S. Epstein, "A scoring system for probability forecasts of ranked categories," *Journal of Applied Meteorology*, vol. 8, no. 6, pp. 985–987, 1969. DOI: 10.1175/1520-0450(1969)008<0985:ASSFPF>2.0.CO;2

[74] A. J. Vickers and E. B. Elkin, "Decision curve analysis: A novel method for evaluating prediction models," *Medical Decision Making*, vol. 26, no. 6, pp. 565–574, 2006. DOI: 10.1177/0272989X06295361

[75] C. Strobl et al., "Conditional variable importance for random forests," *BMC Bioinformatics*, vol. 9, p. 307, 2008. DOI: 10.1186/1471-2105-9-307

[76] S. M. Lundberg and S.-I. Lee, "A unified approach to interpreting model predictions," in *Advances in Neural Information Processing Systems 30*, 2017, pp. 4765–4774. arXiv: 1705.07874.

[77] A. L. Goldberger et al., "PhysioBank, PhysioToolkit, and PhysioNet: Components of a new research resource for complex physiologic signals," *Circulation*, vol. 101, no. 23, e215–e220, 2000. DOI: 10.1161/01.cir.101.23.e215

[78] Republic of Cameroon, *Loi n° 2024/017 du 23 décembre 2024 portant protection des données à caractère personnel au Cameroun*, Dec. 23, 2024.

[79] Republic of Cameroon, *Loi n° 2010/012 du 21 décembre 2010 relative à la cybersécurité et la cybercriminalité au Cameroun*, Dec. 21, 2010.

[80] G. S. Collins et al., "TRIPOD+AI statement: Updated guidance for reporting clinical prediction models that use regression or machine learning methods," *BMJ*, vol. 385, e078378, 2024. DOI: 10.1136/bmj-2023-078378

[81] K. G. M. Moons et al., "PROBAST+AI: An updated quality, risk of bias, and applicability assessment tool for prediction models using regression or artificial intelligence methods," *BMJ*, vol. 388, e082505, 2025. DOI: 10.1136/bmj-2024-082505

APPENDIX A

Data Dictionary

Douala charts. One row per reading, transcribed by hand from the ward observation charts without names (Table A.1). The file is not published.

Table A.1 Columns of the Douala charts file.

Column	Content	Use
`patient`	integer code assigned at transcription	grouping of the samples
`age, sex`	years; F or M	features; subgroups; ages grouped in five-year bands before sharing
`height_cm, weight_kg`	empty in every row	not used
`date`	date of the reading	time since the patient's first reading
`time`	M (morning) or S (evening, *soir*)	slot; the two slots are placed 12 h apart
`sbp, dbp`	systolic and diastolic pressure, mmHg	channels
`hr`	pulse, beats per minute	channel
`glucose_gl`	capillary glucose as written on the chart, g/L	source of the glucose channel
`glucose`	glucose in mmol/L, from g/L with 18.0156 mg/dL per mmol/L	channel
`glucose_state`	meal status; *casuel* (random) in every row	not used: in-hospital thresholds do not depend on it
`notes`	free text, 5 rows	not used
`label`	level under the earlier unsourced rule	comparison of the two rules only (Table 4.2)

eICU demonstration subset. Version 2.0.1 [46], checked against its published checksums before every run. Table A.2 lists the tables and columns read. A stay is eligible when the patient is an adult and hypertension or diabetes is recorded in the diagnoses or the past history; hypertensions of another nature (pulmonary, intracranial, portal, ocular, intra-abdominal, gestational) are excluded.

Table A.2 Tables and columns of eICU used.

Table	Columns	Use
`patient`	`patientunitstayid, uniquepid, hospitalid, age, gender`	stays, patients, hospitals; age and sex
`diagnosis`	`diagnosisstring`	eligibility
`pastHistory`	`pasthistorypath`	eligibility
`vitalAperiodic`	`noninvasivesystolic, noninvasivediastolic, observationoffset`	cuff pressure at each slot
`vitalPeriodic`	`heartrate, observationoffset`	pulse at each slot
`lab`	`labname (bedside glucose), labresult, labresultoffset`	glucose at each slot, mg/dL to mmol/L

APPENDIX B

Benchmark Configuration

The benchmark is declared in `configs/benchmark.yaml` of the research package before any run (Table B.1). Window length and class weights are searched for every candidate; the number of configurations tried per candidate and outer fold is fixed. Seed 20261019; five outer folds grouped by hospital, four inner folds grouped by patient, three repeats.

Table B.1 Search spaces and number of configurations. Ranges marked log are searched on a logarithmic scale.

Candidate	Trials	Structural	Training
All	–	window 24, 48, 72, 96 h	class weights none or balanced
Ordinal logistic	15	–	$C \in [10^{-3}, 10^{2}]$, log
PCA + ordinal logistic	15	components 1–8	$C \in [10^{-3}, 10^{2}]$, log
Random forest	20	max features $\sqrt{p}$, 0.5, 1.0; min leaf 1–50, log; depth 3, 6, 12, none	–
CatBoost	25	depth 2–8; iterations 100–1 000, log	learning rate 0.01–0.3, log; L2 1–30, log
Value forecast, ridge	15	–	$\alpha \in [10^{-3}, 10^{3}]$, log
Value forecast, CatBoost	25	as CatBoost	as CatBoost
GRU-D	20	hidden units 4, 8, 16, 32	dropout 0–0.5; learning rate 10^{-3}–0.03, log; weight decay 10^{-6}–10^{-2}, log; auxiliary loss weight 0–1
Masked LSTM	20	hidden units 4, 8, 16, 32; layers 1, 2	as GRU-D

Rerunning the results. From `telemonitor-research`, every number and figure of the thesis is produced by a script (Table B.2).

Table B.2 Scripts of the research package and what they produce.

Script	Produces
`01_douala_audit.py`	cohort description, quality flags, terminal digits, label uncertainty (Section 4.1)
`02_label_rules.py`	the earlier rule against the sourced rule (Table 4.2)
`04_sample_size.py`	criteria of Riley et al. for both datasets (Table 4.4)
`05_eicu_cohort.py`	eligibility, harmonisation and samples of eICU
`06_benchmark.py`	nested benchmark and external validation; resumes after an interruption
`07_app_parity_fixture.py`	the fixture replayed by the application's parity tests
`08_figures.py;`	figures and tables of a saved run
`13_results_tables.py`	
`09_kappa_sample_size.py`	size of the physician's relabelling
`10_sensitivity_flags_slots.py`	shrinkage constant and slot tolerance (Table 4.3)
`11_export_model.py`	the model served by the risk service
`12_state_comparison.py`	benchmark of the measures of state evolution
`14_thesis_data_figures.py`	data of Figures 3.3, 4.4 and 4.5

APPENDIX C

Record Schema

The fields of a reading and of an event in Cloud Firestore (Figure 3.9). The security rules validate every field on the server.

Table C.1 Fields of a reading, `users/{uid}/readings/{id}`.


Field	Type	Meaning
`recordedAt`	time	measurement time stated by the patient, at most 30 days before arrival
`timestamp`	time	arrival time, set by the server
`entryMode`	text	manual (Bluetooth entry is specified, not built)
`sbp, dbp, hr`	numbers	retained reading of the protocol pair; mmHg and beats per minute
`glucose, context`	number; text	mmol/L; fasting, before a meal, after a meal or random, required with glucose
`readingCount`	number	1 or 2 readings in the session
`firstSbp, firstDbp, firstHr`	numbers	first reading of the pair, kept for audit
`supersedes`	reference	the reading this one corrects
`notes`	text	free text of the patient
`qualityFlag`	text	set by the worker: ok, out_of_range, unit_suspect, duplicate, carried_forward, extreme

Table C.2 Fields of an event, `users/{uid}/events/{id}`.


Field	Type	Meaning
`patientId`, `clinicianId`	references	the patient and the assigned clinician
`triggerSource`	text	rule, model or non_transmission
`tier`, `score`	text; number	yellow or red; the model's probability when the model raised it
`readingId`, `trigger`	reference; map	the triggering reading and values, or the profile and its means
`thresholdVersion`, `modelVersion`	text	versions in force when the event was raised
`tsTriggered`, `status`	time; text	creation; open or closed
`verdict`, `action`, `comment`	text	confirmed, not_significant or uncertain; none, advice, treatment_change, consultation or referral
`tsReviewed`, `reviewedBy`	time; reference	closure of the event, once
`escalatedAt`, `escalatedTo`	time; reference	escalation to the named alternate

APPENDIX D


Application Screens




(a) Sign-in




(b) Enrolment




(c) Patient home




(d) New reading




(e) Correction



(f) History



(g) Clinician queue



(h) Nearby care

Figure D.1 Screens of the application: (a) sign-in, shared by patients and clinicians; (b) enrolment with the clinician's code; (c) patient home with the level, the instruction and the verification profiles; (d) entry of the two readings of the protocol; (e) correction, a new reading that supersedes the old one; (f) history with level and corrections; (g) clinician queue, red before yellow; (h) nearby care, opened in the phone's maps. Rendered by the widget tests with fictitious patients; model names and probabilities are test values.

APPENDIX E

TRIPOD+AI Checklist

The modelling study is reported according to TRIPOD+AI [80]; its risk of bias can be assessed with PROBAST+AI [81].

Table E.1 TRIPOD+AI items and where they are addressed.

Item	Topic	Where
1	Title	title page
3a–c	Background, intended use, inequalities	Introduction; Section 1.9 (Central Africa absent from the evidence)
4	Objectives	Introduction (objectives, research questions, hypotheses)
5a–b	Data sources, dates	Section 3.2: Douala 2026; eICU 2014–2015; Appendix A
6a–b	Setting, eligibility	Sections 3.2 and 3.2.3; Appendix A
6c	Treatments	not recorded in either dataset
7	Data preparation	Section 3.3
8a–c	Outcome, assessment, blinding	Section 2.2; blinded relabelling by the partner physician planned (100–150 readings), not yet done
9a–c	Predictors	Section 3.4; instruments in Section 3.3.2
10	Sample size	Sections 3.2.2 and 4.1.6
11	Missing data	Section 3.3.5
12a–g	Partitioning, handling, model type, clusters, measures, updating, computation	Sections 3.5 to 3.7 and 3.9; outer folds by hospital; intercept update; Appendix B
13	Class imbalance	class weights and their correction (Section 3.6)
14	Fairness	results by sex, age band and hospital (Section 4.2)
15	Output, thresholds	Section 2.4
16	Development against evaluation data	Section 4.1
17	Ethical approval	Section 3.10
18c–f	Protocol, registration, sharing	protocol fixed before results; not registered; Section 3.9
19	Patient and public involvement	none
20–27	Results, discussion	Chapter 4