SAE decoder spectra largely inherit the activation covariance

Ayur Pulle

November 2026

Abstract

The power-law eigenspectrum of SAE decoder directions reported by Li, Michaud et al. (2024) is real, and in the top ∼150 directions most of it is inherited from the activation covariance Σ. We measure the decoder's eigen-slope relative to Σ's as γ. In ranks 11–150 at layer 13, an SAE trained on a Gaussian with the same Σ reproduces real γ at $k \geq 40$, and a decline over k = 8–128 of similar size (-0.088 against -0.097 per doubling). A dictionary drawn to match only the covariance of Gemma Scope's directions gives about half to three-quarters of the decline under Gemma Scope's firing statistics. Real activations differ from such nulls in the top 50 directions, at small k, and above all in the bulk of the spectrum (ranks 31–1000), where the nulls pile decoders into Σ's top subspace. Two regularities are kept apart below. In ranks 11–150, γ is about 1 with one active latent per token and falls about 0.1 per doubling of L0 in Gemma-2-2B. In ranks 101–1000, the decoder covariance is approximately Σ^γ as a matrix, though much less co-diagonal with Σ than an exact Σ^γ cloud. Frequency-driven allocation fits planted circle features at k = 1 but does not separate them from point features.

1 Verdict

We asked whether the power-law eigenspectrum of SAE decoder directions is predicted by frequency-driven latent allocation (slope $\approx (1+\alpha)/(1+b)$, Michaud, Gorton, McGrath 2025) or is an estimator artifact. The best-supported answer is a third one: in ranks 11–150 the spectrum is mostly Σ passed through the SAE objective, so in that window the slope Li et al. reported is largely set by the activation covariance and L0. It is not a pure artifact: real activations differ from Σ-matched nulls in ranks 4–50, at small k, and most of all in the bulk windows. Allocation fits planted circles at k = 1 but does not distinguish them from point features, and it fails per decoder cluster on Gemma. The mid-stack peak in the top windows goes with cluster structure beyond what the covariance implies; it holds for Gemma Scope and our 16k SAEs but, by our window rule, not for our 65k SAEs. A table near the end summarises every hypothesis tested.

2 Related work

Li, Michaud et al. (2024) reported the power-law spectrum of SAE decoder directions and its mid-stack peak, measured on Gemma Scope SAEs (Lieberum et al., 2024). Michaud, Gorton and McGrath (2025) model SAE scaling through the allocation of latents to features of different frequencies, which predicts a spectral slope of about $(1+\alpha)/(1+b)$. Peter and Søgaard (2025) report

latent-frequency exponents steeper than theirs. Our own SAEs follow the TopK recipe with the AuxK loss of Gao et al. (2024). We did not survey the literature systematically, so other work relating the decoder spectrum to the activation covariance may exist.

3 Reproduction

We reproduced Li et al.'s slopes to within 0.02% at layers 0, 12 and 24 of Gemma-2-2B, using the computation in Li et al.'s released notebook (below). In the table, "Paper" is Li et al.'s value.

The original computation, from `galaxy/PointCloudStructure_Galaxy.ipynb` in ejmichaud/feature-geometry:

- SAEs: `gemma-scope-2b-pt-res`, width 65k, one L0 per layer (73, 72 and 124 at layers 0, 12 and 24).
- Point cloud: raw `W_dec` rows (65,536 × 2,304). Gemma Scope rows are already unit norm.
- Covariance: centred, ddof = 1. Eigenvalues sorted descending.
- Fit: least squares of $\log_{10}$ eigenvalue on $\log_{10}$ rank, ranks 11–150 (about 1.1 decades). The null was an N(0, 1) matrix of the same shape.

Layer	Paper	Ours	R^2	Matched isotropic Gaussian
0	−0.2383	−0.2383	0.999	−0.0276
12	−0.4684	−0.4684	0.994	−0.0276
24	−0.2545	−0.2545	0.999	−0.0276

Centring, row normalisation and dead-latent exclusion each change the slope by ≤ 0.002. The fit window matters far more: the spectrum is curved, and moving the window one decade shifts the slope by up to 0.13. Every result below therefore reports four pre-registered windows (ranks 4–50, 11–150, 31–450, 101–1000) and counts a conclusion only if it holds in the primary window and at least two others.

4 What sets the slope: L0, then layer, barely width

Across 288 existing Gemma Scope SAEs, the slope is set mainly by L0 (average active latents per token), then by layer; width matters little at the top of the spectrum.

L0. At layer 12, slope is nearly linear in log L0 at every width from 16k to 1m: about −0.74 at L0 ≈ 12, −0.47 at L0 ≈ 72, −0.24 at L0 ≈ 445, roughly +0.10 per doubling. Li et al.'s −0.47 is one point on this line.

Width. At matched L0, width moves the top-window slope by ≤ 0.05 over six doublings (16k to 1m). In the deeper windows it keeps flattening with no plateau (ranks 101–1000 at L0 ≈ 40: −0.59 at 16k, −0.45 at 1m). "The slope converges with width" holds in only 2 of the 4 windows.




Figure 1: Layer-12 decoder eigen slope against L0, one line per width (16k to 1m), in four rank windows.

Layer, at matched L0. The mid-stack peak survives L0 matching in the top windows, peaking at layers 13–15 (L0 = 40: −0.34 at layer 0, −0.57 at layers 13–15, −0.34 at layer 23, 16k). The bulk windows show the opposite trend: steepest at layers 2–7, shallower after. The peak's amplitude shrinks with L0 (0.23 at L0 = 40, 0.06 at L0 = 140), so layer and sparsity interact. Li et al.'s uneven L0 choice overstated the late-layer contrast by about 0.06.




Figure 2: Decoder eigen slope against layer, interpolated to L0 = 20, 40, 70 and 140, at 16k and 65k, in four rank windows; stars mark the SAEs Li et al. used.

5 Real data against nulls: where they differ

SAEs trained on structureless data with the same covariance reproduce part of the real geometry, not all of it. The two differ most in the bulk windows, where the nulls over-concentrate the cloud; at Li et al.'s operating point in the top window they nearly agree.

We trained identical TopK SAEs on three inputs: real activations; a Gaussian $\mathrm{N}(\mu, \Sigma)$ with μ and Σ estimated from them; and scrambled activations (each coordinate permuted independently across tokens, keeping its marginal but destroying correlations). Scrambling destroys Σ's off-diagonal

structure, so scrambled runs are compared against diag(Σ), not Σ. A random rotation is not a null here, since the geometry is rotation-invariant.

Pythia-70m, layer 3 (k = 32, 15M tokens, 3 seeds). Slope, ranks 6–50:

Width	Real	Gaussian $\mathbf{N}(\mu, \Sigma)$	Scrambled
1k	-0.311 ± 0.007	-0.299 ± 0.001	-0.248 ± 0.014
4k	-0.329 ± 0.006	-0.350 ± 0.004	-0.220 ± 0.013
16k	-0.322 ± 0.007	-0.391 ± 0.004	-0.249 ± 0.006

Gemma-2-2B, layers 3, 13, 23 (k = 40, 28.5M tokens, 16k and 65k, 3 seeds, H100). Seed-to-seed sd is ≤ 0.015 on every value. Our real-data SAEs reach eval FVU 0.18–0.29 and give $\gamma = 0.43$ at layer 3, against 0.41 for Gemma Scope at the same layer and L0, so a different architecture and recipe reproduces the geometry there.

- In the bulk (ranks 31–1000), the Gaussian and scrambled SAEs pile decoders into Σ's top subspace (γ 1.1–2.7). Real data spreads them out (γ 0.29–0.81), at every layer and width. This is the largest real-vs-null difference in the project.
- The SAE objective applied to Σ alone does generate most of the slope's magnitude: the Σ-matched Gaussian already gives -0.3 to -0.4 in Pythia, against -0.13 for an isotropic Gaussian of the same shape. In Pythia, what real structure adds is width stability, not steepness; our Gemma runs are not more width-stable than the null (below).

Confounds checked. The nulls do not differ because they keep fewer latents: they have 0–1 dead latents out of 16k or 65k, against 2–1,385 for real data, and every run, real or null, has $\leq 0.4\%$ near-duplicate latents (cosine > 0.5). Part of the deep-window width drift is estimation: cutting 1m-wide decoders to 16k random rows moves the ranks 101–1000 slope 30–50% of the way toward the 16k SAEs' value. The rest is geometry.

Where the nulls match, and width. The difference is scoped. In the top window the Gaussian null nearly reproduces real γ at layer 13 for $k \geq 40$ (0.536 against 0.509 at $k = 40$, 0.382 against 0.354 at $k = 128$; see the mechanism section). The separation lies in the bulk windows, at $k \leq 16$, and at layers 3 and 23. Real γ is also not more width-stable than the null in our Gemma runs: from 16k to 65k in ranks 11–150 it falls $0.430 \to 0.364$, $0.508 \to 0.387$ and $0.389 \to 0.315$ at layers 3, 13 and 23, against $0.627 \to 0.562$, $0.533 \to 0.485$ and $0.677 \to 0.663$ for the Gaussian (table in the mid-stack section). Width stability rests on Pythia's three widths. Gemma Scope's own drift of at most 0.05 over six doublings suggests our 65k runs (28.5M tokens) are undertrained.

6 The mechanism: Σ raised to a sparsity-set power

Two regularities, measured in different regions. In the top window (ranks 11–150), rank-window γ, the ratio of the two sorted spectra's slopes, is about 1 at L0 = 1 and falls about 0.1 per doubling of L0. In the bulk (ranks 101–1000), the statement holds approximately as a matrix one: in Σ's eigenbasis, the decoder cloud's variances scale as Σ's eigenvalues to a power γ at every L0; in the

top directions this holds only at low L0. The equation below summarises both, but its two halves come from different ranks.

$$\lambda_i^{\mathrm{dec}} \propto \left(\lambda_i^{\Sigma}\right)^{\gamma(L_0)}, \qquad \gamma \approx \gamma_1 - c \log_2 L_0$$

The matrix test. Two slopes can agree by coincidence, so we tested the statement as a matrix one. We rotated the decoder covariance into Σ's eigenbasis, checked how close to diagonal it is there, regressed its log diagonal on $\log \lambda_\Sigma$, and measured the overlap of the two top eigenvector subspaces. Over ranks 101–1000 the power law holds at every L0 (R^2 0.94–0.99). In the top directions it holds only up to L0 $\approx$ 85. Above that, γ measured in the basis drops to about 0 and the overlap of the two top-10 subspaces falls to 0.09–0.13: the decoders stop tracking Σ's leading directions. Planted data shows the same decoupling at the same ratio k/d $\approx$ 0.06–0.12. Pythia-70m's decoders ignore Σ's top direction, which carries 36% of the variance.

Limits of the matrix test. We benchmarked it on the layer-12 16k SAEs against two synthetic clouds of the same size (16,384 unit rows): an exact Σ^γ cloud, and the decoder's own eigenvalues in a random basis. The R^2 is not automatic: in a random basis it is about 0, against 0.93–0.99 for the real decoders at L0 $\leq$ 82, and 0.94–0.97 in ranks 101–1000 at every L0. But the law is only approximate. The rotated decoder covariance puts 0.62–0.74 of its weight on the diagonal, against 0.89–0.95 for an exact Σ^γ cloud and 0.24–0.62 for the random basis; at high L0 the real value is close to the random one. The top-10 subspace overlap falls from 0.51 to 0.11 as L0 rises, while an exact cloud keeps 0.86–0.98, so the decoupling is not a sampling effect. The test and the law also cover different regions: the 0.1-per-doubling law is rank-window γ in ranks 11–150, while the matrix test holds at every L0 only in ranks 101–1000, and even at low L0 basis γ sits 0.08–0.12 below rank γ (0.65 against 0.73 at L0 = 22). The bulk basis γ also inherits the 200k-token Σ's bias (next paragraph).

Σ already carries the layer profile. We measured Σ at all 26 Gemma layers (200k tokens). Its slope reproduces the decoder profile window for window, including the reversal: a mid-stack hump in the top windows (-0.72 at layer 0 to -0.93 at layer 14) and an early-layer peak in the bulk (-1.15 at layers 2–6). Decoder slope correlates moderately with Σ slope across layers (r = $+0.5$ to $+0.8$), and γ = decoder slope / Σ slope averages 0.53 at L0 = 40 and 0.45 at L0 = 70 in every window.

The 200k-token Σ is too steep in the bulk. 200k tokens at context 1,024 is about 200 documents. At layer 12, Σ from 1M tokens, or from either half of them split by document, is flatter in the bulk windows (ranks 31–450: -0.73 against -0.81; ranks 101–1000: -0.72 against -0.84), and the two halves agree to 0.002. The top windows move by at most 0.03. So every bulk-window γ computed against the 200k Σ (the Gemma Scope analyses, this layer profile, the bulk basis γ) is 10–17% too low at layer 12, and the bulk part of the layer profile is uncertain. Our own SAEs' γ used each training set's Σ (at least 10M tokens) and is unaffected. Only layer 12 could be rechecked locally.

At layer 12, γ depends on L0 almost alone. Across all 28 layer-12 Gemma Scope SAEs (16k to 1m), $\gamma = 1.245 - 0.114 \cdot \log_2$ L0 in ranks 11–150, $R^2 = 0.984$; the width spread at fixed L0 is ≤ 0.07 against an L0-driven range of 0.58. Rescored on 16,384 random decoder rows per SAE, so every width has the same number of points, it is $1.207 - 0.110 \cdot \log_2$ L0 (R^2 0.981). Across layers this does not hold: at fixed L0, γ still varies with layer (see the mid-stack section).

Measured directly, with our own SAEs (layer 13, 16k, k = 1 to 128, 2 seeds). γ in the primary




Figure 3: Σ eigen slope against the decoder slope across the 26 layers (top), their scatter (middle), and γ by layer (bottom), in four rank windows.




Figure 4: γ against L0 at layer 12 for seven widths, with log-linear and power-law fits, in four rank windows.

window:

k (= L0)	**1**	**2**	**4**	**8**	**16**	**40**	**128**
Real	1.034	0.893	0.824	0.732	0.665	0.509	0.354
Gaussian N(μ, Σ)	1.248	1.292	1.231	0.826	0.448	0.536	0.382

Real data follows $\gamma = 1.016 - 0.094 \cdot \log_2 k$ ($R^2 = 0.994$). The functional form matches Gemma Scope's; the intercept depends on the training recipe. The fit holds in the two top windows (R^2 0.95 in ranks 4–50, 0.99 in 11–150) but not in the deeper two (0.76 and 0.36). By our own rule it is therefore a top-window law. The Gaussian null does not follow it at small k: it sits near 1.25 for $k \leq 4$, then drops abruptly, and it is not monotone (0.448 at k = 16, 0.536 at k = 40), which we have not traced. At $k \geq 40$ it is close to real data. At $k \leq 8$ most latents die (672 of 16,384 alive at k = 1), so only the top windows are well defined there.

Live-count bias. Measured γ depends on how many points the cloud has next to d. A finite cloud adds sampling spread to every eigenvalue, which pulls a high γ down and pushes a γ near zero up. In simulation at d = 2,304, a cloud with true $\gamma = 0.5$ measures 0.32 at 672 points, 0.41 at 4,786 and 0.47 at 16,384, while an exactly isotropic cloud measures 0.10 at 4,786 points and 0.06 at 16,384. The live count rises with k (672 at k = 1 here), so at small k the bias works against the decline: the true fall from k = 1 is likely steeper than measured. Rescoring the 28 Gemma Scope layer-12 SAEs on a fixed 16,384 rows leaves their L0 law almost unchanged (above), and in the resynthesis control a fixed 8,192 live latents moved the k-slopes by at most 0.002. Our real-data k sweep has not been rescored, because its decoders were not kept.

An empirical fit, not a zero-parameter law. The fit lies within 0.034 of $1 - \ln k / \ln d$ with d = 2304 at every k, which suggested $\gamma = \log_d(d/k)$. A dimension sweep rules that out. We projected the same layer-12 activations onto their top d' principal components and trained at k = 8 and k = 32. The gap between the two k values is 0.01, 0.06, 0.13 and 0.18 at $d' = 256$, 512, 1024 and 2304, where the law predicts 0.25, 0.22, 0.20 and 0.18. The $d' = 2304$ control reproduces the long GPU runs (0.70 vs 0.73 at k = 8; 0.52 vs 0.51 at k = 32), so the training budget is adequate. Truncation squeezes the same features into fewer dimensions, so it is not the same as a model with smaller d; Pythia (d = 512, γ 0.33 against 0.44 predicted) points the same way. We report the k-law as an empirical fit at d = 2304.

A likely reason $\gamma = 1$ at k = 1. With one latent per token, each decoder direction is a centroid of the activations it reconstructs, so the cloud samples Σ's directions and inherits its spectrum. With more latents per token, each explains a residual left by the others, pushing decoders into lower-variance directions and flattening the cloud. This is our reading, not a derivation. Planted data suggested a second route to low γ, frequency-dependent allocation of latents to features, but point features reach the same γ without it (next section).

7 Planted features: allocation and γ

Synthetic activations with known features let us match latents to features and test the allocation theory directly, which is impossible on Gemma. Each token is a sparse sum of planted features with frequencies $p_i \propto i^{-1.5}$, plus small noise, and every dataset has a Gaussian twin with the same Σ. TopK SAEs, width 4,096, d = 256, one seed; γ over ranks 6–50.

Dense regime: too Gaussian. 4,096 features, 32 active per token, directions crowded into about 37 effective dimensions. γ(k) is log-linear and close to $1 - \ln k / \ln 256$ at k = 16–64, but the Gaussian twin behaves almost identically, and at k = 4–32 at most 13% of latents match a feature. One result carries over: the overlap between the top-10 subspaces of the decoder cloud and Σ collapses between k/d = 0.06 and 0.12. Gemma Scope's layer-12 decoders stop tracking Σ at the same ratio (L0 $\approx$ 150–270).

Sparse regime: features recoverable. 512 features, 4 active per token, random directions, so Σ's spectrum comes from the frequencies. If feature i gets n_i latents, the decoder cloud's eigenvalues scale as n_i and γ equals the exponent c in $n_i \propto p_i^c$. For circle features (each a ring in a 2-D plane), tiling a ring with n latents leaves error $\propto n^{-2}$, so the allocation law predicts c = 1/3 at k = 1 (our derivation). Point features need one latent each, so c $\approx$ 0.

k	1	2	4	8	16	32
Circles: allocation exponent c, top 50 features	0.30	0.32	0.23	0.11	0.07	−0.01
Circles: c, all covered features	0.33	0.29	0.26	0.21	0.28	0.35
Circles: γ	0.32	0.21	0.19	0.12	0.08	0.08
Circles, Gaussian twin: γ	1.46*	1.33	0.88	0.58	0.42	0.31
Points: allocation exponent c, top 50 features	0.46*	0.44	0.11	−0.02	−0.14	−0.14
Points: γ	0.30*	0.20	0.16	0.06	0.04	0.06
Points, Gaussian twin: γ	1.25*	1.20	0.88	0.64	0.45	0.33

* Fewer than 1,000 live latents.

- **The prediction holds at k = 1 for circles, but γ does not discriminate.** Over all covered circle features c = 0.334, against 1/3 predicted (the top 50 give 0.30), and γ is 0.32–0.35 in all three windows. But point features, where allocation predicts c $\approx$ 0, give the same γ (0.30 at k = 1 and 0.20 at k = 2, against 0.32 and 0.21 for circles), and their top-50 c is 0.46. A matching γ is therefore not evidence of tiling.
- **Allocation and γ part ways with k, depending on the estimator.** Top-50 c falls with γ (0.30 $\rightarrow$ −0.01, against γ 0.32 $\rightarrow$ 0.08), but c over all covered features stays at 0.21–0.35 through k = 32. The five most frequent circles get 11–25 latents at k = 1 and 6–9 at k $\geq$ 8. So "γ follows allocation" holds only for the top-50 estimator.
- **The Gaussian twin has a steeper k-law with no features at all:** $\gamma = 1.35\text{–}1.45 - 0.22\text{–}0.25 \cdot \log_2 k$, against $0.22\text{–}0.29 - 0.04\text{–}0.05 \cdot \log_2 k$ for planted data. A k-law alone is not evidence for allocation.
- **Gemma's intercept sits between the two regimes; its k-slope does not.** Its fit ($1.02 - 0.094 \cdot \log_2 k$ at layer 13) has an intercept between the sparse and dense planted data, but it falls with k about half as fast as any planted run at a similar intercept.
- Caveats: one seed and one width. Up to 62% of live latents in the point runs match no feature even at cosine 0.3; they fire rarely and mostly fit the noise, adding a near-isotropic floor to the cloud.

Sweeps over density, width and dimension (details in the appendix). Across every planted setting, $\gamma(k) \approx A \cdot (1 - \log_2 k / \log_2 k_0)$: density and d set the amplitude A, while width and d

move k_0 only slowly. At Gemma's d and width, the densest setting runs parallel to Gemma in ranks 31–450. But once Σ is matched to Gemma's power law, no planted setting reproduces Gemma's $\gamma(k)$ in ranks 11–150: the best RMS against Gemma is 0.27 with random directions and 0.24 with planted clusters, against a pre-registered 0.15.

8 What does not explain it

Three candidate explanations failed direct tests on Gemma. For the first, frequency allocation, the Gemma test is underdetermined rather than decisive (see the caveat below); the planted-data section tests it directly.

Candidate	**Test**	**Result**
Frequency-driven latent allocation (slope $\approx (1+\alpha)/(1+b)$)	Firing frequencies of 22 layer-12 SAEs on 1M Pile tokens	The rank-frequency exponent stays at -0.6 to -0.8 while the eigen slope moves from -0.24 to -0.75; b would have to carry all the L0 dependence
Feature splitting: near-parallel latents act as large spikes	Greedy merge of latents with cosine > 0.5, then refit	Up to 34% of latents removed (131k, L0 = 12); slope changes by ≤ 0.010 in all 22 SAEs
Massive activations (residual dim 1170)	Remove latents aligned with dim 1170; separately, train without the tokens carrying its top 0.1% or 1% of values	Slope changes by 0.000 and γ by ≤ 0.005

Two caveats. The frequency test uses latent frequencies, while the allocation theory concerns feature frequencies p_i, so on Gemma it leaves that theory underdetermined rather than refuted; the planted-data test measures p_i directly. Near-duplicates do become more common at low L0 (2% to 59% of latents with a neighbour above cosine 0.5), but the merge test shows they carry none of the anisotropy.

The latent-frequency curve is not a single power law: piecewise exponents run from -0.4 to -1.2 depending on the rank window, which accounts for the disagreement in the literature (-0.57 to -0.74 in Michaud et al., arXiv:2509.02565 against -1.0 to -1.2 in Peter and Søgaard, arXiv:2505.00190).

9 The mid-stack effect

At fixed L0, γ itself rises toward the middle of the stack in the top windows, and Σ does not explain that rise: there, a Gaussian with the same Σ moves the opposite way.

For Gemma Scope (65k, L0 = 40, ranks 11–150), γ is about 0.40 at layers 0–3, about 0.65 at layers 12–16 and about 0.43 at layers 23–24. None of Σ's shape statistics track this (top-eigenvalue share $r = -0.01$, participation ratio $r = -0.33$).

Not an L0 mismatch. We measured L0 for all 135 Gemma Scope 16k SAEs on our tokens. The names understate it mid-stack (measured over named 1.08–1.11 at layers 10–15, 0.92–1.03 elsewhere), so at matched true L0, mid-stack γ is slightly higher than the name-based comparison

showed. The peak's height above the edge layers rises from 0.19 to 0.21 (L0 = 40, primary window), and no layer moves by more than 0.023.

At equal budget, γ of our SAEs (mean of 3 seeds, sd $\leq$ 0.006):

γ, ranks 11–150	Layer 3	Layer 13	Layer 23
Real, 16k	0.430	0.508	0.389
Gaussian N(μ, Σ), 16k	0.627	0.533	0.677
Real, 65k	0.364	0.387	0.315
Gaussian N(μ, Σ), 65k	0.562	0.485	0.663




Figure 5: Equal-budget results: γ by layer for real, Gaussian and scrambled SAEs (left); γ against k at layer 13 (centre); γ after removing massive-activation tokens (right).

- In the two top windows (ranks 4–50 and 11–150), the Gaussian null dips at layer 13 and the real-minus-Gaussian gap is largest at layer 13, at both widths. In the bulk windows neither holds: the null is flat across these layers and layer 3's gap is largest.
- Real γ itself peaks at layer 13 in three of four windows at 16k, but only in two at 65k (ranks 11–150 and 101–1000; in ranks 4–50 it is 0.305, 0.291 and 0.375 at layers 3, 13 and 23, and in 31–450 0.497, 0.489 and 0.292). By our window rule the peak in real γ holds at 16k but not at 65k. Our TopK peak (0.51 at 16k) is also smaller than Gemma Scope's JumpReLU peak (0.65): the size of the effect depends on the SAE recipe.
- In an earlier comparison over seven layers, the gap between Gemma Scope's γ and short Gaussian-null TopK runs (width 4,096, 3M tokens) tracked the kurtosis of the massive-activation dimension (r = 0.93). That comparison mixes recipes and is not the equal-budget one in the table, and seven points on one smooth profile give a weak correlation. Removing those tokens, or the latents aligned with them, changes nothing, so the heavy tail at most marks the mid-stack regime; it does not cause the effect.

10 Cluster structure in the decoder cloud

The planted experiments left a gap: random planted directions give decoders much flatter than Σ, so Gemma's top-window geometry needs structure in the decoder cloud itself. We clustered Gemma

Scope decoder directions (spherical k-means, Ward, average linkage, spectral) and compared each layer's cloud with a Gaussian cloud of the same covariance, unit-normalised and clustered the same way. Only the excess over that matched cloud counts as cluster structure, because a more anisotropic cloud makes any clustering look tighter. All of this ran locally on the 16k SAE nearest L0 = 70 at each of the 26 layers, plus four widths at layer 12.

Cluster structure beyond anisotropy tracks the top-window layer profile, not the bulk. Layer 0's cloud has no cluster structure beyond its covariance. The excess rises over layers 1–4, peaks at layers 13–15, falls through layer 23 and jumps again at layer 25, the same shape as the slope in ranks 4–50 and 11–150. Across the 26 layers, the excess between-cluster share and the excess cosine to the cluster centre correlate with the top-window slope at Spearman -0.87 to -0.91 (1,024 clusters; -0.78 to -0.90 at 256). Mid-stack clusters are also lower-dimensional than the matched cloud's: at 16 latents per cluster, their participation ratio sits up to 0.75 below it (14.1 against 14.8, maximum 15). In the deep windows (ranks 31–450, 101–1000) the excess does not track the slope ($|\rho| \leq 0.45$), while the matched cloud's own statistics do (ρ -0.88 to -0.96). This splits the two layer profiles found earlier, top windows against bulk: the bulk follows anisotropy alone, and the mid-stack peak in the top windows goes with genuine cluster structure.

Clusters carry only a minority of the top eigenvalues. The covariance splits exactly into between-cluster and within-cluster parts. At layer 12 with 16 latents per cluster, the between-cluster part explains 27%, 19%, 14% and 12% of the eigenvalues in the four windows. A covariance-matched cloud gives 14%, 10%, 8% and 7%, and isotropic unit vectors about 8% everywhere. Within-cluster spread still carries most of the slope (ranks 11–150: -0.33 within, against -0.42 overall). So the first-order picture of eigenvalues as cluster spikes accounts for under a third of any window. That is consistent with the near-duplicate test: the anisotropy is cloud-wide. Our own TopK SAEs at layer 12 (4,096 latents) show almost no difference between real and Gaussian-twin training data (12% against 10% in ranks 11–150).

The allocation law does not hold per cluster. Allocation predicts that the mean latent frequency in a cluster rises with the cluster's latent count (log-log slope $b > 0$). At layer 12, with 1M-token frequencies at widths 16k to 131k, the slope's sign depends on cluster scale. With 256 latents per cluster it is -0.57 to -1.52 (z -9 to -27 against a frequency-shuffle null): large clusters are made of rarer latents, the opposite of the prediction. Only with 16 latents per cluster at 65k and 131k is it positive ($+0.21$ to $+0.28$). Correlations are weak ($|r| \leq 0.5$), and no single b fits.

Caveats. Clusterings themselves are unstable: k-means against itself across seeds gives an adjusted Rand index of only 0.08–0.17, and methods agree no better. The cloud is a continuum at these scales, and only the summary statistics are robust across methods. The layer result is a correlation over 26 points on one smooth profile. The matched cloud reproduces the real spectrum within 0.02–0.05 in slope. Frequencies were measured only at layer 12.

11 Resynthesis: how much the covariance explains

Planted data never reproduced Gemma's $\gamma(k)$ in ranks 11–150, so we built test data from Gemma itself. We encoded 12M layer-12 tokens with the Gemma Scope 65k JumpReLU SAE (L0 $\approx$ 72), resynthesised activations from its codes, and broke one ingredient at a time. Every variant trained the same TopK SAEs as the real-data baseline: width 16k, k = 4–128, 3 epochs, 500k held-out tokens. About 72 SAEs on A100s.

γ in ranks 11–150 (mean of 2 SAE seeds for R, A, B, E and F; one seed for C and D):

Variant	What it keeps	Σ slope	k = 4	8	16	32	64	128	γ slope per doubling, k 8–128	Retained (γ)	Retained (raw)
R: real activations	everything	−0.89	0.81	0.70	0.67	0.57	0.46	0.36	−0.090	0.95	0.90
A: codes × decoder + bias	what the SAE captures	−0.94	0.78	0.70	0.67	0.54	0.44	0.34	−0.095	1.00	1.00
B: independent firing	per-latent frequencies and values, directions	−0.79	0.53	0.79	0.68	0.48	0.43	0.36	−0.111	1.17	0.98
C: decoder rows permuted	co-occurrence, directions; pairing broken	−0.75	0.67	0.87	0.59	0.50	0.42	0.36	−0.118	1.25	1.00
D: B and C together	frequencies, directions; unpaired	−0.78	0.55	0.85	0.66	0.52	0.45	0.35	−0.122	1.28	1.07
E: random directions	frequencies and values only	−0.84	0.13	0.23	0.08	0.07	0.07	0.06	−0.036	0.38	0.34
F: uniform frequencies, pooled values	directions only	−0.47	1.02	0.92	0.82	0.75	0.68	0.59	−0.080	0.84	0.42

Retained is a variant's per-doubling slope divided by A's, in two units: γ, and the raw decoder slope (γ times the variant's own Σ slope). Above 0.75 counts as keeping the law, below 0.5 as removing it. The first run (A–D) was scored in γ units over all six k: B 0.68, C 0.95, D 0.76. Under independent firing the k = 4 point dips, so the k = 8–128 fit was made primary before the second run (E, F and a second B seed). That switch was made after seeing the first run. B's dip reproduces across SAE seeds (γ 0.527 and 0.524), so it is a property of the independently fired data, not optimisation noise.

The resynthesis is faithful. A matches real data to RMS 0.022 across k, although its codes leave 21.5% of the variance unexplained. The law survives losing that residual, so it lives in what the SAE captures.

Co-occurrence and the frequency–direction pairing are not needed. From k = 16 up, B, C and D overlap the real curve, and they keep the decline in both units (raw retained 0.98–1.07). C and D lose most of the decoder-level cluster excess, yet keep it.

The split was inconclusive. The rule fixed before the second run read the split in γ units: if E keeps and F removes, the firing statistics carry the law; if E removes and F keeps, the dictionary's directions do. In γ units E removes (0.38) and F keeps (0.84), with per-seed slopes within 0.003 of each other. But γ divides by each variant's own Σ slope, and uniform firing halves F's (−0.47 against −0.94). In raw decoder-slope units F retains only 0.42 and E 0.34, so both remove. F is also close to circular. With uniform firing, F's Σ is essentially the Gemma Scope dictionary's own covariance (its slope, −0.469, matches the decoder cloud's −0.4684 in the reproduction), so at low k the new SAE relearns a subsample of that dictionary and γ starts near 1.

Caveats. E's γ of 0.06–0.08 at k $\geq$ 16 is what an exactly isotropic cloud measures at these sizes (0.06 at 16,384 points, 0.10 at 4,786), so on random isotropic directions the decoders show no anisotropy at all: E's "removes" is a floor, not a lost slope. In ranks 4–50 and 31–450, E is partial in γ units (0.62 and 0.67). There is one data draw per variant, and only layer 12 at width 16k.

Matched-covariance control. To separate Gemma Scope's specific directions from their second-order geometry, we drew unit directions from a Gaussian with the mean and covariance of Gemma Scope's decoder rows. We resynthesised with them under Gemma Scope's firing statistics (G, compared with D) and under uniform firing (H, compared with F), retraining D and F alongside. Each SAE was scored on 8,192 randomly chosen live latents (mean of 4 draws), which fixes the live count across k and variants, in both units. The rule, fixed before the run: a variant keeps the decline if it retains at least 0.75 in both units and removes it below 0.5 in both; if H keeps, F's $\gamma(k)$ needs only the dictionary's covariance.

γ in ranks 11–150 at 8,192 live latents (seed 0):

Variant	Σ slope	k = 8	16	32	64	128	Retained (γ)	Retained (raw)
D: Gemma Scope firing, Gemma Scope rows permuted	−0.780	0.838	0.637	0.487	0.425	0.327	—	—
G: Gemma Scope firing, Gaussian-matched directions	−0.770	0.814	0.545	0.436	0.453	0.392	0.76	0.75
F: uniform firing, Gemma Scope rows	−0.469	0.849	0.756	0.683	0.614	0.531	—	—
H: uniform firing, Gaussian-matched directions	−0.460	1.226	1.202	1.069	0.911	0.781	1.52	1.49

By the rule both keep, but G's pass is fragile. D and F reproduce their earlier γ exactly. G retains 0.76 in γ units and 0.7501 in raw units over k = 8–128, at the cut, on one seed and one draw of directions. Its curve is nearly flat after k = 16 (0.545, 0.436, 0.453, 0.392), so its slope comes from the drop between k = 8 and 16; refitted over k = 16–128 it retains 0.44 of D's decline, which would count as removing it. The supported reading is that a dictionary matching only the covariance of Gemma Scope's directions gives about half to three-quarters of D's decline under Gemma Scope's firing statistics. H adds no independent evidence. Its γ starts above 1 (1.23 at k = 8) and falls like the Gaussian null $N(\mu, \Sigma)$ on real activations, and its SAEs recover little (FVU 0.67 at k = 8, against 0.50 for F), because 72 random directions firing together make near-Gaussian data. Together with the Gaussian null, this says that much of γ's decline with k in ranks 11–150 needs only Σ. How much of the rest depends on Gemma Scope's specific directions is not settled by one seed at one layer and width. The earlier reading, "directions, not frequencies", stays withdrawn.

12 Summary of hypotheses

Hypothesis	Verdict	Deciding evidence
Li et al.'s finding: power-law spectrum, steepest mid-stack	Real, revised	Reproduced to 0.02%. The exponent depends on L0, layer and rank window; the mid-stack peak is in the top ∼150 directions only

Hypothesis	**Verdict**	**Deciding evidence**
The top-window spectrum is inherited from Σ	Largely supported in ranks 11–150; not in ranks 4–50	At layer 13 a Σ-matched Gaussian null gives γ within 0.03 of real at $k \geq 40$, and a decline over $k = 8$–128 of -0.088 against -0.097 per doubling. A covariance-matched dictionary gives about half to three-quarters of D's decline. In ranks 4–50 the null is flat in k and far lower
Pure estimator artifact	Not supported, within limits	Real data differs from Σ-matched Gaussian and scrambled nulls in the bulk windows (ranks 31–1000: γ 0.29–0.81 against 1.1–2.7), in ranks 4–50, at $k \leq 16$, and at layers 3 and 23. Width stability holds for Pythia, not for our Gemma runs
Frequency-driven latent allocation	Not supported on Gemma; consistent with planted circles at $k = 1$, but not diagnostic	On Gemma, latent frequencies (not the theory's feature frequencies) stay at -0.6 to -0.8 while the slope moves from -0.24 to -0.75, and per decoder cluster mean latent frequency falls with cluster size at coarse scales (slope -0.6 to -1.5), the opposite of the prediction. On planted circles at $k = 1$, $c = 0.334$ over all covered features (predicted 1/3) and $\gamma = 0.32$, but point features (predicted $c \approx 0$) give the same γ (0.30)
Gemma Scope's specific directions carry the $\gamma(k)$ decline	Withdrawn; partly open	Uniform-frequency resynthesis keeps the decline in γ units (0.84) but not in raw units (0.42), and its Σ is the dictionary's own covariance. Covariance-matched directions retain 0.44–0.76 of D's decline depending on the k range (one seed, one draw)
Feature splitting (near-duplicate clusters)	Ruled out	Removing near-duplicates, up to a third of latents, changes the slope by ≤ 0.01 in 22 SAEs
Massive-activation tokens drive the mid-stack peak	Ruled out	Dropping the top 0.1% or 1% of tokens on dim 1170 changes γ by ≤ 0.005

Hypothesis	Verdict	Deciding evidence
Decoder covariance $\propto \Sigma^{\gamma}$ as a matrix	Approximate, in the bulk; in the top directions only at low L0	In Σ's eigenbasis the decoder covariance follows Σ^{γ} over ranks 101–1000 at every L0 (R^2 0.94–0.99; a random basis gives about 0), but its diagonal share is 0.62–0.74 against 0.89–0.95 for an exact Σ^{γ} cloud of the same size
γ falls about 0.1 per doubling of L0	Supported in ranks 11–150	$1.245 - 0.114 \cdot \log_2$ L0 across 28 Gemma Scope SAEs ($1.207 - 0.110$ at a fixed 16,384 rows); $1.016 - 0.094 \cdot \log_2$ k for our TopK SAEs. Fails the window rule in the deeper windows
Zero-parameter law $\gamma = 1 - \ln k / \ln d$	Not supported	Fits at $d = 2304$, but with the same activations projected to $d' = 256$ the gap between $k = 8$ and $k = 32$ is 0.01, against 0.25 predicted
Mismatched L0 drives the mid-stack peak	Ruled out	Measured L0 runs up to 11% above the SAE names mid-stack; correcting for it raises the peak slightly ($0.19 \to 0.21$)
Mid-stack peak in real γ	Holds for Gemma Scope and our 16k SAEs; fails the window rule at 65k	Layer 13 tops real γ in 3 of 4 windows at 16k and 2 of 4 at 65k; the real-minus-Gaussian gap peaks at layer 13 in the top windows at both widths

Because γ is defined as the ratio of the two slopes, Li et al.'s -0.47 at layer 12 equals Σ's slope there (-0.87) times $\gamma \approx 0.54$ by construction. The matrix test is the stronger form of the claim, and it holds only approximately (see its limits in the mechanism section).

13 Open questions and next steps

The Σ^{γ} law is measured but not derived, and the matrix-test benchmark shows it is only approximate. Allocation fits planted circles at $k = 1$ but does not distinguish them from points, and it fails per cluster on Gemma. Much of γ's decline with k in ranks 11–150 needs only Σ, so it is not by itself a signature of structure; the clearest real-versus-null differences are in ranks 4–50, at small k and in the bulk windows. The mid-stack top-window effect goes with cluster structure beyond anisotropy, but only as a correlation across layers, and planting balanced clusters does not reproduce Gemma's $\gamma(k)$.

1. **What separates real from null γ, if not the k-slope?** Compare real and covariance-matched dictionaries in ranks 4–50, at small k and in the bulk, at a fixed live count, across layers, widths and L0s, with several seeds and draws of directions. Rescoring our real-data k sweep at a fixed live count needs a GPU retrain, since its decoders were not kept.

2. **Do the bulk-window results survive a better Σ?** At layer 12 the 200k-token Σ is too steep in the bulk by 0.08–0.12. Recompute Σ from at least 500k tokens at every layer, then redo the 26-layer profile and the Gemma Scope bulk γ.

3. **What is off-diagonal in the decoder covariance?** In Σ's eigenbasis it puts only 0.62–0.74 of its weight on the diagonal, against 0.89–0.95 for an exact Σ^{γ} cloud. Characterise the off-diagonal part, for instance against the cluster structure.

4. **Are our 65k runs undertrained?** Real γ drifts with width in our Gemma runs but barely in Gemma Scope's. Retrain at 65k with more tokens before reading width effects or the 65k mid-stack result.

5. **Does allocation scale with manifold dimension?** The same tiling argument predicts $c = m/(m+2)$ for m-dimensional manifold features: 1/3 for circles, 1/2 for spheres. A test needs a statistic that separates tiling from point features, which γ does not, plus more seeds and a width ladder.

6. **What makes the k-law depend on dimension?** The k-dependence vanished under PCA truncation but not in $d = 256$ planted data. Compare natural activations from models of different width, or planted data swept over d at fixed sparsity.

7. **Why does the mid-stack effect depend on the recipe** (JumpReLU 0.65 vs TopK 0.51 at layer 13)? Needs a few GPU-hours of JumpReLU training at matched L0.

8. **Can γ(L0) be derived?** A theory should give $\gamma = 1$ at $L0 = 1$, the decline with L0, and the Gaussian's steeper decline at small k.

9. **A systematic literature search** for prior results relating SAE or dictionary-learning spectra to the data covariance; the related-work section here is not one.

14 Methods, code and compute

Item	Detail
Models	Gemma-2-2B residual stream (`resid_post`); Pythia-70m layer 3
Pretrained SAEs	Gemma Scope `gemma-scope-2b-pt-res`, 16k to 1m wide: decoders of 288 SAEs; encoders of all 135 16k SAEs (L0 at every layer) and of the layer-12 SAEs used for firing frequencies
Data	Pile (uncopyrighted subset), BOS position dropped; Gemma context 1024, Pythia 256
Our SAEs	TopK with AuxK (k_{aux} 512), unit-norm decoder, Adam, linear LR decay over the last 20%; inputs scaled so $\mathrm{E}\|x\|^2 = d$
Spectrum	`geometry.eigen_slope`: centred covariance of decoder rows, ddof 1, least-squares log–log fit; the same function in every experiment
γ	Decoder slope divided by the slope of the training data's covariance, per window
Windows	Ranks 4–50, 11–150 (primary), 31–450, 101–1000 for Gemma; 3–25, 6–50 (primary), 11–100 for Pythia and the $d = 256$ planted data
Planted data	synth.py: sparse sums of point or circle features with $p_i \propto i^{-1.5}$, $d = 256$, noise 0.05. Dense: 4,096 features, 32 active per token. Sparse: 512 features, 4 active, random directions. Latents matched to features at cosine > 0.7

Item	Detail
Local compute	Apple M2 Pro, 16 GB, MPS
GPU compute	One H100 80 GB (Runpod Secure Cloud) for 88 runs, $50.06. A100 pods for the d = 2,304 planted, clustered, resynthesis, split and control runs, about $47 including idle and aborted pods. About $97 billed in total

All code, per-run JSON results and a dated research log are in the project repository (`notes/LOG.md`). Every result records its configuration and git commit. Figures show single values without error bars; seed spreads are given in the text where they were measured.

15 Sources

- Li, Michaud et al. (2024), The Geometry of Concepts: Sparse Autoencoder Feature Structure; code: ejmichaud/feature-geometry
- Michaud, Gorton, McGrath (2025), arXiv:2509.02565: latent allocation model of SAE scaling
- Peter, Søgaard (2025), Empirical Evaluation of Progressive Coding for Sparse Autoencoders, arXiv:2505.00190: latent frequency exponents cited for comparison
- Lieberum et al. (2024), Gemma Scope: the pretrained SAEs analysed here
- Gao et al. (2024), Scaling and evaluating sparse autoencoders: the TopK SAE and AuxK loss used for our own SAEs

A Planted-data sweeps

Interpolation: density sets γ's level, not its k-scale. Point features, random directions, d = 256, with features per dimension and active features per token raised together. Each row is a fit $\gamma = A + s \cdot \log_2 k$ over k = 4–64 (TopK, width 4,096, one seed).

Features, active per token	Planted: A	Planted: s	Gaussian twin: A	Gaussian twin: s	−s/A
512, 4	0.14	−0.015	1.20	−0.178	0.11*, 0.15
1,024, 8	0.55	−0.078	1.17	−0.175	0.14, 0.15
2,048, 16	1.17	−0.176	1.39	−0.210	0.15, 0.15
4,096, 32	2.11	−0.314	1.60	−0.224	0.15, 0.14
4,096, 32, crowded directions (dense regime)	1.30	−0.184	1.16	−0.166	0.14, 0.14
Gemma-2-2B layer 13, TopK	1.02	−0.094			0.09

* Poor fit (R^2 0.27): γ is near its floor at every k.

- **Density moves planted γ steadily from the allocation regime to Gaussian-like and past it.** The intercept rises 0.14 → 0.55 → 1.17 → 2.11. Planted data reaches its Gaussian twin near 2,048 features and 16 active, and overshoots it at 4,096 and 32 (2.11 vs 1.60). With

crowded directions at that density, planted γ falls back to near its twin, so crowding, not density, is what pins γ to the Gaussian value.

- **Every planted fit lies on one ray: $s \approx -0.15 \cdot A$.** Equivalently $\gamma(k) \approx A \cdot (1 - \log_2 k / \log_2 k_0)$ with $k_0 \approx 2^{6.7} \approx 100$ for every density, both direction settings, planted and Gaussian alike. Density changes the amplitude A only, never the k at which γ would reach zero.
- **The path misses Gemma.** Gemma's $-s/A = 0.09$ puts its k_0 near $2^{10.8} \approx 1{,}800$, off the planted ray. The blend hypothesis is half supported: density does carry γ from the allocation value to Gaussian-like, but Gemma's slow decline with k needs a separate scale that this generator does not vary. Candidates are d, SAE width and the training recipe; d alone is doubtful, since Gemma truncated to 256 PCA dimensions had an almost flat k-dependence rather than the planted one.

Width moves k_0, but only slightly. At the density nearest Gemma's intercept (2,048 features, 16 active), we swept SAE width from 1,024 to 16,384. Planted $\log_2 k_0$ rises about 0.25 per doubling of width, from 6.37 at 2,048 to 7.11 at 16,384, against Gemma's 10.8 at the same width. The Gaussian twin follows up to 8,192 and then levels off at 6.8. Width tilts $\gamma(k)$ rather than shifting it: wider SAEs lower γ at small k ($k = 4$: $0.97 \to 0.67$) and leave it almost unchanged at $k \geq 16$. So the intercept and the slope shrink together. At width 8,192 the intercept matches Gemma's (1.02) but the slope is -0.148; at 16,384 the slope is closer (-0.122) but the intercept has fallen to 0.87. Most of Gemma's slow decline with k must come from something else: d, the data, or the recipe. One caveat: Gemma's fit uses ranks 11–150 and k = 1–128 at d = 2,304, so k_0 is not strictly comparable across the two.

Dimension sets the amplitude; k_0 rises only slowly. We then held density fixed as ratios to d (8d features, d/16 active) and swept d from 64 to 1,024, scoring γ in a rank window that scales with d (ranks d/43 to d/5). The amplitude A rises steadily with d, from 0.44 at d = 64 to 2.08 at d = 1,024. $\log_2 k_0$ stays at 6.5–7.0 up to d = 512 and reaches 8.05 only at d = 1,024, about +0.24 to +0.35 per doubling, driven mostly by that last point. So k_0 is set in absolute k, about 100–250 latents per token, not as a fraction of d.

Putting the sweeps together: across every planted setting, $\gamma(k) \approx A \cdot (1 - \log_2 k / \log_2 k_0)$. Density and d set A. Width and d move k_0 slowly. Extrapolated to Gemma's d = 2,304 and width 16k, the planted k_0 reaches about $2^{8.5}$. Gemma's own k_0 depends on the window and the k range: $2^{8.9–9.2}$ in ranks 4–50, $2^{10.8}$ in ranks 11–150, and $2^{9.4–12.1}$ in ranks 31–450. Dimension and width therefore account for part of Gemma's slow decline with k, possibly most of it in the top window, but not robustly in the bulk. Gemma's intercept (1.02) is below the planted A at $d \geq 512$, so a Gemma-like intercept in this family needs lower density, which moves A without moving k_0. One seed, one data family, and the d trend rests mostly on d = 1,024.

At Gemma's own d and width, k_0 reaches Gemma's range. We trained 60 SAEs on an A100 at d = 2,304 and width 16k. That covered three densities (2d features with d/64 active, 4d with d/32, 8d with d/16), k = 4–128, real data and Gaussian twins, and three seeds at one density. The top rank windows turned out to be unusable for this generator. Feature probabilities are capped at 0.5, and at this d the cap saturates the 25–101 most frequent features. That gives Σ a plateau over its top ranks, with slope -0.1 to -0.4 in ranks 4–50 against Gemma's -0.98, so γ there is a ratio over a near-zero denominator. In the d-scaled window (ranks 55–461), where Σ falls with slope -1.0 to -1.3, the planted $\log_2 k_0$ is 9.2 at the sparsest density and 15.3 at the densest. Gemma's is 13.0 in its nearest window (ranks 31–450, k = 4–128). The densest planted curve runs

parallel to Gemma's, at -0.093 vs -0.083 per doubling, about 0.2–0.35 higher. So the k-scale gap seen at $d = 256$ is mostly a d-and-width effect. Two qualifications remain. The planted curves are less clean than Gemma's law (one density is flat until $k = 64$), partly because many latents die at small k with 6M training tokens. And the comparison in ranks 11–150, where Gemma's law is cleanest, needs a generator whose Σ is a power law from rank 1. Seed spread is ≤ 0.02 in almost every cell.

With Σ matched to Gemma's, the ranks 11–150 law is not reproduced. We reran the three densities with feature magnitudes set so that feature variance falls as i^{-1}. That removes the plateau and gives Σ slopes of -0.95, -0.90, -0.82 and -0.80 in ranks 4–50, 11–150, 31–450 and 55–461 (Gemma -0.98, -0.95, -0.75, -0.71). We used 12M tokens and 3 epochs, 30 SAEs in all (about 3.6 A100-hours). In ranks 11–150, real planted γ is flat in k at every density: 0.1–0.3 at the two sparser densities and 0.36–0.46 at the densest (2 seeds, sd ≤ 0.02). Gemma falls from 0.82 at $k = 4$ to 0.35 at $k = 128$. The best planted RMS against Gemma is 0.27. Only the densest setting in the wide windows looks like Gemma: in ranks 31–450 it falls from 0.88 at $k = 8$ to 0.38 at $k = 128$ (fit over $k \geq 8$: $A = 1.28$, slope -0.124 per doubling, R^2 0.98, $\log_2 k_0 = 10.3$, against Gemma's 1.08, -0.083, 13.0). Its $k = 4$ point (0.44) breaks the trend. The Gaussian twin, by contrast, now has the same shape as Gemma's Gaussian twin ($2.90 \to 0.73$ in ranks 31–450 vs Gemma's $2.20 \to 0.70$). So with Σ a power law, the null carries over to planted data and the real-data curve does not. At the sparser densities planted decoders carry only a fraction of Σ's anisotropy ($\gamma \approx 0.1$–0.3 in every window). That is what one expects when an SAE recovers random feature directions, whatever their variances. Gemma's γ of 0.35–1.0 in the top window needs decoders that inherit much more of Σ, which points to structure the planted family lacks, such as correlated or clustered feature directions. Dead latents remain a confound: at $k \leq 8$ only 1,900–9,100 of 16,384 latents stay alive.

Planting cluster structure raises γ but does not give it Gemma's k-dependence. We added clusters to the power-law generator: features in groups of 16, each at cosine $\rho = 0.4$ or 0.6 to a random centre. This leaves Σ's slopes within 0.03 of the unclustered version. We trained 30 SAEs at the two denser settings ($k = 4$–128; 2 seeds on one setting; about 4 A100-hours). The test, fixed before the run, was RMS against Gemma below 0.15 in ranks 11–150 plus a slope within 0.04 of Gemma's -0.094 per doubling. Every setting fails. The best RMS is 0.24, and every slope lies between -0.02 and $+0.03$ per doubling, in every window. Clusters move the level, not the shape. At $\rho = 0.6$, mean γ in ranks 11–150 rises from 0.16 to 0.26 at the sparser setting (4d features) and from 0.41 to 0.48 at the denser one (8d); in ranks 31–450 at 4d it rises from 0.22 to 0.44. The sparser setting at $\rho = 0.4$ is the telling case. Its trained decoders carry Gemma's level of cluster structure beyond their covariance (excess between-cluster share $+0.04$ to $+0.08$, against Gemma's $+0.054$), yet its γ stays near 0.19 and flat. At the denser setting, which has more features than latents, the decoders lose the planted clusters (excess ≈ 0 for $k \geq 16$). So balanced, frequency-independent clusters do not produce Gemma's $\gamma(k)$.