

The complete chloroplast genome sequence of *Populus simonii*, a medicinal plant for anti-lung cancer activity

Yi Zhuo*, Jian-Bo Lin*, Yin-He Yin and Fan-Cai Lai

Department of Thoracic Surgery, First Affiliated Hospital of Fujian Medical University, Fuzhou, Fujian, China

ABSTRACT

Lung cancer is one of the most common malignant tumors. It is clinically divided into two types: small cell lung cancer and non-small cell lung cancer. *Populus simonii* distributed in East Asia region including China used as traditional medicine, which is an important medicinal plant for anti-lung cancer activity. The complete chloroplast genome sequence of *P. simonii* was characterized from Illumina pair-end sequencing. The chloroplast genome of *P. simonii* was 156,559 bp in length, containing a large single-copy region (LSC) of 84,825 bp, a small single-copy region (SSC) of 47,561 bp, and two inverted repeat (IR) regions of 16,494 bp. The overall GC content is 36.70%, while the corresponding values of the LSC, SSC, and IR regions are 34.5%, 30.0%, and 42.0%, respectively. The genome contains 131 complete genes, including 86 protein-coding genes (68 protein-coding gene species), 37 tRNA genes (29 tRNA species), and 8 rRNA genes (4 rRNA species). The neighbour-joining phylogenetic analysis showed that *P. simonii* and *P. qamdoensis* clustered together as sisters to other *Populus* species.

ARTICLE HISTORY

Received 16 June 2020
Accepted 20 June 2020

KEYWORDS



Populus simonii; chloroplast genome; phylogenetic analysis; anti-lung cancer activity

Introduction

Lung cancer is one of the most common malignant tumors. The latter is the most common type of lung cancer pathology. Lung cancer occupies the number one cause of cancer death in the world. Although there have been some advances in the diagnosis and treatment of lung cancer, there are currently no specific treatment drugs. *Populus simonii* distributed in East Asia region including China used as traditional medicine, which is an important medicinal plant for anti-lung cancer activity, which has persisted largely in an undomesticated state that is highly resistant to different environmental stresses. Most patients with non-small cell lung cancer are not sensitive to radiotherapy and chemotherapy, so it is urgent to develop anti-cancer drugs with high efficiency and low toxicity and side effects based on the research of tumor molecular signaling pathways. Natural products are an important source of anti-tumor chemical drugs. Lung cancer refers to a disease caused by environmental and genetic factors, with dementia as the main clinical phase. Modern pharmacological studies have shown that *P. simonii* can significantly increase the number of capillary networks and accelerate blood flow, thereby restoring the function of microcirculation. At the same time, it can also reduce plasma lactic acid content and improve metabolic disorders caused by cell hypoxia. It plays an important role for anti-lung cancer activity. Studies have shown that *P. simonii* can reduce the content of lipid peroxide and erythrocyte sorbitol, increase the level of superoxide dismutase, reduce the

oxidative stress reaction, and has an antioxidant effect. Studies have also found that *P. simonii* can improve the role of lung cancer. The mechanism may be: the antioxidant effect of drugs, by restoring the structure and function of nerve cells, inhibiting lipid peroxidation, improving blood rheology, and protecting against ischemia. Damaged neurons improve the pathological response of for lung cancer. Since it contains selaginellin, flavonoids and sumafavone, it also presented anti-cancer, inhibition of immediate allergic reactions, and antihyperglycemic activities. *P. simonii* has high ecological and economic value with high levels of intraspecific genetic diversity. *P. simonii* has wide geographic distribution, high intraspecific polymorphism, adaptability to different environments, combined with a relatively small genome size. Consequently, *P. simonii* represents an excellent model for understanding how different evolutionary forces have sculpted the variation patterns in the genome during the process of population differentiation and ecological speciation (Neale and Antoine 2011). Moreover, we can develop conservation strategies easily when we understand the genetic information of *P. simonii*. In the present research, we constructed the whole chloroplast genome of *P. simonii* and understood many genome variation information about the species, which will provide beneficial help for population genetics studies of *P. simonii*.

The fresh leaves of *P. simonii* were collected from Yanzhou (116°35'E; 35°43'N). Fresh leaves were silica-dried and taken to the laboratory until DNA extraction. The voucher specimen

CONTACT Fan-Cai Lai  laifancaimd@163.com  Department of Thoracic Surgery, First Affiliated Hospital of Fujian Medical University, Fuzhou, Fujian, China
*These authors contributed equally to this work.

References

- Campos FS, Kluge M, Franco AC, Giongo A, Valdez FP, Saddi TM, Brito WMED, Roehe PM. 2016. Complete genome sequence of porcine parvovirus 2 recovered from swine sera. *Genome Announc.* 4(1): e01627.
- Doyle J. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull.* 19(1):11–15.
- Hahn C, Bachmann L, Chevreux B. 2013. Reconstructing mitochondrial genomes directly from genomic next-generation sequencing reads—a baiting and iterative mapping approach. *Nucleic Acids Res.* 41(13): e129–e129.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30(4):772–780.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol.* 33(7): 1870–1874.
- Neale DB, Antoine K. 2011. Forest tree genomics: growing resources and applications. *Nat Rev Genet.* 12(2):111–122.
- Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile metagenomic assembler. *Genome Res.* 27(5):824–834.
- Wyman SK, Jansen RK, Boore JL. 2004. Automatic annotation of organellar genomes with DOGMA. *Bioinformatics.* 20(17):3252–3255.