



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Polyphenols as promising biologically active substances for preventing SARS-CoV-2:
A review with research evidence and underlying mechanisms

Taha Mehany, Ibrahim Khalifa, Hassan Barakat, Sami A. Althwab, Yousef M. Alharbi,
Sobhy El-Sohaimy



PII: S2212-4292(21)00016-X

DOI: <https://doi.org/10.1016/j.fbio.2021.100891>

Reference: FBIO 100891

To appear in: *Food Bioscience*

Received Date: 17 June 2020

Revised Date: 12 January 2021

Accepted Date: 12 January 2021

Please cite this article as: Mehany T., Khalifa I., Barakat H., Althwab S.A., Alharbi Y.M. & El-Sohaimy S., Polyphenols as promising biologically active substances for preventing SARS-CoV-2: A review with research evidence and underlying mechanisms, *Food Bioscience*, <https://doi.org/10.1016/j.fbio.2021.100891>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Elsevier Ltd. All rights reserved.

TH, IK and HB, Conceptualization, Methodology, and Writing-Original draft. HB, SA, and SE Visualization, Reviewing, and Editing. YA, Language editing and proofing.

Polyphenols as promising biologically active substances for preventing SARS-CoV-2: A review with research evidence and underlying mechanisms

Running title: The potential role of polyphenols against SARS-CoV-2

**Taha Mehany^{a#}, Ibrahim Khalifa^{b#}, Hassan Barakat^{b,c*}, Sami A. Althwab^c,
Yousef M. Alharbi^d, Sobhy El-Sohaimy^{a,e}**

^a Food Technology Department, Arid Lands Cultivation Research Institute, City of Scientific Research and Technological Applications, 21934, Alexandria, Egypt.

^b Food Technology Department, Faculty of Agriculture, Benha University, Moshtohor, 13736, Egypt.

^c Department of Food Science and Human Nutrition, College of Agriculture and Veterinary Medicine, Qassim University, Buraydah 51452, Saudi Arabia.

^d Department of Veterinary Medicine, College of Agriculture and Veterinary Medicine, Qassim University, Buraydah 51452, Saudi Arabia.

^e Department of Technology and Organization of Public Catering, Institute of Sport, Tourism and Service, South Ural State University 454080, Chelyabinsk, Russia.

* Correspondence should be addressed to Dr. Hassan Barakat

Email: hassan.barakat@fagr.bu.edu.eg

haa.mohamed@qu.edu.sa

Authors are equally contributed.

Abstract

Currently, antiviral drugs and/or vaccines are not yet available to treat or prevent severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). In this review, we narrated the available data, from credible publishers, regarding the possible role of polyphenols and natural extracts-containing polyphenols in the prevention of coronavirus disease 2019 (COVID-19), and their immune-boosting properties. It was revealed that polyphenols could be considered as promising biologically active substances for the prevention of COVID-19. The underlying potential mechanism behind this action is mostly due to the antiviral activities and the immune-regulation functions of polyphenols against COVID-19-infections. Antivirus polyphenolic-based medications can mitigate SARS-CoV-2-enzymes, which are vital for virus duplication and infection. It was also found that triterpenoid, anthraquinone, flavonoids, and tannins are possible keys to scheming antiviral therapies for inhibiting SARS-CoV-2-proteases. The identified pharmacophore structures of polyphenols could be utilized in the explanation of novel anti-COVID-19 designs. The advantage of using mixtures containing polyphenols is related to the high-safety profile without having major side-effects, but further randomized controlled trials are required in the upcoming studies.

Keywords: SARS-CoV2; Polyphenols; Functional foods; Boosting immune functions.

Abbreviations: Severe acute respiratory syndrome coronavirus 2: SARS-CoV-2; Coronavirus disease 2019: COVID-19; Coronaviruses: CoVs; Severe observation of acute respiratory syndrome: SARS-CoV; Middle East Respiratory Syndrome: MERS-CoV.

1. Introduction

SARS-CoV-2 affects all peoples, especially those who had weak immune systems and/or week immune-based responses. Coronaviruses (CoVs) belonged to the sub-family of Ortho-coronavirinae, under the Coronaviridae family and in order of Nidovirales, and this sub-family contains alpha-, beta-, gamma-, and delta-CoVs (Banerjee *et al.*, 2019). The deadliness of CoVs has been identified by the outbreaks of severe observation of acute respiratory syndrome (SARS-CoV) in 2002/2003 and the Middle East Respiratory Syndrome (MERS-CoV) in 2012/2013. SARS-CoV and MERS-CoV are going to β -CoV (Zhang *et al.*, 2020). Freshly, a new flu-like CoV (COVID-19) associated with SARS-CoV and MERS-CoV was realized in the late of 2019 and spread worldwide with a pandemic (Cohen and Normile, 2020). The total number of cases globally was 91,322,850 and 1,952,979 confirmed deaths in WHO situation report COVID-19 in January 12, 2021 (worldmeters.info).

Hitherto, no vaccine and actual SARS-CoV-2 medications based on anti-SARS-CoV therapy by the scientific community were confirmed (Jo *et al.*, 2020), while scientists are trying to develop efficient vaccines or drugs for the SARS-CoV-2 virus. The general viral infection treatment involving dietary interferences and other immune enhancers which were utilized to promote immunity towards the viral poisons. Thus, they may also be utilized to contest SARS-CoV-2 by the nutritional status of each patient and should be assessed before applying the overall treatments (Zhang *et al.*, 2020). Recently, numerous studies have exposed that phytonutrients play a vital part in the stoppage of chronic syndromes, as most can be related to nourishments (Fraga *et al.*, 2019). Various countries have entered the notion of functional-based food not only essential for human life but also to boost the physiological functions and immunity responses (Del Bo *et al.*, 2019; Quero *et al.*, 2020). The importance of the proper functioning of the body and the well-balanced immune system for the maintenance of human

life has been remarkably evident over the last decades (Lopez-Varela *et al.*, 2002; Wichers, 2009).

In addition to phytonutrients mainly presented in fruit and vegetables, polyphenols have recently been targeted due to their role in the prevention of several diseases as described (Fraga *et al.*, 2019). Interestingly, polyphenols have attracted much attention in recent years since their dietary consumption has been associated with the prevention of some chronic and degenerative diseases that are significant causes of death and incapacity in developing countries (Santos-Buelga *et al.*, 2019). In addition, due to the biological activities of polyphenols, particularly antimicrobial, antioxidant, anti-inflammatory, pro-apoptotic behavior, anti-proliferative, and hormonal control ability among polyphenols, countless plants have been considered and investigated because of the classification of new substances with nutraceutical possessions (Forni *et al.*, 2019). Dietary polyphenols signify several bioactive substances broadly found in plant-based food which have the capabilities to prevent the beginning and development of several diseases, and to decrease and control their signs which become a central issue in food research (Del Bo *et al.*, 2019; Quero *et al.*, 2020).

The outbreak of the COVID-19 pandemic caused by SARS-CoV-2, along with the lack of targeted medicaments and vaccines, forced the scientific world to seek new antiviral formulations. This article reviews the feasibility of using polyphenols extracted from plants as a source of bioactive components (e.g., gallates and quercetin), effective in mitigating SARS-CoV-2 and boosting the immune system. There is potential for the utilization of plant-derived polyphenols as functional foods and pharmaceutical formulations. The method of employing such formulations is fast because the raw materials for their preparation, namely herbs and useable plants, are permitted for human consumption worldwide. Approximately 100 research papers were selected from Science Direct, Scopus, MDPI, PubMed, and Web of

Science to review the recently published papers on the related topic with the aid of relative keywords.

2. Functional foods and their efficiency in boosting the immune response

In fact, malnutrition is the most common cause of immunodeficiency concerns worldwide (Schaible and Stefan, 2007). The immune system, which is joined into all physiological arrangements, defends the human body towards microorganisms' infections and other internal and external abuses (Castelo-Branco and Soveral, 2014; Maggini *et al.*, 2018). And suitable nutrition at all phases of life is an important factor to guarantee a sufficient stock of energy, macro- and micronutrients essential for the growth, upkeep, and countenance of the immune response (Maggini *et al.*, 2007). Furthermore, undernutrition is leading to weakening immune function which can be due to consumption of fast food, insufficient intakes of energy and macronutrients, and/or shortages in the essential micronutrients (Alpert, 2017; Calder, 2013; Pereira, 2003). **Fig S1.** showed the sources of variation in the immune system and its relation with suboptimal nutrient status on an increased risk of infection, as previously described (Calder and Kew, 2007; Maggini *et al.*, 2018), showing the vital role of the functional foods.

In this context, functional foods are used to strengthen other physiological functions including the immunity responses (Roberfroid, 2000). The characteristics of functional foods were tabulated in **Table 1**. Functional food components can be useful components that originate logically in or complement food as functional elements such as vitamins, minerals, dietary fiber, fatty acids, amino acids, polyphenols, phytopigments, non-saponifiable fractions, and functional plant-based proteins (Guiné *et al.*, 2009; London, 2010). **Table 2** summarises the functional foods-based products which were quoted from the available data. These foods and/or ingredients boost the immune response by different mechanisms (Powell *et al.*, 2000). Meanwhile, the deficiencies of one or more essential fatty and amino acids,

vitamins, Se, Zn, Fe, Cu, and phytonutrients will affect almost all types of immunity (London, 2010; Maggini *et al.*, 2018; Zhang *et al.*, 2020). Regarding COVID-19 pandemic, these functional ingredients may mitigate SARS-CoV-2 through their immunoenhancing and antiviral activity. For example, Zhang *et al.* (2020) had claimed that vitamin A and/or its derivatives like retinoic acid could provide a promising alternative for CoV-therapy. Vitamin D has several immune-modulatory effects and might work as additional therapeutic against COVID-19 (Moriguchi and Muraga, 2000; Zhang *et al.*, 2020). Zhang and Liu (2020) had stated that Zn may affect not only COVID-19-linked symptoms like inferior respiratory tract infection and diarrhea but also SARS-CoV-2 replication. Dosing the patients with Se could be an active choice for the cure of COVID-19 as previously stated by Zhang *et al.* (2020), for instance. These functional compounds are essential substances at the molecular and cellular levels that play critical functions.

3. The potential uses of plant extracts containing polyphenols against SARS-CoV-2

3.1. Overview of the health-benefits of polyphenols

Polyphenols constituents are categorized by the attendance of one or more hydroxy group on an aromatic circle. These components are categorized by their molecular mass, chemical construction, and intricacy to flavonoids (flavones, flavonols, flavanones, flavanonols, iso-flavonoids, flavanols, anthocyanidins, and chalcones) and non-flavonoid ones (phenolic acids, stilbenes, curcuminoids, lignans, and tannins) (Quero *et al.*, 2020). **Fig S2.** portrayed the plant polyphenolics classifications according to Phenol-Explorer (<http://www.phenol-explorer.eu>). Such polyphenols are valuable to human health and are capable to decrease the diseases perils (Rasouli *et al.*, 2017). Thus, polyphenols have recently attracted much attention to the scientific community due to their dietary intake which is related to the deterrence of some chronic and wasting diseases that establish the main reasons of death. The

unique properties of their nature and position of the substituents have concerned for biological utilization, namely, antiviral, antibacterial, antioxidant, and anti-inflammatory activities. These polyphenols have been isolated from a great number of plants, such as fruits, vegetables, legumes, nuts, seeds, herbs, spices, tea, coffee, dark chocolate, and so on (Quero *et al.*, 2020; Rasouli *et al.*, 2017; Santos-Buelga *et al.*, 2019; Silva and Pogačnik, 2020). Among the sources of polyphenols, citrus is a promising source due to its content from bioflavonoids including apigenin, acacetin, luteolin, diosmetin, chrysoeriol, and glycosides which play key roles in providing an extensive range of health useful effects. However, polyphenols' super-healthy properties depend on the amount ingested and their bioavailability (Silva and Pogačnik, 2020).

3.2. Polyphenols and immune system

Many factors cause immune disorders, including environmental stress and changes in the immune regulators that can lead to many human diseases. Bioactive components can play a vital role in immunity through intrusive with immune cell regulation and pro-inflammatory cytokine synthesis. Recently, a huge number of *in vitro* and *in vivo* studies demonstrated the immunomodulatory part of polyphenols. These studies have shown that people with a specific diet, especially polyphenols-rich ones, are at reduced risk for a variety of diseases (Rasouli *et al.*, 2017; Yahfoufi *et al.*, 2018). Additionally, polyphenols are developing therapeutic tools for controlling and protecting the body from several autoimmune disorders due to their immunomodulatory effects (Khalifa *et al.*, 2020; Khan *et al.*, 2019). The immune modulation effect of polyphenols is related to the impact on immune cells, pro-inflammatory gene expression, and modulate cytokines production (Yahfoufi *et al.*, 2018). In addition to the antioxidant properties such as reactive oxygen species scavenging of polyphenols and their donations to the rule of inflammatory signing. Polyphenols also showed high-effects in increasing the natural killers in fruits (Yahfoufi *et al.*, 2018). For their effect on the balance

between pro and anti-inflammatory cytokine secretion, polyphenols compounds such as catechins and quercetin have been added to various foods. Additionally, numerous flavonoids can inhibit the expression of various pro-inflammatory cytokines and chemokines (Comalada *et al.*, 2006). Furthermore, the epigallocatechin gallate and curcumin can induce epigenetic changes in cells. Thus, polyphenols compounds can be efficiently used to control the immune response.

3.3.The antiviral activities of polyphenols

The literature review demonstrated that polyphenols have been investigated for their potential antiviral efficiency against SARS-CoV-2 in molecular modeling studies, cell-free polyphenol–protein interaction studies, and in cell-based virus infection studies (Chojnacka *et al.*, 2020; Paraiso *et al.*, 2020). Convincing data shows that polyphenols such as epigallocatechin gallate, resveratrol, and curcumin are prime candidates for preclinical and clinical trials. As a cautionary note, the efficacy of tested polyphenols against human viral disease *in silico* and *in vitro* screening methods are not validated. Potential regulation of COVID-19 intensity by polyphenols controlling ACE2 *in vivo* has been suggested (Horne and Vohl, 2020; Manoharan *et al.*, 2020) but very few studies have been performed to examine the antiviral activity of polyphenols against SARS-CoV-2 *in vivo*. Pudilan Xiaoyan Oral Liquid (PDL), a traditional Chinese medicine containing four herbs and more than 180 ingredients, showed potent anti-SARSCoV-2 activity in infected hACE2 mice (Deng *et al.*, 2020). It has also been stated that a nebulized formula of quercetin and N-acetylcysteine substantially relieved the respiratory symptoms of SARS-CoV-2 in a patient treated with hydroxychloroquine and antibiotics (Schettig *et al.*, 2020). This reinforces the value of more clinical trials to determine the potential of polyphenol-based nutraceuticals as an adjuvant or key treatment for COVID-19. High-throughput screening methods will speed the *in vitro* development of lead candidates by reducing the number of polyphenol libraries. Screening of

polyphenol-rich plant extracts is an alternative, commonly used method, but it has the drawback that extracts contain a multitude of natural products with inherent difficulties of not being able to reliably distinguish active concepts and the potential for pharmacological antagonism. These problems can be solved by integrating classical bioassay-guided fractionation with machine learning approaches that expose the identity of bioactive natural products in extracts without the need for homogeneity purification. In our view, the above hybrid method aims to accelerate research because many antiviral *in vitro* experiments can be conducted without the treatment of live viruses and because the identification of polyphenols (and other natural products) has been made simpler over the past few years thanks to developments in plant metabolism and ever-growing natural product databases such as phenol-explorer (Rothwell *et al.*, 2016), knapsack (Shinbo *et al.*, 2006), and the global natural product social networking database (Wang *et al.*, 2016).

Interestingly, there are no known therapies for CoVs-infection and defensive vaccines are still being discovered even discovered still need extensive evaluation. Thus, the situation illustrates the need for appropriate antivirals and/or immune-enhancers to be produced for prophylaxis CoV-infection. Recently, the antiviral activity of numerous polyphenols from different sources was evaluated against several viruses (Khaerunnisa *et al.*, 2020; Maryam *et al.*, 2020). **Table 3** shows the antiviral activities of polyphenols, their main sources, and mode of action (Khaerunnisa, 2020). In a related study, Sundararajan *et al.* (2010) evaluated three pomegranate extracts including pomegranate juice, concentrated liquid extract, and dried extract for their anti-influenza activity. He found that the anti-influenza activity of pomegranate polyphenols is substantially modified by changes in covering glycoproteins. Likewise, Vela *et al.* (2010) who utilized isoflavone (genistein) and general kinase inhibitor for treating animals, and reported that general kinase inhibitor genistein may use as an anti-arenavirus component. It was also found that theaflavins, which are the polyphenols of black

tea, had antiviral effects on vaccinia virus, influenza virus, poliovirus-1, herpes simplex virus, coxsackievirus, and human rotavirus. Morán-Santibañez *et al.* (2018) reported that polyphenol-rich extracts isolated from Mexican seaweed have *in vitro* significant virucidal effects on the Measles virus. Virucidal activity of these extracts is not only a prophylactic tactic before viral infection but can also be fruitful as a dealing after contagion and for avoiding virus spreading.

3.4.The role of polyphenols against SARS-CoV, MERS-CoV, and SARS-CoV-2

Regarding SRARS-CoV, saikosaponins, naturally occurring triterpene glycosides derived from medicinal plants such as *Heteromorpha* spp., *Bupleurum* spp., and *Scrophularia scorodonia*, showed several biological activities, specifically antihepatoma, anti-hepatitis, antinephritis, anti-inflammation, immunomodulation, and antimicrobial effects. Thus, Cheng *et al.* (2006) recommended the use of saikosaponins as *in vitro* antiviral activity against CoVs-22E9. The extracts of *Cibotium barometz*, *Gentiana scabra*, *Dioscorea batatas*, *Cassia tora*, and *Taxillus chinensis* inhibited the SARS-CoV replication better than other 200 extracts (Wen *et al.*, 2011). Likewise, the extracts of *Gentianae radix*, *Dioscoreae rhizoma*, *Cassiae Semen*, *Loranthi Ramus*, and *Rhizoma Cibotii* in the concentrations from 25 to 200 $\mu\text{g mL}^{-1}$ inhibited SARS-CoV. Bioflavonoids from *Torreya nucifera* inhibited SARS-CoV 3CLpro by 62% at 100 $\mu\text{g mL}^{-1}$ (Ryu, Jeong, *et al.*, 2010), where he identified 8 diterpenoids and 4 bioflavonoids by fluorescence resonance energy transfer analysis. The activity of flavones (apigenin, luteolin, and quercetin) was also investigated with IC_{50} -values of 280.8, 20.2, and 23.8 μM , separately, and the presence of the apigenin moiety at the C-30 position of flavones seems to be more effective. Ethyl acetate fraction of *Houttuynia cordata* Thunb. contribute to the superior antiviral efficacy *in vitro* and *in vivo* presenting the potential for the development of antiviral agents against coronavirus and dengue infections (Chio *et al.*, 2016) with an IC_{50} -value of 0.98 mg mL^{-1} . The IC_{50} -values for quercetin, epigallocatechin

gallate, and galusatechin gallate against SARS-CoV were 73, 73, and 47 μM , respectively, and the galooyl group should be present at 3-OH-position to be more efficient (Nguyen *et al.*, 2012). Quercetin derivatives such as 7-O-arylmethylquercetin and quercetin-3- β -galactoside could combat as anti-SARS-CoV agent (Chen *et al.*, 2006; Park *et al.*, 2012). Flavonoids found in *Galla chinensis* or *Veronica lin rrifolia* extracts may interact with the surface of the spiky protein of the SARS-CoV, avoiding virus particles from powerful the cell (Wang and Liu, 2014). Phlorotannins isolated from brown algae display an inhibitory effect of SARS-CoV-3CLpro, regulating the virus replication. In particular, dieckol showed high inhibitory activity with an IC_{50} value of 2.7 μM (Park *et al.*, 2013). Tetra-O-galoilo- β -d-glucose and luteolin have been confirmed to be active against SARS-CoV (Lin *et al.*, 2005). In particular, *Lycoris radiata* extract showed to be a good entrant against SARS-CoV as it showed the highest efficacy, mainly due to the lycorin component (Lin *et al.*, 2005). In addition, Park *et al.* (2017) isolated 10 polyphenolics from *Broussonetia papyrifera* as CoVs-protease inhibitors against 3-chymotrypsin-like (3CLpro) and-papain-like-CoVs-cysteine-proteases (PLpro). It was also found that all polyphenols have more effectiveness against PLpro than 3CLpro; the results suggested polyphenols of *B. papyrifera* have anti-CoVs activity. Likewise, Ryu *et al.* (2010) examined the capacity for SARS-CoV inhibition of flavonoids from the *Torreya nucifera* leaves. The ethanolic extract of *T. nucifera* leaves-inhibited the SARS-CoV-3CLprotease with inhibition values of 62% at 100 g mL^{-1} . Additionally, this study confirmed that amentoflavone is more effective than the respective flavones (luteolin and apigenin) and bioflavonoids derivatives on SARS-CoV-3CLpro. Thus, amentoflavone may be a good component for development as a natural therapeutic medicine against SARS-CoV infection. Jo *et al.* (2020) studied the antiviral activity of three flavonoids against SARS-CoV. Rhoifolin, herbacetin, and pectolinarin have been found to block SARS-CoV-3CLpro's enzyme activity. The interaction of these bioflavonoids was established using a

tryptophan-based fluorescence method, and the three flavonoids were recommended to be templates to strategy functionally developed inhibitors. **Fig 1A.** presented the binding pose of amentoflavone in SARS-CoV 3CLpro cited from the study of Ryu *et al.* (2010). As for MERS-CoV, flavonoids were considered as potential inhibitors of MERS-CoV (Jo *et al.*, 2019), where each of herbacetin, isobavachalcone, quercetin 3- β -d-glucoside, and helichristetine blocked the enzymatic activity of MERS-CoV-3CLpro. It was noted that flavonoid derivatives with hydrophobic or carbohydrate groups devoted to their core structures inhibit MERS-CoV. The computational and experimental study showed that chalcone and flavonol are desired supports to bind with the catalytic site of MERS-CoV-3CLpro. Therefore, this study suggested that flavonoids with these properties can be used as models to develop powerful MERS-CoV-3CLpro inhibitors. Regarding SARS-CoV-2, Negida, and Bahbah (2020) reported that saikosaponins are considered as active substances for SARS-CoV-2 due to their immunomodulatory, anti-inflammatory, and antiviral activities. The water extract from *Isatis indigotica* root, containing a number of phenolic compounds, namely sinigrin (IC₅₀: 217 μ M) and hesperetin (IC₅₀: 8.3 μ M) showed anti-SARS-CoV-2-3CLpro. Through sequential matching, it has been proven that SARS-CoV-2 major protease has approximately a 96% similarity to SARS-CoV-2-Mpro (Xu *et al.*, 2020). The inhibition of Mpro was ascribed to flaronates, flavonoids, and pseudo-o-peptides. The most hopeful inhibitors of SARS-CoV-2 were found to be oligomers of phloroglucin (1,3,5-trihydroxybenzene) derived from brown algae *Sargassum spinuligerum*. In turn, the most active inhibitors of SARS-CoV-2 are compounds from the florotannin group (8,8'-Bieckol, 6,6'-Bieckol, Dieckol) isolated from the brown algae *Ecklonia cava* (Gentile *et al.*, 2020). Khalifa, Nawaz, *et al.* (2020) revealed that phacelianin, gentiodelphin, cyanodelphin, and tecophilin authentically interacted with the receptor binding site and catalytic dyad (Cys145 and His41) of SARS-CoV-2-3CLpro. Khalifa, Zhu, *et al.* (2020) also discovered that

pedunculagin, tercatain, and castalin intensely interacted with the receptor binding site and catalytic dyad (Cys145 and His41) of SARS-CoV-2-3CLpro. These studies unleashed that anthocyanins and/or tannins with specific structure could be used as effective anti-COVID-19 natural components. Meanwhile, **Fig 1B**. portrayed the interaction between tannins with SARS-CoV-2-3CLpro (Khalifa, Nawaz, *et al.*, 2020; Khalifa, Zhu, *et al.*, 2020). We summarized the antiviral activities from the recent studies of natural polyphenols from several plants and their mode of action against CoVs (**Table 4**).

4. The underlying mechanism of the antiviral activity of polyphenols

After the outbreak of SARS-CoV-2, there is an abundance of literature on plant-based materials such as traditional Chinese herbs as a possible cure for symptoms related to SARS like fever control and faster chest clearing. Nonetheless, further clinical trials are needed in treating acute respiratory tract infections (Wu *et al.*, 2008). It remains important to recognize new therapeutic and functional agents of infectious diseases using natural sources (Cragg and Newman, 2013). During the last decade, scientists have made significant efforts to classify several polyphenols with anti-SARS-CoV activity. However, because of SARS-CoV and SARS-CoV-2 homology, these earlier findings regard naturally occurring substances could be capable of inhibiting SARS-CoV-2. A 3CLpro is essential for virus replication and is thus a potential drug goal for the production of therapeutic materials for SARS-CoV as well as SARS-CoV-2 (Yang *et al.*, 2020). It could be explained that bioactive compounds extracted from herbs were effectively inhibited the enzymatic activity of SARS-3CLpro in a dose-dependent manner. Rhubarb extracts (an IC_{50} of $13.76 \pm 0.03 \mu\text{g mL}^{-1}$) water extract (Luo *et al.*, 2009), *Houttuynia cordata* water extract (Lau *et al.*, 2008), litchi seeds-flavonoid extracted (Gong *et al.*, 2008), and extracted from *Isatis indigotica* root (Lin *et al.*, 2005) were efficiently effective against CoVs. Further, subsequent herb-derived occurring complexes-

318 includes herbacetin, -rhoifolin, pectolinarin (Jo *et al.*, 2020), sinigrin (IC₅₀: 217 µM), indigo
 319 (IC₅₀: 752 µM), -aloe-emodin (IC₅₀: 366 µM), hesperetin- (IC₅₀:8.3-µM) (Lin *et al.*, 2005),
 320 quercetin- (IC₅₀: 73 µM), epigallocatechin-gallate- (IC₅₀: 73 µM), and gallocatechin-gallate
 321 (IC₅₀: 47 µM) (Nguyen *et al.*, 2012), inhibited SARS 3CLpro effects. In addition, flavonoids,
 322 including isobavaschalcone, herbacetin, helichrysetin, and quercetin 3-β-D-glucoside, could
 323 inhibit the enzymatic action of MERS-CoV 3CLpro (Jo *et al.*, 2019). Helicase protein is often
 324 thought to be a possible focus for the production of anti-HCoV agents. Yu *et al.* stated that
 325 myricetin and scutellarein had a strong inhibition of nsP13 (SARS-CoV-helicase-protein) *in*
 326 *vitro* by influencing the function of ATPase (Yu *et al.*, 2012). Wu *et al.* conducted a massive-
 327 scale review of emerging medicines, natural products, and synthetic substances (>10000
 328 substances) to classify successful anti-SARS-CoV agents via a cell-based technique of
 329 SARS-CoV and Vero-E6-cells. Ginsenoside Rb1 extracted from *Panax-ginseng*, aescin
 330 extracted from horse chestnut, reserpine in the eucalyptus, Rauwolfia genus extracts, and
 331 *Lonicera japonica* blocked SARS-CoV replication at non-toxic concentrations (Wu *et al.*,
 332 2004). Same as SARS-CoV and HCoV-NL63, SARS-CoV-2-uses the ACE2-host receptor for
 333 cellular entry (Letko *et al.*, 2020). Thus, traditional Chinese medicine with ACE₂ goal
 334 capability keeps a pledge to avoid SARS-CoV-2 infection. In addition, disruption of the 3a
 335 receptor by kaempferol-juglanin derivatives may avoid viral release from contaminated cells
 336 (Schwarz *et al.*, 2014). Significantly, in compromised patients with SARS-CoV, MERS-CoV,
 337 and SARS-CoV-2 infections, the overpowering inflammatory responses have remarkably
 338 occurred. As a consequence, anti-inflammatory drugs are expected to reduce the incidence
 339 and mortality risk (Lu, 2020). In addition, a variety of anti-CoV agents have been reported
 340 from natural herbs, but the mechanisms of action have not yet been explained. Recently, the
 341 Chinese Health Commission has officially confirmed that natural medicinal herbs could be
 342 used in combination with traditional medicine for the treatment of COVID-19 patients (Yang

et al., 2020). Currently, experimental research on the efficacy of natural and medicinal plants focuses on the therapeutic potential of polyphenols against SARS-CoV-2 (Yang *et al.*, 2020). As the homogeneity of SARS-CoV is closely similar to SARS-CoV-2, many experiments demonstrating the anti-SARS-CoV-2 behavior of polyphenols or are planned to be released in the immediate future. Medicinal plants provide helpful and preventive treatments but are still desperately needed for the care of patients with SARS-CoV-2 infection, and experience with polyphenols is certainly worth researching. More analysis of polyphenols may contribute to the discovery of innovative anti-CoV compounds that could probably prove effective in the treatment of SARS-CoV-2 or other evolving lethal viral diseases as traditional therapeutic agents. Experimental searching with medicinal plant fields still has some risks and needs massive and further experiments. Polyphenols therapies traditionally used to treat the viral respiratory infection can include specific anti-SARS-CoV-2 substances. This research described many medicinal plants labeled as antiviral / pneumonia-effective which could specifically inhibit SARS-CoV-2 (Zhang *et al.*, 2020). Letko *et al.* (2020) confirmed how dissimilar lineage B-viruses can recombine to increase passage into human cells and have agreed that human ACE₂ is the receptor for the recent SARS-CoV-2 development. However, polyphenols including flavonoids as antiviral agents against MERS, SARS-CoV, and SARS-CoV-2 are not hardly investigated and more practical experiments are required. **Fig 2.** illustrated the role of polyphenols against MERS, SARS-CoV, and SARS-CoV-2.

5. Future needs

This review delivers a comprehensive range of evidence on the utilization of polyphenols extracted from medicinal plants for the reserve of CoVs. The cases obviously demonstrate that the designs with specific components of polyphenols and/or herb extracts that have been established for centuries in traditional medicine can be useful as the basis of anti-SARS-CoV

components in the treating and/or deterrence of COVID-19. Natural polyphenols can act as anti-enzymes, antioxidants, and through binding-blocking virus exterior protein receptors. Different plants can be utilized as a source of polyphenols with a wide range of biological activity. These components offer hope that polyphenol extracts accessible in this review may have antiviral activity, also against CoVs-family. The wide range of natural polyphenols presented in different sources, which are undervalued in conformist medicine, may establish a nearly infinite source of remedies. The latest tendencies in medicine and biotechnology have allowed the usage of polyphenols to a greater level, than in the earlier decades, mainly as nutritional complements and nutraceuticals. Their toxicity altogether with ingesting time is yet to be explored by aims of multidisciplinary tactics (proteomics, omics science-genomics, and metabolomics). New formulation technologies and processing might aid to improve the antiviral polyphenols solubility, delivery tactics, and therapeutic effects to adjust them as antiviral functional foods and remedies. The research achieved so far have added to the information of the chemical structure of antiviral polyphenols. Bioinformatic studies of the polyphenols-pharmacophores structures can chief to scheming new antiviral polyphenols drugs. The usage of plant extracts might deliver synergistic health useful effects i.e., anti-inflammation, antioxidant, antimutagenic, and antiviral among occurring polyphenol blends, which should be considered and must be intensely discovered.

6. Conclusion

Numerous instances display that nature proposes the greatest solutions. Many plant species hold biologically active polyphenols, which in synergistic blends are active in battling many diseases. They already have been discovered by ancient medicine and/or folk. Recognizing individual components and revealing the underlying mechanisms is a challenge. This delays the product's standardization, namely since plant extracts hold a wide variety of different

components, happening in fluctuating concentrations, with seemingly synergistic effects. Despite several unknowns, studies decidedly show that polyphenols inhibit the life cycle of CoVs. This review indicates the huge possibility of polyphenols in the expansion of functional foods and/or pharmaceutical preparations. Laboratory model trials are required to verify their efficiency in inhibiting CoVs-replication. The upcoming phase should be to conduct clinical trials on COVID-19-patients and remarking the possibility to decrease virus multiplication in the patient's organism and thus recover clinical signs. It appears that the addition of polyphenols into food and clinical practice should be fairly fast because those are mixtures of natural origin, attained from plants approved for use as herbs and food. It is worth stating that the advantage of using polyphenols is related to the high-safety profile without causing major side effects.

Acknowledgments

This work was self-funded.

Declaration of competing interest

The authors declared that there is no conflict of interest.

References

- Alpert, P. T. (2017). The role of vitamins and minerals on the immune system. *Home Health Care Management & Practice*, 29(3), 199-202.
- Banerjee, A., Kulcsar, K., Misra, V., Frieman, M. & Mossman, K. (2019). Bats and coronaviruses. *Viruses*, 11(1), 41.
- Calder, P. C. (2013). Feeding the immune system. *Proceedings of the Nutrition Society*, 72(3), 299-309.

- Calder, P. C. & Kew, S. (2007). The immune system: A target for functional foods?. *British Journal of Nutrition*, 88(S2), S165-S176.
- Castelo-Branco, C. & Soveral, I. (2014). The immune system and aging: A review. *Gynecological Endocrinology*, 30(1), 16-22.
- Chen, C.J., Michaelis, M., Hsu, H.K., Tsai, C.C., Yang, K. D., Wu, Y.C., Cinatl Jr, J. & Doerr, H. W. (2008). Toona sinensis roem tender leaf extract inhibits sars coronavirus replication. *Journal of Ethnopharmacology*, 120(1), 108-111.
- Chen, L., Li, J., Luo, C., Liu, H., Xu, W., Chen, G., Liew, O. W., Zhu, W., Puah, C. M. & Shen, X. (2006). Binding interaction of quercetin-3- β -galactoside and its synthetic derivatives with SARS-CoV 3CLpro: Structure–activity relationship studies reveal salient pharmacophore features. *Bioorganic & medicinal chemistry*, 14(24), 8295-8306.
- Chen, Z. & Nakamura, T. (2004). Statistical evidence for the usefulness of chinese medicine in the treatment of sars. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 18(7), 592-594.
- Chen, C., Zuckerman, D. M., Brantley, S., Sharpe, M., Childress, K., Hoiczky, E., & Pendleton, A. R. (2014). *Sambucus nigra* extracts inhibit infectious bronchitis virus at an early point during replication. *BMC veterinary research*, 10(1), 1-12.
- Cheng, Y. T., Rodak, D., Wong, C. & Hayden, C. (2006). Effects of micro-and nano-structures on the self-cleaning behaviour of lotus leaves. *Nanotechnology*, 17(5), 1359.
- Chiew, K., Phoon, M., Putti, T., Tan, B. K. & Chow, V. T. (2016). Evaluation of antiviral activities of houttuynia cordata thunb. Extract, quercetin, quercetrin and cinanserin on murine coronavirus and dengue virus infection. *Asian Pacific journal of tropical medicine*, 9(1), 1-7.

- Chojnacka, K., Witek-Krowiak, A., Skrzypczak, D., Mikula, K. & Młynarz, P. (2020). Phytochemicals containing biologically active polyphenols as an effective agent against covid-19-inducing coronavirus. *Journal of Functional Foods*, 73, 104146.
- Chowdhury, P., Sahuc, M. E., Rouillé, Y., Rivière, C., Bonneau, N., Vandeputte, A., ... & Séron, K. (2018). Theaflavins, polyphenols of black tea, inhibit entry of hepatitis C virus in cell culture. *PloS one*, 13(11), e0198226.
- Cohen, J. & Normile, D. 2020. "New sars-like virus in china triggers alarm." In.: American Association for the Advancement of Science. *Science*, 367(6475), 234-235.
- Comalada, M., Ballester, I., Bailón, E., Sierra, S., Xaus, J., Gálvez, J., Medina, F. S. d. & Zarzuelo, A. (2006). Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure–activity relationship. *Biochemical Pharmacology*, 72(8), 1010-1021.
- Cragg, G. M. & Newman, D. J. (2013). Natural products: A continuing source of novel drug leads. *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1830(6), 3670-3695.
- Del Bo, C., Bernardi, S., Marino, M., Porrini, M., Tucci, M., Guglielmetti, S., Cherubini, A., Carrieri, B., Kirkup, B. & Kroon, P. (2019). Systematic review on polyphenol intake and health outcomes: Is there sufficient evidence to define a health-promoting polyphenol-rich dietary pattern?. *Nutrients*, 11(6), 1355.
- Deng, J., Tang, X., Wei, X., Yu, C., Zhang, X., Shi, L. & Song, Y. (2020). Analysis method for the operation health of all-fiber optical current transformer. In 2019 International Conference on Optical Instruments and Technology: Optical Sensors and Applications (Vol. 11436, p. 114360T). International Society for Optics and Photonics.
- Forni, C., Facchiano, F., Bartoli, M., Pieretti, S., Facchiano, A., D'Arcangelo, D., Norelli, S., Valle, G., Nisini, R. & Beninati, S. J. B. r. i. (2019). Beneficial role of phytochemicals

on oxidative stress and age-related diseases. *BioMed Research International*, 2019, 8748253.

Fraga, C. G., Croft, K. D., Kennedy, D. O. & Tomás-Barberán, F. A. (2019). The effects of polyphenols and other bioactives on human health. *Food & Function*, 10(2), 514-528.

Gentile, D., Patamia, V., Scala, A., Sciortino, M. T., Piperno, A. & Rescifina, A. (2020). Putative inhibitors of SARS-CoV-2 main protease from a library of marine natural products: A virtual screening and molecular modeling study. *Marine drugs*, 18(4), 225.

Ge, M., Xiao, Y., Chen, H., Luo, F., Du, G., & Zeng, F. (2018). Multiple antiviral approaches of (–)-epigallocatechin-3-gallate (EGCG) against porcine reproductive and respiratory syndrome virus infection in vitro. *Antiviral research*, 158, 52-62.

Ghosh, R., Chakraborty, A., Biswas, A. & Chowdhuri, S. (2020). Evaluation of green tea polyphenols as novel corona virus (SARS CoV-2) main protease (Mpro) inhibitors—an in silico docking and molecular dynamics simulation study. *Journal of Biomolecular Structure Dynamics*, 1-13.

Gong, S., Su, X., Wu, H., Li, J., Qin, Y., Xu, Q. & Luo, W. (2008). A study on anti-SARS-CoV 3CL protein of flavonoids from litchi chinensis sonn core. *Chinese Pharmacological Bulletin*, 24(5), 699-700.

Guiné, R., Lima, M. & Barroca, M. (2009). Role and health benefits of different functional food components. *Millenium*, 1-17.

Hasan, A., Nahar, N., Jannat, K., Bondhon, T. A., Jahan, R. & Rahmatullah, M. (2020). In silico studies on phytochemicals of *pimpinella anisum* L.(apiaceae) as potential inhibitors of SARS-CoV-2 3C-like protease. *International Journal of Ayurveda*, 5(11),08-14.

- Horne, J. R. & Vohl, M.-C. (2020). Biological plausibility for interactions between dietary fat, resveratrol, ACE2, and SARS-CoV illness severity. *American Journal of Physiology-Endocrinology and Metabolism*, 318(5), E830-E833.
- Iwatani, S. & Yamamoto, N. (2019). Functional food products in japan: A review. *Food Science and Human Wellness*, 8(2), 96-101.
- Jo, S., Kim, H., Kim, S., Shin, D. H. & Kim, M. S. (2019). Characteristics of flavonoids as potent MERS-CoV 3C-like protease inhibitors. *Chemical Biology & Drug Design*, 94(6), 2023-2030.
- Kesel, A. J. (2005). Synthesis of novel test compounds for antiviral chemotherapy of severe acute respiratory syndrome (SARS). *Current Medicinal Chemistry*, 12(18), 2095-2162.
- Khaerunnisa, S., Kurniawan, H., Awaluddin, R., Suhartati, S. & Soetjipto, S. (2020). Potential inhibitor of COVID-19 main protease (Mpro) from several medicinal plant compounds by molecular docking study. *Preprints*. doi10. 20944/preprints202003.0226. v1, 1-14.
- Khalifa, I., Nawaz, A., Sobhy, R., Althwab, S. A. & Barakat, H. (2020). Polyacylated anthocyanins constructively network with catalytic dyad residues of 3CLpro of 2019-NCOV than monomeric anthocyanins: A structural-relationship activity study with 10 anthocyanins using *in-silico* approaches. *Journal of Molecular Graphics and Modelling*, 100, 107690.
- Khalifa, I., Zhu, W., Mohammed, H. H. H., Dutta, K. & Li, C. (2020). Tannins inhibit SARS-CoV-2 through binding with catalytic dyad residues of 3CLpro: An *in silico* approach with 19 structural different hydrolysable tannins. *Journal of Food Biochemistry*, 44(10), e13432.

- 512 Khalifa, I., Xia, D., Dutta, K., Peng, J., Jia, Y., & Li, C. (2020). Mulberry anthocyanins exert
 513 anti-AGEs effects by selectively trapping glyoxal and structural-dependently blocking
 514 the lysyl residues of β -lactoglobulins. *Bioorganic Chemistry*, 96, 103615.
- 515 Khan, H., Sureda, A., Belwal, T., Çetinkaya, S., Süntar, İ., Tejada, S., Devkota, H. P., Ullah,
 516 H. & Aschner, M. (2019). Polyphenols in the treatment of autoimmune diseases.
 517 *Autoimmunity Reviews*, 18(7), 647-657.
- 518 Kim, H. Y., Eo, E. Y., Park, H., Kim, Y. C., Park, S., Shin, H. J., & Kim, K. (2010).
 519 Medicinal herbal extracts of Sophorae radix, Acanthopanax cortex, Sanguisorbae
 520 radix and Torilis fructus inhibit coronavirus replication *in vitro*. *Antivir Ther*, 15(5),
 521 697-709.
- 522 Kim, D. W., Seo, K. H., Curtis-Long, M. J., Oh, K. Y., Oh, J. W., Cho, J. K., Lee, K. H. &
 523 Park, K. H. (2014). Phenolic phytochemical displaying SARS-CoV papain-like
 524 protease inhibition from the seeds of psoralea corylifolia. *Journal of Enzyme Inhibition*
 525 *and Medicinal Chemistry*, 29(1), 59-63.
- 526 Lau, K.M., Lee, K.M., Koon, C.M., Cheung, C. S.F., Lau, C.P., Ho, H.M., Lee, M. Y.H., Au,
 527 S. W.N., Cheng, C. H.K. & Bik-San Lau, C. (2008). Immunomodulatory and anti-sars
 528 activities of houttuynia cordata. *Journal of Ethnopharmacology*, 118(1), 79-85.
- 529 Letko, M., Marzi, A. & Munster, V. (2020). Functional assessment of cell entry and receptor
 530 usage for SARS-CoV-2 and other lineage b betacoronaviruses. *Nature Microbiology*,
 531 5(4), 562-569.
- 532 Liang, W., He, L., Ning, P., Lin, J., Li, H., Lin, Z., Kang, K. & Zhang, Y. (2015). (+)-
 533 Catechin inhibition of transmissible gastroenteritis coronavirus in swine testicular cells
 534 is involved its antioxidation. *Research in veterinary science*, 103, 28-33.

- 535 Lin, C.W., Tsai, F.J., Tsai, C.H., Lai, C.C., Wan, L., Ho, T.Y., Hsieh, C.C. & Chao, P.D. L.
 536 (2005). Anti-sars coronavirus 3c-like protease effects of isatis indigotica root and plant-
 537 derived phenolic compounds. *Antiviral Research*, 68(1), 36-42.
- 538 London, C. (2010). Functional foods that boost the immune system. *Functional Food Product*
 539 *Development*, 2, 295-321.
- 540 Lopez-Varela, S., Gonzalez-Gross, M. & Marcos, A. (2002). Functional foods and the
 541 immune system: A review. *European Journal of Clinical Nutrition*, 56(3), S29-S33.
- 542 Lu, H. (2020). Drug treatment options for the 2019-new coronavirus (2019-ncov). *BioScience*
 543 *Trends* 14(1), 69-71.
- 544 Luo, W., Su, X., Gong, S., Qin, Y., Liu, W., Li, J., Yu, H. & Xu, Q. (2009). Anti-sars
 545 coronavirus 3c-like protease effects of rheum palmatum l. Extracts. *Bioscience Trends*,
 546 3(4), 124-126.
- 547 Maggini, S., Pierre, A. & Calder, P. C. (2018). Immune function and micronutrient
 548 requirements change over the life course. *Nutrients*, 10(10), 1531.
- 549 Maggini, S., Wintergerst, E. S., Beveridge, S. & Hornig, D. H. (2007). Selected vitamins and
 550 trace elements support immune function by strengthening epithelial barriers and cellular
 551 and humoral immune responses. *British Journal of Nutrition*, 98 Suppl 1, S29-35.
- 552 Manoharan, Y., Haridas, V., Vasanthakumar, K., Muthu, S., Thavoorullah, F. F. & Shetty, P.
 553 (2020). Curcumin: A wonder drug as a preventive measure for covid19 management.
 554 *Indian Journal of Clinical Biochemistry*, 35(3), 373-375.
- 555 Maryam, M., Te, K. K., Wong, F. C., Chai, T. T., Gary, K. & Gan, S. C. (2020). Antiviral
 556 activity of traditional chinese medicinal plants dryopteris crassirhizoma and morus alba
 557 against dengue virus. *Journal of Integrative Agriculture*, 19(4), 1085-1096.
- 558 Mohd, A., Zainal, N., Tan, K. K., & AbuBakar, S. (2019). Resveratrol affects Zika virus
 559 replication *in vitro*. *Scientific reports*, 9(1), 1-11.

- 560 Morán-Santibañez, K., Peña-Hernández, M. A., Cruz-Suárez, L. E., Ricque-Marie, D.,
 561 Skouta, R., Vasquez, A. H., Rodríguez-Padilla, C. & Trejo-Avila, L. M. (2018).
 562 Virucidal and synergistic activity of polyphenol-rich extracts of seaweeds against
 563 measles virus. *Viruses*, 10(9), 465.
- 564 Pekmezci, D. (2011). Vitamin E and immunity. In *Vitamins & Hormones* (Vol. 86, pp. 179-
 565 215). Academic Press, New York, USA.
- 566 Negida, A. & Bahbah, E. I. 2020. "Comments on the hydroxychloroquine/azithromycin study
 567 and lessons for future clinical trials of covid-19 treatments." Preprint.
- 568 Nguyen, T. T. H., Woo, H.-J., Kang, H.-K., Kim, Y.-M., Kim, D.-W., Ahn, S.-A., Xia, Y. &
 569 Kim, D. (2012). Flavonoid-mediated inhibition of sars coronavirus 3c-like protease
 570 expressed in pichia pastoris. *Biotechnology Letters*, 34(5), 831-838.
- 571 Palamara, A. T., Nencioni, L., Aquilano, K., De Chiara, G., Hernandez, L., Cozzolino, F.,
 572 Ciriolo, M.R. & Garaci, E. (2005). Resveratrol inhibits Influenza A virus replication *in*
 573 *vitro* and *in vivo*. *Journal of Infectious Diseases*, 191, 1719-1729.
- 574 Pandey, P., Rane, J. S., Chatterjee, A., Kumar, A., Khan, R., Prakash, A. & Ray, S. (2020).
 575 Targeting SARS-CoV-2 spike protein of covid-19 with naturally occurring
 576 phytochemicals: An in silico study for drug development. *Journal of Biomolecular*
 577 *Structure Dynamics*, 1-11.
- 578 Paraiso, I. L., Revel, J. S. & Stevens, J. F. (2020). Potential use of polyphenols in the battle
 579 against covid-19. *Current Opinion in Food Science*, 32, 149-155.
- 580 Park, H. R., Yoon, H., Kim, M. K., Lee, S. D. & Chong, Y. (2012). Synthesis and antiviral
 581 evaluation of 7-o-arylmethylquercetin derivatives against sars-associated coronavirus
 582 (scv) and hepatitis c virus (hcv). *Archives of pharmacal research*, 35(1), 77-85.
- 583 Park, J.-Y., Kim, J. H., Kwon, J. M., Kwon, H.-J., Jeong, H. J., Kim, Y. M., Kim, D., Lee, W.
 584 S. & Ryu, Y. B. (2013). Dieckol, a SARS-CoV 3CLpro inhibitor, isolated from the

edible brown algae ecklonia cava. *Bioorganic & Medicinal Chemistry*, 21(13), 3730-3737.

Park, J.-Y., Yuk, H. J., Ryu, H. W., Lim, S. H., Kim, K. S., Park, K. H., Ryu, Y. B. & Lee, W. S. (2017). Evaluation of polyphenols from broussonetia papyrifera as coronavirus protease inhibitors. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 32(1), 504-512.

Pereira, P. C. M. (2003). Interaction between infection, nutrition and immunity in tropical medicine. *Journal of Venomous Animals and Toxins including Tropical Diseases*, 9, 163-173.

Pillaiyar, T., Meenakshisundaram, S., & Manickam, M. (2020). Recent discovery and development of inhibitors targeting coronaviruses. *Drug discovery today*, 25(4), 668-688.

Powell, J., Borchers, A. T., Yoshida, S., & Gershwin, M. E. (2000). Evaluation of the immune system in the nutritionally at-risk host. In *Nutrition and Immunology* (pp. 21-31). Humana Press, Totowa, NJ.

Quero, J., Mármol, I., Cerrada, E., & Rodríguez-Yoldi, M. J. (2020). Insight into the potential application of polyphenol-rich dietary intervention in degenerative disease management. *Food & Function*, 11(4), 2805-2825.

Raekiansyah, M., Buerano, C. C., Luz, M. A. D., & Morita, K. (2018). Inhibitory effect of the green tea molecule EGCG against dengue virus infection. *Archives of virology*, 163(6), 1649-1655.

Rasouli, H., Farzaei, M. H. & Khodarahmi, R. (2017). Polyphenols and their benefits: A review. *International Journal of Food Properties*, 20(sup2), 1700-1741.

Roberfroid, M. B. (2000). Concepts and strategy of functional food science: The european perspective. *The American journal of clinical nutrition*, 71(6), 1660S-1664S.

- Rothwell, J. A., Urpi-Sarda, M., Boto-Ordoñez, M., Llorach, R., Farran-Codina, A., Barupal, D. K., Neveu, V., Manach, C., Andres-Lacueva, C. & Scalbert, A. (2016). Systematic analysis of the polyphenol metabolome using the phenol-explorer database. *Molecular Nutrition and Food Research*, 60(1), 203-211.
- Ryu, Y. B., Jeong, H. J., Kim, J. H., Kim, Y. M., Park, J.-Y., Kim, D., Naguyen, T. T. H., Park, S.-J., Chang, J. S., Park, K. H., Rho, M.-C. & Lee, W. S. (2010). Biflavonoids from *torreya nucifera* displaying SARS-CoV 3CLpro inhibition. *Bioorganic & Medicinal Chemistry*, 18(22), 7940-7947.
- Ryu, Y. B., Park, S.-J., Kim, Y. M., Lee, J.-Y., Seo, W. D., Chang, J. S., Park, K. H., Rho, M.-C. & Lee, W. S. (2010). SARS-CoV 3CLpro inhibitory effects of quinone-methide triterpenes from *tripterygium regelii*. *Bioorganic & Medicinal Chemistry Letters*, 20(6), 1873-1876.
- Santos-Buelga, C., González-Paramás, A. M., Oludemi, T., Ayuda-Durán, B. & González-Manzano, S. (2019). Plant phenolics as functional food ingredients. *Advances in Food and Nutrition Research*, 90, 183-257.
- Sekizawa, H., Ikuta, K., Ohnishi-Kameyama, M., Nishiyama, K., & Suzutani, T. (2019). Identification of the Components in a *Vaccinium oldhamii* Extract Showing Inhibitory Activity against Influenza Virus Adsorption. *Foods*, 8(5), 172.
- Schaible, U. E. & Stefan, H. (2007). Malnutrition and infection: Complex mechanisms and global impacts. *PLoS Medicine*, 4(5), e115.
- Schettig, R., Sears, T., Klein, M., Tan-Lim, R., Matthias Jr, R., Aussems, C., Hummel, M., Sears, R., Poteet, Z. & Warren, D. (2020). Covid-19 patient with multifocal pneumonia and respiratory difficulty resolved quickly: Possible antiviral and anti-inflammatory benefits of quercinex (nebulized quercetin-nac) as adjuvant. *Advances in Infectious Diseases*, 10(3), 45-55.

- Schwarz, S., Sauter, D., Wang, K., Zhang, R., Sun, B., Karioti, A., Bilia, A. R., Efferth, T. & Schwarz, W. (2014). Kaempferol derivatives as antiviral drugs against the 3a channel protein of coronavirus. *Planta Medica*, 80(02/03), 177-182.
- Shinbo, Y., Nakamura, Y., Altaf-Ul-Amin, M., Asahi, H., Kurokawa, K., Arita, M., ... & Kanaya, S. (2006). KNApSACk: a comprehensive species-metabolite relationship database. In *Plant metabolomics* (pp. 165-181). Springer, Berlin, Heidelberg.
- Solnier, J., & Fladerer, J. P. (2020). Flavonoids: A complementary approach to conventional therapy of COVID-19?. *Phytochemistry Reviews*, 1-23.
- Silva, R. F. M. & Pogačnik, L. (2020). Polyphenols from food and natural products: Neuroprotection and safety. *Antioxidants*, 9(1), 61.
- Sundararajan, A., Ganapathy, R., Huan, L., Dunlap, J. R., Webby, R. J., Kotwal, G. J. & Sangster, M. Y. (2010). Influenza virus variation in susceptibility to inactivation by pomegranate polyphenols is determined by envelope glycoproteins. *Antiviral Research*, 88(1), 1-9.
- Vela, E. M., Knostman, K. A., Mott, J. M., Warren, R. L., Garver, J. N., Vela, L. J. & Stammen, R. L. (2010). Genistein, a general kinase inhibitor, as a potential antiviral for arenaviral hemorrhagic fever as described in the pirital virus-syrian golden hamster model. *Antiviral Research*, 87(3), 318-328.
- Villagomez, J. (2017). *In Vitro* Antiviral Activity of Black Tea Polyphenols on Sindbis Virus in Vero Cells. MSc Thesis, Montclair State University, New Jersey, USA.
- Wang, M., Carver, J. J., Phelan, V. V., Sanchez, L. M., Garg, N., Peng, Y., ... & Porto, C. (2016). Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nature biotechnology*, 34(8), 828-837.
- Wang, X. & Liu, Z. (2014). Prevention and treatment of viral respiratory infections by traditional chinese herbs. *Chinese medical journal*, 127(7), 1344-1350.

- 660 Wen, C.-C., Shyur, L.-F., Jan, J.-T., Liang, P.-H., Kuo, C.-J., Arulselvan, P., Wu, J.-B., Kuo,
 661 S.-C. & Yang, N.-S. (2011). Traditional chinese medicine herbal extracts of cibotium
 662 barometz, gentiana scabra, dioscorea batatas, cassia tora, and taxillus chinensis inhibit
 663 SARS-CoV replication. *Journal of traditional and complementary medicine*, 1(1), 41-
 664 50.
- 665 Weng, J. R., Lin, C. S., Lai, H. C., Lin, Y. P., Wang, C. Y., Tsai, Y. C. & Lin, C. W. (2019).
 666 Antiviral activity of sambucus formosananakai ethanol extract and related phenolic acid
 667 constituents against human coronavirus NL63. *Virus research*, 273, 197767.
- 668 Wichers, H. (2009). Immunomodulation by food: Promising concept for mitigating allergic
 669 disease? *Analytical and Bioanalytical Chemistry*, 395(1), 37-45.
- 670 Wu, C.-Y., Jan, J.-T., Ma, S.-H., Kuo, C.-J., Juan, H.-F., Cheng, Y.-S. E., Hsu, H.-H., Huang,
 671 H.-C., Wu, D. & Brik, A. (2004). Small molecules targeting severe acute respiratory
 672 syndrome human coronavirus. *Proceedings of the National Academy of Sciences*,
 673 101(27), 10012-10017.
- 674 Wu, T., Yang, X., Zeng, X. & Poole, P. (2008). Traditional chinese medicine in the treatment
 675 of acute respiratory tract infections. *Respiratory Medicine*, 102(8), 1093-1098.
- 676 Xu, Z., Peng, C., Shi, Y., Zhu, Z., Mu, K., Wang, X. & Zhu, W. (2020). Nelfinavir was
 677 predicted to be a potential inhibitor of 2019-ncov main protease by an integrative
 678 approach combining homology modelling, molecular docking and binding free energy
 679 calculation. *BioRxiv*, 2020.01.27.921627.
- 680 Yahfoufi, N., Alsadi, N., Jambi, M. & Matar, C. (2018). The immunomodulatory and anti-
 681 inflammatory role of polyphenols. *Nutrients*, 10(11), 1618.
- 682 Yang, Y., Islam, M. S., Wang, J., Li, Y. & Chen, X. (2020). Traditional chinese medicine in
 683 the treatment of patients infected with 2019-new coronavirus (SARS-CoV-2): A review
 684 and perspective. *International Journal of Biological Sciences*, 16(10), 1708-1717.

- Yi, L., Li, Z., Yuan, K., Qu, X., Chen, J., Wang, G., Zhang, H., Luo, H., Zhu, L. & Jiang, P. (2004). Small molecules blocking the entry of severe acute respiratory syndrome coronavirus into host cells. *Journal of Virology*, 78(20), 11334-11339.
- Yu, M.-S., Lee, J., Lee, J. M., Kim, Y., Chin, Y.-W., Jee, J.-G., Keum, Y.-S. & Jeong, Y.-J. (2012). Identification of myricetin and scutellarein as novel chemical inhibitors of the sars coronavirus helicase, nsp13. *Bioorganic & Medicinal Chemistry Letters*, 22(12), 4049-4054.
- Zhang, D.H., Wu, K.L., Zhang, X., Deng, S.Q. and Peng, B., 2020. *In silico* screening of Chinese herbal medicines with the potential to directly inhibit 2019 novel coronavirus. *Journal of Integrative Medicine*, 18(2), 152-158.
- Zhang, L., & Liu, Y. (2020). Potential interventions for novel coronavirus in China: A systematic review. *Journal of Medical Virology*, 92(5), 479-490.

Table 1. The Japanese "FOSHU" criteria for functional foods, (Iwatani and Yamamoto, 2019).

-
1. They can or should be consumed as a portion of a normal diet.
 2. They are foods: not capsules, pills, or powder-based on natural food components.
 3. They have a clear function on the human organism such as:
 - 3.1. Improving the immunity function.
 - 3.2. Supporting recovery from specific diseases.
 - 3.3. Preventing specific diseases.
 - 3.4. Slowing down the aging process.
 - 3.5. Controlling physical complaints.
-

Table 2. Overview of vital functional food components (non-polyphenols) in boosting up the immune response, virus targeted, and functions related, quoted from (London, 2010; Maggini *et al.*, 2018; Zhang and Liu, 2020).

Food components	Health properties	Virus targeted and related functions
Ω-3 fatty acids	<ul style="list-style-type: none"> Stimulation of the immune system. 	Influenza and human immunodeficiency virus.
Vitamin C	<ul style="list-style-type: none"> Stimulates production, function, and movement of leukocytes (e.g., lymphocytes, neutrophils, and phagocytes). Antimicrobial, natural killer cell activities, and chemotaxis. Regenerates other important antioxidants such as glutathione and vitamin E to their active state. Promotes collagen synthesis. Increases serum levels of complement proteins. Involved in apoptosis and clearance of spent neutrophils from sites of infection by macrophages. 	Avian coronavirus.
Vitamin D	<ul style="list-style-type: none"> Stimulates immune cell proliferation and cytokine production and helps protect against infection caused by pathogens. Expressed in innate immune cells (e.g., monocytes, macrophages, dendritic cells). 	Bovine coronavirus.
Vitamin A	<ul style="list-style-type: none"> Maintains the structural and functional integrity of mucosal cells in innate barriers (e.g., skin, and respiratory tract). 	Measles virus, human immunodeficiency virus, and avian coronavirus.
Vitamin B	<ul style="list-style-type: none"> Boosting the immune response of the host. 	MERS-CoV.

Vitamin E	<ul style="list-style-type: none"> • Impairs humoral and cell-mediated aspects of adaptive immunity, i.e., B and T cell function. 	Coxsackievirus and bovine coronavirus.
Se	<ul style="list-style-type: none"> • Important for the antioxidant host defense system affecting leukocyte. 	Influenza virus and avian coronavirus.
Fe	<ul style="list-style-type: none"> • Stimulation of the immune system. 	Viral mutations.
Zn	<ul style="list-style-type: none"> • Stimulation of the immune system. 	Measles and SARS-CoV.

Table 3. The antiviral activities of polyphenols, some examples from the available studies.

Polyphenols	Main sources	Conclusion	References
Theaflavins	Black tea	Theaflavins inhibited the Sindbis Virus infection by 99% at a concentration of 14.6mM. Act directly on HCV-viral particles and inhibit the ability to bind to the receptor surface.	Villagomez (2017) Chowdhury <i>et al.</i> (2018)
Epigallocatechin gallate, epicatechingallate and gallocatechin-3-gallate	Green tea	EGCG effectively inhibits porcine reproductive and respiratory syndrome virus infection and replication in porcine alveolar macrophages. It prevents MARC-145 cells. EGCG directly interacts with the Dengue virus molecule causing virus deformation and thus preventing the virus from infecting further cells.	Ge <i>et al.</i> (2018) Raekiansyah <i>et al.</i> (2018)
Different polyphenols	<i>Radix Sophorae</i> , <i>rhrizome</i> <i>Acanthopanacis</i> , <i>radix</i> <i>Sanguisorbae</i> and extract	Inhibits the effects on replication stages and/or their influence on cellular signal pathways of Mouse hepatitis virus.	(Kim <i>et al.</i> , 2010)

Torilis fructus

Quercetin and rutin	<i>Houttuynia cordata Thunb</i>	Inhibitory effects on the ATPase of mouse hepatitis and DENV type 2 virus. (Chiou <i>et al.</i> , 2016)
(+)-Catechin	Teas and pome fruits	Inhibiting effect on transmissible gastroenteritis virus proliferation <i>in vitro</i> and associated with its anti-oxidation. (Liang <i>et al.</i> , 2015)
Caffeic, chlorogenic, and gallic acids	<i>Sambucus Formosana Nakai</i>	Inhibition of the replication of HCoV NL63 in a cell-type independent manner. (Weng <i>et al.</i> , 2019)
Anthocyanins	<i>Sambucus nigra</i>	Inhibition of avian infectious bronchitis virus replication (Chen <i>et al.</i> , 2014)
Resveratrol	Skins of red fruits	Resveratrol inhibits the binding of Zika virus particles to cells and reduces circulating Zika virus particles, which underlines its potential of limiting disease severity during the viraemic phase. (Mohd <i>et al.</i> , 2019)
Resveratrol	Skins of red fruits	Resveratrol exhibited potent inhibitory effects against influenza in MDCK cells via the blockade of nuclear-cytoplasmic translocation of viral ribonucleoprotein complexes, (Palamara <i>et al.</i> , 2019)

the decrease in the expression of late viral proteins, and the inhibition of cellular protein kinase C activity and its dependent pathways.

Flavonoids	Elderberry and blueberry	Several blueberry varieties exert anti-influenza viral activities, especially viral adsorption, which positively correlated with the total polyphenol content in Madin-Darby canine kidney cells.	(Sekizawa <i>et al.</i> , 2019)
------------	--------------------------	---	---------------------------------

Table 4. The role of polyphenols against SARS-CoV, MERS-CoV, and SARS-CoV-2.

Polyphenols	Representative	Form/Source	Virus	Mechanism	Reference
Phenolic acids					
Hydrobenzoic acids	Gallic acid	Tetra- <i>O</i> -galloyl- β -D glucose from <i>Galla chinensis</i>	SARS-CoV	Avidly binds with surface spike protein of SARS-CoV.	(Yi <i>et al.</i> , 2004)
	Hydrobenzoic acid	Desmethoxyreserpine	SARS-CoV-2	Inhibit replication of 3CLpro, and entry.	(Kesel, 2005)
Flavonoids					
Flavonols	Kaempferol	Kaempferol derivatives, Kaempferol	SARS-CoV	Inhibit 3a ion channel of CoVs.	(Schwarz <i>et al.</i> , 2014)
			MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	(Park <i>et al.</i> , 2017)
	Quercetin	Quercetin, Quercetin 3- β -D-glucoside,	MERS	Inhibit cleavage activity of MERS-3CLpro	(Jo <i>et al.</i> , 2019)

	isobavachalcone, and helichrysetin		enzyme.	
	Quercetin, Quercetin- β - galactoside	MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	(Park <i>et al.</i> , 2017)
	Quercetin and TSL-1 from <i>Toona sinensis</i> Roem	SARS-CoV	Inhibit the cellular entry of SARS-CoV.	(Chen <i>et al.</i> , 2008)
	Quercetin	SARS-CoV-2	PLpro and 3CLpro enzyme.	(Park <i>et al.</i> , 2013)
Myricetin	Myricetin	SARS-CoV	Inhibit nsP13 by affecting the ATPase activity. SARS-CoV helicase inhibitor.	(Yu <i>et al.</i> , 2012)
Herbacetin	Herbacetin	MERS	Inhibit cleavage activity of MERS-3CLpro	(Jo <i>et al.</i> , 2019)

		enzyme.		
		SARS-CoV	block the enzymatic activity of SARS-CoV 3CLpro.	(Jo <i>et al.</i> , 2020)
Papyriflavonol A	<i>Broussonetia papyrifera</i>	MERS-CoV,	Inhibit PLpro.	(Park <i>et al.</i> ,
Kazinol A, B, F and J, and broussoflavan A		SARS-CoV	Inhibit SARS-3CLpro activity.	2017)
Amentoflavone,	<i>Torreya nucifera</i> leaves	SARS-CoV		(Ryu, Park,
		3CL pro inhibitor		<i>et al.</i> , 2010)
Kaempferol, quercetin, luteolin-7-glucoside, demethoxycurcumin, naringenin, apigenin-7- glucoside, oleuropein, curcumin, catechin,	Traditional herbs	Inhibitors of SARS-CoV-2-Mpro	block the enzymatic activity of SARS-CoV 3CLpro.	(Khaerunnisa <i>et al.</i> , 2020)

	epicatechingallate, zingerol, gingerol, and allicin				
Flavones	Apigenin	<i>Ocimum basilicum</i>	SARS-CoV	Inhibit PLpro. Inhibit SARS-CoVpro activity.	(Pandey <i>et al.</i> , 2020)
	Baicalin	<i>Scutellaria baicalensis</i>	SARS-CoV	Inhibit Angiotensin- converting enzyme.	(Chen and Nakamura, 2004)
	Scutellarein	<i>Scutellaria lateriflora</i>	SARS-CoV	Inhibit nsP13 by affecting the ATPase activity.	(Yu <i>et al.</i> , 2012)
	Rhoifolin	<i>Rhus succedanea.</i>	SARS-CoV	Inhibit SARS-3CLpro activity.	(Gong <i>et al.</i> , 2008; Nguyen <i>et al.</i> , 2012)

	Luteolin	luteolin, from <i>Veronica linifolia</i>	SARS-CoV	Avidly binds with surface spike protein of SARS-CoV.	(Yi <i>et al.</i> , 2004)
	Daidzein	Plant-derived phenolic compounds and Root extract of <i>Isatis indigotica</i>	SARS-CoV	Not active.	(Lin <i>et al.</i> , 2005)
	30-(3-methylbut-2-enyl)-30,4,7-trihydroxyflavone	<i>Broussonetia papyrifera</i>	MERS-CoV, SARS-CoV	Inhibition of cysteine proteases CoV	(Park <i>et al.</i> , 2017)
	neobavaisoflavone	<i>Psoralea corylifolia</i>	SARS-CoV	Inhibitory activity toward SARS-CoV PLpro.	(Kim <i>et al.</i> , 2014)
Flavanones	Herbacetin, Rhoifolin pectolinarin Tetra-O-galoyl- β -d-glucose (TGG)	Plant-derived phenolic compounds and Root extract of <i>Isatis indigotica</i> ,	SARS-CoV	Inhibit the cleavage activity of the SARS-3CLpro enzyme.	(Jo <i>et al.</i> , 2020; Lin <i>et al.</i> , 2005)

luteoline					
	Pelargonidin	<i>Pimpinella anisum</i>	SARS-CoV-2	Binding affinities to 3C-like protease of SARS-CoV-2	(Hasan <i>et al.</i> , 2020)
	Bavachinin	<i>Psoralea corylifolia</i>	SARS-CoV	Inhibitory activity toward SARS-CoV PLpro.	(Kim <i>et al.</i> , 2014)
Anthocyanidins	10 polyacylated and monomeric anthocyanins	Bure components	SARS-CoV-2	Constructively network with catalytic dyad residues of 3CLpro of SARS-CoV-2.	(Khalifa, Nawaz, <i>et al.</i> , 2020)
Flavanols	Epigallocatechin gallate	Green tea	SARS-CoV	Inhibit SARS-3CLpro activity.	(Gong <i>et al.</i> , 2008; Nguyen <i>et al.</i> , 2012)
	Gallocatechin gallate	Green tea	SARS-CoV	Inhibit SARS-3CLpro	(Gong <i>et al.</i> ,

	and epicatechingallate			activity.	2008; Nguyen <i>et al.</i> , 2012)
	galocatechin-3-gallate	Green tea	SARS-CoV	Inhibit SARS-3CLpro activity.	(Ghosh <i>et al.</i> , 2020)
Chalcone	Isoliquiritigenin	<i>Glycyrrhiza glabra</i>	MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	(Park <i>et al.</i> , 2017)
	Broussochalcone B, broussochalcone A, and 4- hydroxyisolonchocarpin	<i>Broussonetia papyrifera</i>	MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	(Park <i>et al.</i> , 2017)
	isobavachalcone	<i>Psoralea corylifolia</i>	SARS-CoV	inhibitory activity toward SARS-CoV PLpro.	(Kim <i>et al.</i> , 2014)
	4'-O-	<i>Broussonetia papyrifera</i>	MERS-CoV,	Inhibit PLpro.	(Park <i>et al.</i> ,

	methylbavachalcone		SARS-CoV	Inhibit SARS-3CLpro activity.	2017)
Tannins	19 hydrolysable tannins	Bure components	SARS-CoV-2	Efficacious and selective anti-COVID-19 therapeutic compounds.	(Khalifa, Zhu, <i>et al.</i> , 2020)

Figures legends:

Fig 1. The binding poses of amentoflavone in SARS-CoV 3CLpro. Ribbon plot of amentoflavone complexed to 3CLpro with hydrogen bonding (A). Fig B. represented the binding between tannins and 3CLpro of SARS-CoV-2, adopted from Khalifa *et al.* (2020) and Khalifa *et al.* (2020).

Fig 2. Underlying of the most important structural proteins of CoV are spike protein (S), membrane protein (M), envelop protein (E), and the nucleocapsid protein (N), the RNA genome of CoV is packed in the nucleocapsid protein and further covered with envelope. The hydroxyl group of flavonoids at 7-position looks important to bind at the binding site against 3-chymotrypsin-like protease (3CLpro), and papain-like protease (PLpro). Flavonoids as a vast reservoir of therapeutically active constituents as antiviral candidates against RNA viruses, adopted from Rasouli *et al.* (2017), Solnier *et al.* (2020), and Pillaiyar *et al.* (2020).




Fig 1.




Fig 2.

Highlights

Polyphenols possess many potential health benefits, including antiviral effects.

Polyphenols consider promising bioactive substances for COVID-19 prevention.

Polyphenol-rich foods will be helpful during the COVID-19's treatment.

Polyphenols assist as blocking agent for protease's enzymatic activity of COVID-19.

Conflict of interest

No conflict of interest among authors.