



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1766>

Title:	Observation of $\tau \rightarrow \pi^- \nu \tau e^+ e^-$ and search for $\tau \rightarrow \pi^- \nu \tau \mu^+ \mu^-$
Authors:	Bhardwaj, V. (/jspui/browse?type=author&value=Bhardwaj%2C+V.)
Keywords:	Statistical Efficiencies Mechanisms
Issue Date:	2019
Publisher:	American Physical Society
Citation:	Physical Review D, 100(7).
Abstract:	We present the first measurements of branching fractions of rare tau-lepton decays, $\tau \rightarrow \pi^- \nu \tau \ell^+ \ell^-$ ( $\ell = e$ or $\mu$ ), using a data sample corresponding to 562 fb <sup>-1</sup> collected at a center-of-mass energy of 10.58 GeV with the Belle detector at the KEKB asymmetric-energy $e^+ e^-$ collider. The $\tau \rightarrow \pi^- \nu \tau e^+ e^-$ decay is observed for the first time with 7.0 $\sigma$ significance. The partial branching fraction determined by the structure-dependent mechanisms mediated by either a vector or an axial-vector current for the mass region $M_{\pi e e} > 1.05$ GeV/c <sup>2</sup> is measured to be $B(\tau \rightarrow \pi^- \nu \tau e^+ e^-) [M_{\pi e e} > 1.05 \text{ GeV/c}^2] = (5.90 \pm 0.53 \pm 0.85 \pm 0.11) \times 10^{-6}$ , where the first uncertainty is statistical, the second is systematic, and the third is due to model dependence. In the full phase space, due to the different detection efficiencies for the structure-dependent mechanisms mediated by axial-vector and vector currents, the branching fraction varies from $B_A(\tau \rightarrow \pi^- \nu \tau e^+ e^-) = (1.46 \pm 0.13 \pm 0.21) \times 10^{-5}$ to $B_V(\tau \rightarrow \pi^- \nu \tau e^+ e^-) = (3.01 \pm 0.27 \pm 0.43) \times 10^{-5}$ , respectively. An upper limit is set on the branching fraction of the $\tau \rightarrow \pi^- \nu \tau \mu^+ \mu^-$ decay, $B(\tau \rightarrow \pi^- \nu \tau \mu^+ \mu^-) < 1.14 \times 10^{-5}$ , at the 90% confidence level.
Description:	Only IISERM authors are available in the record.
URI:	<a href="https://journals.aps.org/prd/abstract/10.1103/PhysRevD.100.071101">https://journals.aps.org/prd/abstract/10.1103/PhysRevD.100.071101</a> ( <a href="https://journals.aps.org/prd/abstract/10.1103/PhysRevD.100.071101">https://journals.aps.org/prd/abstract/10.1103/PhysRevD.100.071101</a> ) <a href="http://hdl.handle.net/123456789/1766">http://hdl.handle.net/123456789/1766</a> ( <a href="http://hdl.handle.net/123456789/1766">http://hdl.handle.net/123456789/1766</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/1766/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/1766/1/Need%20to%20add%20pdf.odt\)](#)

[Show full item record \(/jspui/handle/123456789/1766?mode=full\)](#)

[Statistics \(/jspui/handle/123456789/1766/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.