



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/3172>


Title:	Radio observations of magnetic cataclysmic variables
Authors:	Singh, Kulinder Pal (/jspui/browse?type=author&value=Singh%2C+Kulinder+Pal)
Keywords:	Cataclysmic variables Radio continuum Stars Stars Activity Stars Magnetic fields
Issue Date:	2020
Publisher:	Elsevier Ltd
Citation:	Advances in Space Research, 66(5), pp.1226-1234.
Abstract:	The NSF's Karl G. Jansky Very Large Array (VLA) is used to observe 122 magnetic cataclysmic variables (MCVs) during three observing semesters (13B, 15A, and 18A). We report radio detections of 33 stars with fluxes in the range 6–8031 μ Jy. Twenty-eight stars are new radio sources, increasing the number of radio detected MCVs to more than 40. A surprising result is that about three-quarters (24 of 33 stars) of the detections show highly circularly polarized radio emission of short duration, which is characteristic of electron cyclotron maser emission. We argue that this emission originates from the lower corona of the donor star, and not from a region between the two stars. Maser emission enables a more direct estimate of the mean coronal magnetic field of the donor star, which we estimate to be 1–4 kG assuming a magnetic filling factor of 50%. A two-sample Kolmogorov-Smirnov test supports the conclusion that the distribution function of radio detected MCVs with orbital periods between 1.5 and 5 h is similar to that of all MCVs. This result implies that rapidly-rotating ($P_{\text{spin}} < 10$ days), fully convective stars can sustain strong magnetic dynamos. These results support the model of that the change in angular momentum loss across the fully convective boundary at $P_{\text{orb}} \approx 3$ h is due to a change in the magnetic field structure of the donor star from a low-order to high-order multipolar field.
Description:	Only IISERM authors are available in the record.
URI:	https://www.sciencedirect.com/science/article/pii/S0273117720302453?via%3Dihub (https://www.sciencedirect.com/science/article/pii/S0273117720302453?via%3Dihub) http://hdl.handle.net/123456789/3172 (http://hdl.handle.net/123456789/3172)
Appears in	Research Articles (/jspui/handle/123456789/9)
Collections:	

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/3172/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/3172/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/3172?mode=full>)

 (</jspui/handle/123456789/3172/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.