



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4665>

Title:	Well-Defined Ni(0) and Ni(II) Complexes of Bicyclic (Alkyl)(Amino)Carbene (MeBICAAC): Catalytic Activity and Mechanistic Insights in Negishi Cross-Coupling Reaction
Authors:	Thakur, Sandeep Kumar (/jspui/browse?type=author&value=Thakur%2C+Sandeep+Kumar) Kaur, Mandeep (/jspui/browse?type=author&value=Kaur%2C+Mandeep) Manar, Krishna Kumar (/jspui/browse?type=author&value=Manar%2C+Krishna+Kumar) Adhikari, Manu (/jspui/browse?type=author&value=Adhikari%2C+Manu) Choudhury, Angshuman Roy (/jspui/browse?type=author&value=Choudhury%2C+Angshuman+Roy) Singh, Sanjay (/jspui/browse?type=author&value=Singh%2C+Sanjay)
Keywords:	Catalytic Activity Mechanistic Insights Cross-Coupling Reaction
Issue Date:	2022
Publisher:	Wiley
Citation:	Chemistry - A European Journal, 28(59), 2202237
Abstract:	Negishi cross-coupling reaction of organozinc compounds as nucleophiles with aryl halides has drawn immense focus for C–C bond formation reactions. In comparison to the well-established library of Pd complexes, the C–C cross-coupling of this particular approach is largely primitive with nickel-complexes. Herein, we describe the syntheses of Ni(II) complexes, [(MeBICAAC) ₂ NiX ₂] (X=Cl (1), Br (2), and I (3)) by employing the bicyclic (alkyl)(amino)carbene (MeBICAAC) ligand. The reduction of complexes 1–3 using KC ₈ afforded the two coordinate low valent, Ni(0) complex, [(MeBICAAC) ₂ Ni(0)] (4). Complexes 1–4 have been characterized by spectroscopic techniques and their solid-state structures were also confirmed by X-ray crystallography. Furthermore, complexes 1–4 have been applied in a direct and convenient method to catalyze the Negishi cross-coupling reaction of various aryl halides with 2,6-difluorophenylzinc bromide or phenylzinc bromide as the coupling partner in the presence of 3 mol % catalyst. Comparatively, among all-pristine complexes, 1 exhibit high catalytic potential to afford value-added C–C coupled products without the use of any additive. The UV-vis studies and HRMS measurements of controlled stoichiometric reactions vindicate the involvement of Ni(I)–Ni(III) cycle featured with a penta-coordinated Ni(III)-aryl species as the key intermediate for 1 whereas Ni(0)/Ni(II) species are potentially involved in the catalytic cycle of 4.
Description:	Only IISERM authors are available in the record
URI:	https://doi.org/10.1002/chem.202202237 (https://doi.org/10.1002/chem.202202237) http://hdl.handle.net/123456789/4665 (http://hdl.handle.net/123456789/4665)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:


File	Description	Size	Format

Need To Add...Full Text_PDF.
(/jspui/bitstream/123456789/4665/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF.)

15.36 Unknown
kB

[View/Open \(/jspui/](#)

[Show full item record \(/jspui/handle/123456789/4665?mode=full\)](#)

 [\(/jspui/handle/123456789/4665/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.