



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5160>


Title:	Synthesis, in vitro anti-plasmodial potency, in-silico-cum-SPR binding with inhibition of PfPyridoxal synthase and rapid parasitocidal action by 3,5-bis(E) arylidene-N-methyl-4-piperidones
Authors:	Joshi, Mayank (/jspui/browse?type=author&value=Joshi%2C+Mayank) Choudhury, Angshuman Roy (/jspui/browse?type=author&value=Choudhury%2C+Angshuman+Roy)
Keywords:	Synthesis in vitro anti-plasmodial potency PfPyridoxal
Issue Date:	2021
Publisher:	New Journal Chemistry
Citation:	New Journal of Chemistry, 45(47), 22150–22165.
Abstract:	Twenty-five (Ia–Iu, IIa–IIb, IIIa, and IVa) diarylidene-N-methyl-4-piperidones (DANMPs) were synthesized and characterized via UV, FT-IR, NMR, and MS while Id was characterized also by single crystal XRD. Twenty-one compounds shortlisted after initial in vitro anti-plasmodial activity successive screenings at 100 μ M and 10 μ M were evaluated for their IC ₅₀ s against chloroquine-sensitive Pf3D7, chloroquine-resistant PfINDO, and artemisinin-resistant PfMRA-1240 strains. The four most promising compounds were Ie (IC ₅₀ s μ M 0.35MRA, 1.39INDO, 1.923D7), If (IC ₅₀ s μ M 1.07MRA, 1.36INDO, 3.393D7), Ir (IC ₅₀ s μ M 0.74MRA, 2.45INDO, 1.443D7), and In (IC ₅₀ s μ M 1.27MRA, 1.8INDO, 1.73D7). Resistance indices as low as 0.2 to 0.5 for these potent compounds and <1 for most other compounds suggest their greater potency against drug resistant strains than the drug sensitive strain. The parasitocidal action of Ir was seen within 4 h against the trophozoite stage of the parasite, which is known to express the highest levels of PLP synthase. In silico docking scores of –7.0 to –8.0 kcal mol ^{–1} between potent DANMPs and PfPLP synthase, the direct binding of Ir studied by SPR to recombinantly expressed and purified PfPdx-1 and inhibition of Pdx1 enzymatic activity by Ir suggest this vital enzyme to be a probable target for the DANMPs. The non-hemolytic nature of Ir and conformity of most DANMPs to Lipinski's parameters indicate their potential as new anti-plasmodial leads with PfPLP synthase as one of their targets.
Description:	Only IISER Mohali authors are available in the record.
URI:	https://doi.org/10.1039/d1nj04604g (https://doi.org/10.1039/d1nj04604g) http://hdl.handle.net/123456789/5160 (http://hdl.handle.net/123456789/5160)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format

Need To Add...Full Text_PDF. (/jspui/bitstream/123456789/5160/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF.)	Only IISER Mohali authors are available in the record.	15.36 kB	Unknown	View/Open (/jspui/
--	--	-------------	---------	------------------------------------

[Show full item record \(/jspui/handle/123456789/5160?mode=full\)](#)

 [\(/jspui/handle/123456789/5160/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.