



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2713>


Title:	Vibrio cholerae porin ompu induces caspase-independent programmed cell death upon translocation to the host cell mitochondria
Authors:	Gupta, Shelly (/jspui/browse?type=author&value=Gupta%2C+Shelly) Prasad, G.V.R.Krishna (/jspui/browse?type=author&value=Prasad%2C+G.V.R.Krishna) Mukhopadhyaya, Arunika (/jspui/browse?type=author&value=Mukhopadhyaya%2C+Arunika)
Keywords:	Vibrio cholerae porin cell death cell mitochondria
Issue Date:	2015
Publisher:	American Society for Biochemistry and Molecular Biology Inc.
Citation:	Journal of Biological Chemistry, 290(52)
Abstract:	Porins, a major class of outer membrane proteins in Gramnegative bacteria, primarily act as transport channels. OmpU is one of the major porins of human pathogen, Vibrio cholerae. In the present study, we show that V. cholerae OmpU has the ability to induce target cell death. Although OmpU-mediated cell death shows some characteristics of apoptosis, such as flipping of phosphatidylserine in the membrane as well as cell size shrinkage and increased cell granularity, it does not show the caspase-3 activation and DNA laddering pattern typical of apoptotic cells. Increased release of lactate dehydrogenase in OmpU-treated cells indicates that the OmpU-mediated cell death also has characteristics of necrosis. Further, we show that the mechanism of OmpU-mediated cell death involves major mitochondrial changes in the target cells. We observe that OmpU treatment leads to the disruption of mitochondrial membrane potential, resulting in the release of cytochrome c and apoptosis-inducing factor (AIF). AIF translocates to the host cell nucleus, implying that it has a crucial role in OmpU-mediated cell death. Finally, we observe that OmpU translocates to the target cell mitochondria, where it directly initiates mitochondrial changes leading to mitochondrial membrane permeability transition and AIF release. Partial blocking of AIF release by cyclosporine A in OmpU-treated cells further suggests that OmpU may be inducing the opening of the mitochondrial permeability transition pore. All of these results lead us to the conclusion that OmpU induces cell death in target cells in a programmed manner in which mitochondria play a central role. © 2015 by The American Society for Biochemistry and Molecular Biology, Inc.
URI:	<a href="https://www.jbc.org/content/290/52/31051">https://www.jbc.org/content/290/52/31051</a> ( <a href="https://www.jbc.org/content/290/52/31051">https://www.jbc.org/content/290/52/31051</a> ) <a href="http://hdl.handle.net/123456789/2713">http://hdl.handle.net/123456789/2713</a> ( <a href="http://hdl.handle.net/123456789/2713">http://hdl.handle.net/123456789/2713</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2713/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2713/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/2713?mode=full>)

 (</jspui/handle/123456789/2713/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.