



## Library Indian Institute of Science Education and Research Mohali



## DSpace@IISERMohali / Thesis & Dissertation / Master of Science / MS-17

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/4248

Title: Study of self - assembly and photoisomerization of polymeric amphiphiles at liquid crystal- aqueous interfaces for the applications in biosensing

Authors: Aggarwal, Aditi

Keywords: photoisomerization

polymeric amphiphiles liquid crystal

interfaces

Issue

Apr-2022

Date:

IISER Mohali

Publisher:
Abstract:

It has been established from past studies that LC-aqueous interfaces are quite responsive to a variety of biological phenomena. It is detected by observing the change in the surface alignment of the liquid crystal molecules using the polarized optical microscope. Firstly, the amphiphilic polymers were studied where it was established that amphiphilic polymers show interactions with the liquid crystal molecules. But there still existed a need to develop a promising approach to introduce specific biorecognition groups at these interfaces that show interactions with the specific biological species. Therefore, new biotin functionality was introduced to these amphiphilic polymers which are known to show specific binding interactions with the avidin protein. We have prepared the doped mixtures of the biotinylated polymer into parent polymer so as to obtain an optimum concentration for further optical studies. Consequently, we were able to detect the presence of avidin protein at the LC-aqueous interfaces by observing the optical signals. Also, by using LC-sensing as a platform, we were able to show that polymer architecture plays an important role in influencing the surface anchoring of LC molecules. Hence two different architectures of polymers (linear and hyperbranched) were differentiated. In addition to this, the effect of photo-induced isomerization of amphiphilic polymers on the ordering of LC within the polymer doped LC films at aqueous interfaces was investigated. Overall, the results of the experiments provide a new approach for the introduction of biorecognition groups at the LC-aqueous interfaces for studying their interactions with biological species including protein.

URI: http://hdl.handle.net/123456789/4248

Appears in MS-17

Collections:

Files in This Item:

File	Description	Size	Format	
Yet to obtain consent.pdf		144.56 kB	Adobe PDF	View/Open

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.



Customized & Implemented by - Jivesna Tech