



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/3129>


Title:	A Trinuclear Zinc–Schiff Base Complex: Biocatalytic Activity and Cytotoxicity
Authors:	Kaur, Gurpreet (/jspui/browse?type=author&value=Kaur%2C+Gurpreet) Choudhury, A.R. (/jspui/browse?type=author&value=Choudhury%2C+A.R.)
Keywords:	Zinc Schiff bases Catecholase activity Drug delivery Antitumor agents Medicinal chemistry
Issue Date:	2014
Publisher:	Wiley-VCH Verlag
Citation:	European Journal of Inorganic Chemistry, 2014(21), pp.3350-3358.
Abstract:	<p>A novel trinuclear zinc(II) complex $[\text{Zn}_3\text{L}_2(\mu\text{-O}_2\text{CCH}_3)_2(\text{CH}_3\text{OH})_4]$ (1) that contains an N,O-donor Schiff base ligand $\{\text{H}_2\text{L} = 2\text{-}[(2\text{-hydroxyphenylimino)methyl]}\text{-6-methoxyphenol}\}$ has been synthesized and crystallographically characterized. The X-ray crystal structure of 1 contains three zinc(II) centers, which have distorted-octahedral coordination geometry, and the molecule crystallizes in the Pbcn space group. The zinc(II) complex displays significant catecholase oxidation activity in methanolic medium through a ligand-centered radical pathway. This is the first example of catecholase oxidation through a trinuclear zinc(II)–Schiff base complex by means of the formation of a mononuclear intermediate as $[\text{ZnL}(\text{dtbc})]$ (dtbc = 3,5-di-tert-butylcatechol). The fluorescence property of 1 indicates that it can serve as a potential photoactive material. It effectively cleaves the double strand of pBR 322 plasmid DNA at a given concentration (25 μM). The complex shows remarkable cytotoxicity against a human hepatocarcinoma cell line (HepG2). A crystallographically characterized trinuclear zinc(II) complex $[\text{Zn}_3\text{L}_2(\mu\text{-O}_2\text{CCH}_3)_2(\text{CH}_3\text{OH})_4]$ (1) that contains an N,O-donor Schiff base ligand $\{\text{H}_2\text{L} = 2\text{-}[(2\text{-hydroxyphenylimino)methyl]}\text{-6-methoxyphenol}\}$ exhibits potential ligand-centered catalytic activity relevant to catechol oxidase. The molecule shows remarkable cytotoxicity against a human hepatocarcinoma cell line (HepG2).</p>
Description:	Only IISERM authors are available in the record.
URI:	https://chemistry-europe.onlinelibrary.wiley.com/doi/abs/10.1002/ejic.201402158 (https://chemistry-europe.onlinelibrary.wiley.com/doi/abs/10.1002/ejic.201402158) http://hdl.handle.net/123456789/3129 (http://hdl.handle.net/123456789/3129)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/3129/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/3129/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/3129?mode=full>)

 (</jspui/handle/123456789/3129/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.