



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Thesis & Dissertation (/jspui/handle/123456789/1)

/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)

/ PhD-2012 (/jspui/handle/123456789/955)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/972>

Title: Crystal Growth and Magnetic Property Study of layered honeycomb materials with Kitaev-like interactions

Authors: Mehlawat, K. (/jspui/browse?type=author&value=Mehlawat%2C+K.)

Keywords: honeycomb materials
quantum magnets
thermodynamic
electrical transport
Crystal growth

Issue Date: 1-Sep-2018

Publisher: IISERM

Abstract: Kitaev-like bond-directional exchange interactions are novel and quite different from the ubiquitous Heisenberg interactions found in most magnets. These kind of interactions open up new possibilities in the exploration and designing of new quantum magnets which can host novel quantum ground states like spin-liquids. However, Kitaev-like interactions have remained only a theoretical construct until recently. Honeycomb iridates $A_2\text{IrO}_3$ ($A = \text{Na}, \text{Li}$), offer potential realizations of such novel exchange coupling. In this thesis I have synthesized single crystals of $A_2\text{TO}_3$ ($A = \text{Na}, \text{Li}, \text{K}$, and $T = \text{Ir}, \text{Ru}$) and studied their electrical transport, magnetic, and thermal properties. Our work provides several new results: (i) the first thermodynamic evidence of possible fractionalization of electrons in Na_2IrO_3 because of proximity to the Kitaev spin-liquid state, (ii) evidence through magnetic impurity doping, of fragile magnetic order and importance of nearest-neighbour interactions and spin-orbit coupling in deciding the magnetic ground state in Na_2IrO_3 , (iii) a novel method (reactive ion etching) of surface doping Na_2IrO_3 and possibly other layered oxides has been discovered. The surface conductivity of Na_2IrO_3 crystals could be increased by 11 orders of magnitude by varying etching times. The samples which turned metallic show transport anomalies consistent with charge density wave or structural instabilities, (iv) first crystal growth of Li_2RuO_3 is reported where the Xtals crystallize in the $P2_1/m$ structure and show the expected high temperature magneto-structural transition, and (v) design and crystal growth of a new layered honeycomb lattice iridate K_2IrO_3 with an interlayer separation between Ir honeycomb planes which is more than a factor of 2 larger than in Na_2IrO_3 . Magnetic measurements on crystals reveal localized effective spin $S = 1/2$ interacting strongly at $\theta = -210 \text{ K}$ but without magnetic order down to 1.8 K . Thus K_2IrO_3 is a new Kitaev spin liquid candidate

URI: <http://hdl.handle.net/123456789/972> (<http://hdl.handle.net/123456789/972>)

Appears in PhD-2012 (/jspui/handle/123456789/955)
Collections:

Files in This Item:

File	Description	Size	Format	
PH12136.pdf (/jspui/bitstream/123456789/972/1/PH12136.pdf)		22.87 MB	Adobe PDF	View/Open (/jspui/bitstream/123456789/972/1/F

[Show full item record \(/jspui/handle/123456789/972?mode=full\)](/jspui/handle/123456789/972?mode=full)

[📊 \(/jspui/handle/123456789/972/statistics\)](/jspui/handle/123456789/972/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

[Export \(migrate\) Item](#)

[Export metadata](#)