



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4802>

Title:	Glu289 residue in the pore-forming motif of <i>Vibrio cholerae</i> cytolysin is important for efficient β -barrel pore formation
Authors:	Mondal, Anish Kumar (/jspui/browse?type=author&value=Mondal%2C+Anish+Kumar) Singh, Mahendra (/jspui/browse?type=author&value=Singh%2C+Mahendra) Lata, Kusum (/jspui/browse?type=author&value=Lata%2C+Kusum) Lahiri, Indrajit (/jspui/browse?type=author&value=Lahiri%2C+Indrajit) Chattopadhyay, Kausik (/jspui/browse?type=author&value=Chattopadhyay%2C+Kausik)
Keywords:	Glu289 residue in the pore-forming motif of <i>Vibrio cholerae</i> cytolysin β -barrel pore formation protein structure
Issue Date:	2022
Publisher:	Elsevier
Citation:	Journal of Biological Chemistry, 298(10), 102441.
Abstract:	<p><i>Vibrio cholerae</i> cytolysin (VCC) is a potent membrane-damaging β-barrel pore-forming toxin. Upon binding to the target membranes, VCC monomers first assemble into oligomeric prepore intermediates and subsequently transform into transmembrane β-barrel pores. VCC harbors a designated pore-forming motif, which, during oligomeric pore formation, inserts into the membrane and generates a transmembrane β-barrel scaffold. It remains an enigma how the molecular architecture of the pore-forming motif regulates the VCC pore-formation mechanism. Here, we show that a specific pore-forming motif residue, E289, plays crucial regulatory roles in the pore-formation mechanism of VCC. We find that the mutation of E289A drastically compromises pore-forming activity, without affecting the structural integrity and membrane-binding potential of the toxin monomers. Although our single-particle cryo-EM analysis reveals WT-like oligomeric β-barrel pore formation by E289A-VCC in the membrane, we demonstrate that the mutant shows severely delayed kinetics in terms of pore-forming ability that can be rescued with elevated temperature conditions. We find that the pore-formation efficacy of E289A-VCC appears to be more profoundly dependent on temperature than that of the WT toxin. Our results suggest that the E289A mutation traps membrane-bound toxin molecules in the prepore-like intermediate state that is hindered from converting into the functional β-barrel pores by a large energy barrier, thus highlighting the importance of this residue for the pore-formation mechanism of VCC.</p>
Description:	Only IISERM authors are available in the record
URI:	https://doi.org/10.1016/j.jbc.2022.102441 (https://doi.org/10.1016/j.jbc.2022.102441) http://hdl.handle.net/123456789/4802 (http://hdl.handle.net/123456789/4802)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:


File	Description	Size	Format

Need To Add...Full Text_PDF
(/jspui/bitstream/123456789/4802/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF)

15.36 Unknown
kB

[View/Open \(/jspui/t](#)

[Show full item record \(/jspui/handle/123456789/4802?mode=full\)](#)

 [\(/jspui/handle/123456789/4802/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.