

Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/3230

Title: Allosteric inhibition of MTHFR prevents futile SAM cycling and maintains nucleotide pools in one-

carbon metabolism

Authors: Bhatia, Muskan (/jspui/browse?type=author&value=Bhatia%2C+Muskan)

Suyal, S. (/jspui/browse?type=author&value=Suyal%2C+S.)

Sharma, Mahak (/jspui/browse?type=author&value=Sharma%2C+Mahak) Bachhawat, A.K. (/jspui/browse?type=author&value=Bachhawat%2C+A.K.)

Keywords: Allosteric inhibition

MTHFR SAM Nucleotide

Issue Date: 2020

Publisher: American Society for Biochemistry and Molecular Biology Inc.

Citation: Journal of Biological Chemistry, 295(47) pp. 16037-16057.

Abstract:

Methylenetetrahydrofolate reductase (MTHFR) links the folate cycle to the methionine cycle in one-carbon metabolism. The enzyme is known to be allosterically inhibited by SAM for decades, but the importance of this regulatory control to one-carbon metabolism has never been adequately understood. To shed light on this issue, we exchanged selected amino acid residues in a highly conserved stretch within the regulatory region of yeast MTHFR to create a series of feedbackinsensitive, deregulated mutants. These were exploited to investigate the impact of defective allosteric regulation on one-carbon metabolism. We observed a strong growth defect in the presence of methionine. Biochemical and metabolite analysis revealed that both the folate and methionine cycles were affected in these mutants, as was the transsulfuration pathway, leading also to a disruption in redox homeostasis. The major consequences, however, appeared to be in the depletion of nucleotides. 13C isotope labeling and metabolic studies revealed that the deregulated MTHFR cells undergo continuous transmethylation of homocysteine by methyltetrahydrofolate (CH3THF) to form methionine. This reaction also drives SAM formation and further depletes ATP reserves. SAM was then cycled back to methionine, leading to futile cycles of SAM synthesis and recycling and explaining the necessity for MTHFR to be regulated by SAM. The study has yielded valuable new insights into the regulation of one-carbon metabolism, and the mutants appear as powerful new tools to further dissect out the intersection of one-carbon metabolism with various pathways both in yeasts and in humans

Description: Only IISERM authors are available in the record.

URI: https://www.sciencedirect.com/science/article/pii/S0021925817504305?via%3Dihub (https://www.sciencedirect.com/science/article/pii/S0021925817504305?via%3Dihub)

http://hdl.handle.net/123456789/3230 (http://hdl.handle.net/123456789/3230)

Appears in Collections:

Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File Description Size Format

Need to add pdf.odt (/jspui/bitstream/123456789/3230/1/Need%20to%20add%20pdf.odt)

8.63 OpenDocument kB Text

View/Open (/jspui/bitstream/12345

Show full item record (/jspui/handle/123456789/3230?mode=full)

. I (/jspui/handle/123456789/3230/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.