

## Library Indian Institute of Science Education and Research Mohali



## DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/108

Title: Structural studies of diorganotin(IV) sulfonates: The synthesis of [(n-C4H 9)2Sn{OSO2C 6H 3(CH

3)2-2,5}2] and [(n-C 4H 9)2Sn{OSO 2R)2  $\cdot$  2(hexamethylphosphoric triamide)] [R=CH3, 4-C6H4CH3, 2,5-C6 H3(CH3)2] 2,5,6-C6H2(CH3)3] and crystal structures of [(n-C4H9)2Sn( $\mu$ -OH) (OSO2C6H3(CH3)2-2,5)]2 and (n-C4H9)2Sn{OSO2R}2 $\cdot$  2(hexamethylphosphoric triamide)] [R=4-

C6H4CH3, 2,5-C6H3(CH3)2, 2,4,6-C6H2(CH3)3]

Authors: Kapoor, Ramesh (/jspui/browse?type=author&value=Kapoor%2C+Ramesh)

Keywords: Atomic physics

Atoms Complexation

Computational geometry

Issue Date: 2009

Publisher: Elsevier B.V.

Citation: Journal of Organometallic Chemistry, 694 (5), pp. 623-629.

 $Abstract: \qquad \mbox{The synthesis of [(n-C 4H 9)2Sn{OSO2C6H3(CH3)2-2,5} 2] and [(n-C 4H 9)2Sn{OSO2R)2} + (n-C 4H 9)2Sn{OSO2R} +$ 

2(hexamethylphosphoric triamide)] [R = CH3 (3), 4-C6H4CH 3, (4) 2,5-C6H3(CH3) 2 (5), 2,4,6-C6H2(CH3) 3 (6) and have been carried out to study their structures and to delineate the coordination behavior of the weakly coordinating sulfonate anions. Compound 1 hydrolyzes slowly to [(n-C4H9) 2Sn( $\mu$ -OH)(OSO2C6H3(CH3)2-2,5)]2 (2) when kept in CH2Cl2 for a few days. The crystal structure shows that 2 has a dimeric structure in which tin atoms are bridged by two hydroxy groups and each tin atom is further bonded to two n-Bu groups and a mono-coordinated sulfonate anion thus generating a trigonal bipyramidal geometry at tin atom. However, the coordination geometry at tin can be visualized as a severely distorted octahedron, if a relatively weak Sn-O bond of 2.690 Å between Sn and an O atom of the neighboring sulfonate anion is also considered. Compounds 3-6 are obtained by the consecutive reaction of (n-C4H9)2SnO with the appropriate sulphonic acid and hexamethylphosphoric triamide (HMPA). X-ray crystal structures of 4, 5 and 6 show octahedral geometry around tin atom in the two compounds in which the sulfonate ligands are covalently bonded in a monodentate mode. Compounds 3-6 are nonionic in polar solvents. The Sn-O (sulfonate) bond distances 2.354(2), 2; 2.233(2), 4; 2.237(5), 5 and 2.227(3) Å, 6 suggest some degree of ionic character in the metal-anion bonds. These compounds have also been characterized by multinuclear (1H,13C and119Sn) NMR studies.

Description: Only IISERM authors are available in the record.

URI: http://www.sciencedirect.com/science/article/pii/S0022328X0800764X

(http://www.sciencedirect.com/science/article/pii/S0022328X0800764X) http://dx.doi.org/10.1016/j.jorganchem.2008.11.031

(http://dx.doi.org/10.1016/j.jorganchem.2008.11.031)

Appears in I Collections:

Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File Description Size Format

Need to add pdf.odt (/jspui/bitstream/123456789/108/3/Need%20to%20add%20pdf.odt)

8.63 OpenDocument kB Text

View/Open (/jspui/bitstream/123456

Show full item record (/jspui/handle/123456789/108?mode=full)

d (/jspui/handle/123456789/108/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.