



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2997>


Title:	Transmembrane oligomeric form of <i>Vibrio cholerae</i> cytolysin triggers TLR2/TLR6-dependent proinflammatory responses in monocytes and macrophages
Authors:	Khilwani, B. (/jspui/browse?type=author&value=Khilwani%2C+B.) Mukhopadhyaya, Arunika (/jspui/browse?type=author&value=Mukhopadhyaya%2C+Arunka) Chattopadhyay, K. (/jspui/browse?type=author&value=Chattopadhyay%2C+K.)
Keywords:	Bacterial toxins Macrophages Monocytes Proinflammatory response Toll-like receptors <i>Vibrio cholerae</i> cytolysi
Issue Date:	2015
Publisher:	Biochemical Society
Citation:	Biochemical Journal, 466 pp. 147-161
Abstract:	<p><i>Vibrio cholerae</i> cytolysin (VCC) kills target eukaryotic cells by forming transmembrane oligomeric β-barrel pores. Once irreversibly converted into the transmembrane oligomeric form, VCC acquires an unusual structural stability and loses its cytotoxic property. It is therefore possible that, on exertion of its cytotoxic activity, the oligomeric form of VCC retained in the disintegrated membrane fractions of the lysed cells would survive within the host cellular milieu for a long period, without causing any further cytotoxicity. Under such circumstances, VCC oligomers may potentially be recognized by the host immune cells. Based on such a hypothesis, in the present study we explored the interaction of the transmembrane oligomeric form of VCC with the monocytes and macrophages of the innate immune system. Our study shows that the VCC oligomers assembled in the liposome membranes elicit potent proinflammatory responses in monocytes and macrophages, via stimulation of the tolllike receptor (TLR)2/TLR6-dependent signalling cascades that involve myeloid differentiation factor 88 (MyD88)/interleukin-1-receptor-associated kinase (IRAK)1/tumour-necrosis-factor-receptor-associated factor (TRAF)6. VCC oligomer-mediated proinflammatory responses critically depend on the activation of the transcription factor nuclear factor-κ B. Proinflammatory responses induced by the VCC oligomers also require activation of the mitogen-activated protein kinase (MAPK) family member c-Jun N-terminal kinase, which presumably acts via stimulation of the transcription factor activator protein-1. Notably, the role of the MAPK p38 could not be documented in the proces</p>
URI:	https://pubmed.ncbi.nlm.nih.gov/25431887/ (https://pubmed.ncbi.nlm.nih.gov/25431887/) http://hdl.handle.net/123456789/2997 (http://hdl.handle.net/123456789/2997)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

Need to add pdf.odt (/jspui/bitstream/123456789/2997/1/Need%20to%20add%20pdf.odt)	8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/2997/1/Need%20to%20add%20pdf.odt)
--	------------	----------------------	---

Show full item record (/jspui/handle/123456789/2997?mode=full)

 (/jspui/handle/123456789/2997/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.