



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2887>


Title:	Pd(OAc) ₂ /AgOAc catalytic system based bidentate ligand directed regiocontrolled C-H arylation and alkylation of the C-3 position of thiophene- and furan-2-carboxamides
Authors:	Padmavathi, R. (/jspui/browse?type=author&value=Padmavathi%2C+R.) Sankar, Rathinam (/jspui/browse?type=author&value=Sankar%2C+Rathinam) Gopalakrishnan, B. (/jspui/browse?type=author&value=Gopalakrishnan%2C+B.) Parella, R. (/jspui/browse?type=author&value=Parella%2C+R.) Babu, S.A. (/jspui/browse?type=author&value=Babu%2C+S.A.)
Keywords:	Pd(OAc) ₂ /AgOAc bidentate ligand furan-2-carboxamides
Issue Date:	2015
Publisher:	WILEY-VCH Verlag GmbH
Citation:	European Journal of Organic Chemistry, 2015 (17) pp. 3727-3742
Abstract:	A contemporary method is reported for the Pd(OAc) ₂ /AgOAc catalytic system based bidentate ligand directed, regioselective C-H activation and C-C bond formation at the C-3 position of thiophene- and furan-2-carboxamides, which are derived from 8-aminoquinoline or 2-(methylthio)aniline. The Pd-catalyzed C-H arylation of thiophene- and furan-2-carboxamides with a variety of aryl iodides and heteroaryl iodides was highly regioselective and afforded C-3-arylated thiophene-2-carboxamides and furan-2-carboxamides in good to very good yields. The bidentate ligand directed Pd(OAc) ₂ /AgOAc based strategy was also successfully employed for benzylation and alkylation reactions of the thiophene-2-carboxamides. These reactions occurred with high regioselectivity to afford the C-3-benzylated and C-3-alkylated thiophene-2-carboxamides in good yields. The observed regioselectivity of these reactions was confirmed by X-ray crystal structure analyses of compounds 3e, 5a, and 6a. A contemporary method is reported for the Pd(OAc) ₂ /AgOAc catalytic system based bidentate ligand directed, regioselective C-H activation and C-C bond formation of the C-3 position of thiophene- and furan-2-carboxamides. This protocol was used for the direct C-3 arylation and alkylation reactions of both thiophene- and furan-2-carboxamides. Copyright
URI:	https://chemistry-europe.onlinelibrary.wiley.com/doi/full/10.1002/ejoc.201500249 (https://chemistry-europe.onlinelibrary.wiley.com/doi/full/10.1002/ejoc.201500249) http://hdl.handle.net/123456789/2887 (http://hdl.handle.net/123456789/2887)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2887/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2887/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/2887?mode=full>)

 (</jspui/handle/123456789/2887/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.