



## Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**  
**/ Thesis & Dissertation (/jspui/handle/123456789/1)**  
**/ Master of Science (/jspui/handle/123456789/2)**  
**/ MS-14 (/jspui/handle/123456789/1078)**


Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1125>

Title:	Conformational Analysis of 2-Butyne-1, 4-diol: A Matrix Isolation FTIR and Ab-Initio Study
Authors:	Singh, Himanshi (/jspui/browse?type=author&value=Singh%2C+Himanshi)
Keywords:	Chemical Sciences Matrix Isolation FTIR Hydrogen Bonding Crystallography Nature of Matrix
Issue Date:	23-Sep-2019
Publisher:	IISERM
Abstract:	<p>The work presents, for the first time, the matrix isolation infrared and ab initio studies on the molecular conformations of 2-butyne-1, 4-diol. It is found that when the acetylenic moiety (<math>\text{-C}\equiv\text{C-H}</math>) in propargyl alcohol is replaced by a <math>\text{CH}_2\text{OH}</math> group, it gives rise to many different conformations of ethylene glycol, which are facilitated by strong intramolecular O-H ... O hydrogen bonded interactions. Similarly, when the acidic hydrogen in propargyl alcohol is replaced by a <math>\text{CH}_2\text{OH}</math> group, several conformations arises for 2-butyne-1, 4-diol. Thus, these molecules, propargyl alcohol, ethylene glycol and 2- butyne-1,4-diol provide an interesting case to study the effect of an acetylenic <math>\pi</math> cloud between the two <math>\text{-OH}</math> groups on the conformations, which was absent in ethylene glycol. Starting from only two stable conformers in propargyl alcohol and ten stable conformers in ethylene glycol, we have computed six stable conformers of 2-butyne-1,4-diol in this work at M06-2X and MP2 methods employing 6-311++G** and aug-cc-pVDZ basis sets. Our experiments together with computations revealed the presence of three most stable conformer in the N<sub>2</sub> matrix. AIM and NBO analysis have been performed to understand the nature of interactions in the conformers. Vicinal orbital interactions are found to play deciding role in its conformational preference. Competitive hydrogen bonding because of the presence of multiple bonding sites in 2-butyne-1, 4-diol is another aspect which we have explored from our computational studies on its hydrogen-bonded complexes with water.</p>
URI:	IISER M (IISER M) <a href="http://hdl.handle.net/123456789/1125">http://hdl.handle.net/123456789/1125</a> ( <a href="http://hdl.handle.net/123456789/1125">http://hdl.handle.net/123456789/1125</a> )
Appears in Collections:	MS-14 (/jspui/handle/123456789/1078)

Files in This Item:

File	Description	Size	Format	
MS14084.pdf (/jspui/bitstream/123456789/1125/3/MS14084.pdf)	Full Text.pdf	2.47 MB	Adobe PDF	<a href="#">View/Open (/jspui/bitstream/123456789/1125/3/I</a>

[Show full item record \(/jspui/handle/123456789/1125?mode=full\)](#)

 [\(/jspui/handle/123456789/1125/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.