



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1795>


Title:	A critical role for sorting nexin 1 in the trafficking of metabotropic glutamate receptors
Authors:	Sharma, Rohan (/jspui/browse?type=author&value=Sharma%2C+Rohan) Gulia, R. (/jspui/browse?type=author&value=Gulia%2C+R.) Bhattacharyya, Samarjit (/jspui/browse?type=author&value=Bhattacharyya%2C+Samarjit)
Keywords:	Metabotropic glutamate receptors Neurotransmitter receptors Trafficking Cell Culture Techniques
Issue Date:	2018
Publisher:	Society for Neuroscience
Citation:	Journal of Neuroscience, 38(40) pp. 8605-8620
Abstract:	Group I metabotropic glutamate receptors (mGluRs) function as modulators of neuronal physiology and they have also been implicated in various neuropsychiatric disorders. Trafficking of mGluRs plays important roles in controlling the precise localization of these receptors at specific region of the cell, as well as it regulates the activity of these receptors. Despite this obvious significance, we know very little about the cellular machineries that control the trafficking of these receptors in the CNS. Sorting nexin 1 (SNX1) has been shown to regulate the endosomal sorting of few cell surface receptors either to lysosomes where they are downregulated or back to the cell surface. Using "molecular replacement" approach in hippocampal neurons derived from mice of both sexes, we show here that SNX1 plays critical role in the trafficking of mGluR1, a member of the group I mGluR family. Overexpression of dominant-negative SNX1 or knockdown of endogenous SNX1 resulted in the rapid recycling of the receptor. Importantly, recycling via the rapid recycling route, did not allow the resensitization of the receptors. Our data suggest that both, N-terminal and C-terminal region of SNX1 play critical role in the normal trafficking of the receptor. In addition, we also show here that SNX1 regulates the trafficking of mGluR1 through the interaction with Hrs (hepatocyte growth factor-regulated tyrosine kinase substrate), a protein that has been implicated in both signaling and vesicular trafficking. Thus, these studies reveal a mechanistic role of SNX1 in the trafficking of group I mGluRs and its physiological implications.
URI:	https://www.jneurosci.org/content/early/2018/08/24/JNEUROSCI.0454-18.2018/tab-article-info?versioned=true (https://www.jneurosci.org/content/early/2018/08/24/JNEUROSCI.0454-18.2018/tab-article-info?versioned=true) http://hdl.handle.net/123456789/1795 (http://hdl.handle.net/123456789/1795)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/1795/1/Need%20to%20add%20pdf.odt)		8.04 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/1795/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/1795?mode=full>)

 (</jspui/handle/123456789/1795/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.