



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4912>


Title:	Self-Assembled Oligothiophenes for Photocatalytic Hydrogen Production and Simultaneous Organic Transformation
Authors:	Mishra, Samita (/jspui/browse?type=author&value=Mishra%2C+Samita) Kumar, Amit (/jspui/browse?type=author&value=Kumar%2C+Amit) De, Arijit K. (/jspui/browse?type=author&value=De%2C+Arijit+K.)
Keywords:	Self-Assembled Oligothiophenes Photocatalytic Hydrogen Production and Simultaneous Organic Transformation
Issue Date:	2022
Publisher:	ACS Publications
Citation:	ACS Applied Nano Materials, 5(10), 14746-14758.
Abstract:	Herein, we have fabricated self-assembled semiconducting organic nanomaterials with various morphologies (1D-fiber, 2D-flakes, and 2D-nanosheets) made of small conjugated oligomer 2,2':5',2'':5'',2'''-quaterthiophene (α -QTH) by a simple solution-based coprecipitation method. By simply varying the good-solvent-to-bad-solvent ratio, we can critically tune the self-assembly process and eventually can control the intermolecular interactions of the constituent molecules in these self-assembled nanostructures. Different types of self-assembled nanostructures have been utilized for photocatalytic solar H ₂ production. The H ₂ production efficiencies directly depend on the morphology of self-assembled self-assembled nanomaterials as well as intermolecular interactions of QTH molecules. The overall photocatalytic properties are further correlated with the ongoing photophysical properties by means of detailed steady-state and time-resolved fluorescence spectroscopy and dimer-based time dependent-density functional theory (TD-DFT) calculations. Furthermore, femtosecond transient absorption spectroscopy has been utilized to explore the detailed photoinduced exciton dynamics by global analysis of spectrally resolved pump-probe traces. In addition to that, the overall photocatalytic activities are further supported by an in-depth electrochemical study. Finally, a boost in photocatalytic H ₂ production has been observed by using 4-methylbenzyl alcohol (4-MBA) as a specific hole scavenger for the completion of the redox cycle. Therefore, the present system can be utilized for simultaneous solar H ₂ production and the specific organic transformation through a green and cost-efficient approach.
Description:	Only IISER Mohali authors are available in the record.
URI:	https://doi.org/10.1021/acsanm.2c03061 (https://doi.org/10.1021/acsanm.2c03061) http://hdl.handle.net/123456789/4912 (http://hdl.handle.net/123456789/4912)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF (1) (/jspui/bitstream/123456789/4912/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF%20%281%29)		15.36 kB	Unknown

[View](#)

Show full item record (</jspui/handle/123456789/4912?mode=full>)

 (</jspui/handle/123456789/4912/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.