



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)


Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2714>

Title:	Hydrogen-Bonded Complexes of Phenylacetylene-Acetylene: Who is the Proton Donor
Authors:	Verma, Kanupriya (/jspui/browse?type=author&value=Verma%2C+Kanupriya) Dave, Kapil (/jspui/browse?type=author&value=Dave%2C+Kapil) Viswanathan, K.S. (/jspui/browse?type=author&value=Viswanathan%2C+K.S.)
Keywords:	oncovalent interactions Hydrocarbons Interaction energies Isolation
Issue Date:	2015
Publisher:	American Chemical Society
Citation:	Journal of Physical Chemistry A, 119(51)
Abstract:	Hydrogen-bonded complexes of C ₂ H ₂ and phenylacetylene (PhAc) were studied using matrix isolation infrared spectroscopy and quantum chemical computations. Both C ₂ H ₂ and PhAc, being potential proton donors, the question arises as to which of the two species would be the proton donor in the PhAc-C ₂ H ₂ complex; a question that this work primarily addresses. The molecular structures, vibrational frequencies, and interaction energies of the PhAc-C ₂ H ₂ complexes were calculated at the M06-2X and MP2 levels of theory, employing both 6-311++G(d,p) and aug-cc-pVDZ basis sets. At the M06-2X/aug-cc-pVDZ level, two nearly isoenergetic complexes (BSSE corrected) were indicated to be the global minima; one a C-H... π complex, where C ₂ H ₂ served as a proton donor to the phenyl π -system in PhAc, and the other a C-H... π complex, where C ₂ H ₂ served as a proton donor to the acetylene π -system in PhAc. Of the two, only the second complex was identified in the matrix, evidenced by a characteristic large shift in the \equiv C-H stretch of C ₂ H ₂ . Experiments were also performed using PhAc deuterated at the acetylene hydrogen (PhAcD) to study the isotopic effects on the vibrational spectra of complexes. The isotopic studies further confirmed the structure of the complex trapped in the matrix, thereby presenting unambiguous evidence that C ₂ H ₂ served as the proton donor to the acetylene π -system of PhAc. The theory of atoms-in-molecules (AIM), energy decomposition (EDA), and natural bond orbital (NBO) analysis were performed to understand the nature of the interactions involved in the complexes.
URI:	https://pubs.acs.org/doi/abs/10.1021/acs.jpca.5b08559?src=recsys (https://pubs.acs.org/doi/abs/10.1021/acs.jpca.5b08559?src=recsys) http://hdl.handle.net/123456789/2714 (http://hdl.handle.net/123456789/2714)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/2714/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/2714/1/Need%20to%20add%20pdf.odt)

Show full item record (</jspui/handle/123456789/2714?mode=full>)

 (</jspui/handle/123456789/2714/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.