



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Master of Science / MS-18

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5577>

Title:	Understanding the role of HY5 in maintaing iron homeostasis in Arabidopsis thaliana
Authors:	Kalra, Muskan
Keywords:	Arabidopsis thaliana iron homeostasis
Issue Date:	May-2023
Publisher:	IISER Mohali
Abstract:	<p>iron is the fourth most abundant element in the Earth's crust and is crucial for plants because of its diverse roles; however, its availability to plants is minimal. Almost 1/3 rd of the world's cultivated lands are calcareous, where iron exists in the form of insol- ule ferric ions, making the soil iron deficient. To cope with this deficiency, plants have evolved two strategies which are Strategy 1: Reduction-based and Strategy II: Chelation- based. However, as plants are the only source of dietary iron at higher trophic levels. improving iron uptake in plants is currently the major area of focus. Arabidopsis thaliana follows a reduction-based strategy for iron uptake from the soil and has quite an elaborate iron signaling pathway which the bHLH family of transcription factors majorly governs. Many of these have already been elucidated; however, there is still much to discover. HY5 encodes for a bZIP transcription factor that is known to be involved in photomorphogenesis and seedling development. It inhibits hypocotyl elongation and lateral root development. It is already known that HYS has a role in sulphur, nitrate and copper signaling pathways. Upon checking the phenotype, we found that HYS has a role in the primary root growth and development of the Arabidopsis thaliana under iron-limiting conditions as the hy5 mutant exhibits shortened primary root length, less chlorophyll content and low iron levels in iron- limiting conditions. This phenotype was rescued fully upon the use of the overexpression line of HY5 constructed in the hy5 mutant background (35S:HA:HY5/hy5). This confirmed the role of HY5 in primary root growth and development under iron-limiting conditions. In a recent study, it was shown that the induction of HY5 for iron uptake was dependent on PHYB in the case of tomato. Upon analysing the phenotype of p / t * yB - 9 mutants, we concluded that under iron-limiting conditions, HYS regulates the primary root length inde- pendent of PHYB in Arabidopsis thaliana.</p>
Description:	Embargo Period
URI:	http://hdl.handle.net/123456789/5577
Appears in Collections:	MS-18

Files in This Item:

File	Description	Size	Format	
Need To Add...Full Text_PDF		15.36 kB	Unknown	View/Open

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.