



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Thesis & Dissertation (/jspui/handle/123456789/1)

/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)

/ PhD-2013 (/jspui/handle/123456789/269)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1571>

Title:	Function and mechanism of the unconventional ubiquitin-like protein Hub1
Authors:	Kumar, Kiran K. (/jspui/browse?type=author&value=Kumar%2C+Kiran+K.)
Keywords:	RNA Splicing Assembly of the spliceosome Molecular biology reagents DNA isolation
Issue Date:	Nov-2019
Publisher:	IISER Mohali
Abstract:	Ubiquitin-like proteins (UBLs) control various cellular processes such as protein degradation, DNA repair, autophagy, transcription, RNA splicing, and immune responses. Hub1 is one such UBLs that have been reported to regulate pre-mRNA splicing. However, the function and the mechanism of Hub1 in intron-rich eukaryotes has not been studied yet. We aimed to understand the role of Hub1 in RNA splicing in an intron-rich unicellular eukaryote <i>Schizosaccharomyces pombe</i> , and a multicellular eukaryote, <i>Caenorhabditis elegans</i> . This study demonstrates a conserved genome-wide role of Hub1 in pre-mRNA splicing in <i>S. pombe</i> . Hub1 alters the protein composition of the spliceosome selectively. It promotes splicing of pre-mRNAs that are synthesized faster. It is likely that rapidly synthesizing transcripts require Hub1 to couple transcription with splicing. We identified a functionally conserved Hub1 surface centered at two positively charged residues critical for splicing in <i>S. pombe</i> . We also show that Hub1 directly binds to the Krebs cycle enzyme, Fum1, through another surface. This Hub1 surface also regulates pre-mRNA splicing. Additionally, our study showed the potential role of Hub1 in trans-splicing in the multicellular <i>C. elegans</i> . In summary, Hub1 employs multiple surfaces to facilitate binding of specific factors for its function in pre-mRNA splicing. My findings are highly relevant not only for regulatory and tissue-specific gene expression, but also for understanding new mechanisms of alternative splicing.
URI:	http://hdl.handle.net/123456789/1571 (http://hdl.handle.net/123456789/1571)
Appears in Collections:	PhD-2013 (/jspui/handle/123456789/269)

Files in This Item:


File	Description	Size	Format

PH13037.pdf
(/jspui/bitstream/123456789/1571/3/PH13037.pdf)

7.16 MB Adobe PDF

[View/Open \(/jspui/bitstream/123456789/1571/3/\)](/jspui/bitstream/123456789/1571/3/)

[Show full item record \(/jspui/handle/123456789/1571?mode=full\)](/jspui/handle/123456789/1571?mode=full)

 [\(/jspui/handle/123456789/1571/statistics\)](/jspui/handle/123456789/1571/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

[Edit...](#)

[Export Item](#)

[Export \(migrate\) Item](#)

[Export metadata](#)