



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4352>


Title:	Residential heating emissions (can) exceed paddy-residue burning emissions in rural northwest India
Authors:	Pawar, Harshita (/jspui/browse?type=author&value=Pawar%2C+Harshita) Sinha, Baerbel (/jspui/browse?type=author&value=Sinha%2C+Baerbel)
Keywords:	Crop residue burning Residential heating Solid biofuel India Particulate matter Low-cost sensors
Issue Date:	2021
Publisher:	Elsevier
Citation:	Atmospheric Environment, 269, 118846.
Abstract:	<p>November onwards, the poor air quality over northwest India is blamed on the large-scale paddy residue burning in Punjab and Haryana. However, the emission strength of this source remains poorly constrained due to the lack of ground-based measurements over rural areas and issues in the satellite detection of paddy residue fires. In this study, we report the particulate matter (PM) measurements at Nadampur, a village in the Sangrur district with the highest reported paddy residue fires, from 1 October to 19 December 2019, using the Airveda low-cost PM sensors. The daily average PM<sub>10</sub> and PM<sub>2.5</sub> mass concentration at Nadampur correlated well with the daily sum of Visible Infrared Imaging Radiometer Suite (VIIRS) fire counts (<math>r &gt; 0.7</math>) in a 50 km × 50 km area surrounding the village. Agreement of the Coefficient of Emissions (Ce) estimated in this study (0.038 kg MJ<sup>-1</sup>) with the reported value (0.04 kg MJ<sup>-1</sup>), and a disagreement of the top-down estimate of PM emission factors with the laboratory reported values indicates an underdetection of paddy residue fires. Residential burning of solid fuels such as cow-dung cakes and fuelwood for space heating triggered by a dip in the temperature led to poor air quality from 20 November onwards. Source apportionment performed using Multiple Linear Regression (MLR) and Positive Matrix Factorization (PMF) revealed that paddy residue burning increased the PM<sub>10</sub> (PM<sub>2.5</sub>) at Nadampur by <math>97.0 \pm 36.6 \mu\text{g m}^{-3}</math> (<math>53.4 \pm 16.8 \mu\text{g m}^{-3}</math>) which was more than harvesting activities <math>44.8 \pm 1.7 \mu\text{g m}^{-3}</math> (<math>20.1 \pm 5.2 \mu\text{g m}^{-3}</math>), but less than residential heating emissions <math>151.2 \pm 47.2 \mu\text{g m}^{-3}</math> (<math>120.1 \pm 8.8 \mu\text{g m}^{-3}</math>). Unlike agricultural activities, which typically affect the air quality for roughly one month, heating-related emissions profoundly impact the air quality for multiple months.</p>
Description:	Only IISER Mohali authors are available in the record.
URI:	<a href="https://doi.org/10.1016/j.atmosenv.2021.118846">https://doi.org/10.1016/j.atmosenv.2021.118846</a> ( <a href="https://doi.org/10.1016/j.atmosenv.2021.118846">https://doi.org/10.1016/j.atmosenv.2021.118846</a> ) <a href="http://hdl.handle.net/123456789/4352">http://hdl.handle.net/123456789/4352</a> ( <a href="http://hdl.handle.net/123456789/4352">http://hdl.handle.net/123456789/4352</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

Need To Add...Full Text_PDF..pdf (/jspui/bitstream/123456789/4352/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF..pdf)	Only IISER Mohali authors are available in the record.	15.36 kB	Adobe PDF	<a href="#">View/Open (/jspu</a>
--	--	-------------	--------------	----------------------------------

[Show full item record \(/jspui/handle/123456789/4352?mode=full\)](#)

 [\(/jspui/handle/123456789/4352/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.