



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Doctor of Philosophy (PhD) / PhD-2015

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4084>

Title:	Multimodal inhibition of the weedy fungi in the gardens of <i>Odontotermes obesus</i>
Authors:	Agarwal, Renuka
Keywords:	Multimodal odontotermes
Issue Date:	Nov-2021
Publisher:	IISER Mohali
Abstract:	<p>Termites feed on different lignocellulosic biomass which are chemically recalcitrant and require a nutritional symbiosis with microbes to digest. Fungus-growing termites (Subfamily Macrotermitinae) have evolved to cultivate a specialized fungus, <i>Termitomyces</i>, as food which is cultivated on a spongy substrate called the fungus comb. But these fungal crops face constant threat of invasion by other weedy fungi, like <i>Pseudoxylaria</i>, which can take over the crop fungus. My thesis tries to answer how these weedy fungi are kept in check by secondary bacterial symbionts found in these colonies and what are the relationships between them and the crop and weedy fungi. Using both culture-dependent isolation of bacterial and fungal strains and culture-independent surveys using the Nanopore platform, I report the presence of several bacterial mutualists which can prevent the weedy fungus while keeping the crop fungus relatively unharmed. A sequential analysis of both the micro- and mycobiota of a degrading fungus combs identified some of the roles of these microbes in maintaining a healthy fungus comb where a coupled dynamics in the fungal and bacterial communities were seen. However, this survey also indicated the near absence of any other fungi in the comb, in spite of it being an ideal growth chamber of various other non-specific fungi. Therefore, I hypothesized that as <i>Termitomyces</i> and <i>Pseudoxylaria</i> are strictly termite associated fungi, they could have roles in controlling the growth of non-specific fungi. Through in vitro interaction assays I found that both <i>Termitomyces</i> and <i>Pseudoxylaria</i> can indeed inhibit many of these fungal contaminants. However, as <i>Pseudoxylaria</i> inhibits <i>Termitomyces</i>, I introduce a verbal model to explain the dynamics of the symbiosis. This model postulates that the fungal contaminants are prevented by <i>Termitomyces</i>, <i>Pseudoxylaria</i> as well as some bacterial mutualists. <i>Pseudoxylaria</i>, in its turn, is prevented by some of the bacterial mutualists as well as by the termites.</p>
URI:	http://hdl.handle.net/123456789/4084
Appears in Collections:	PhD-2015

Files in This Item:

File	Description	Size	Format	
It is under embargo period.pdf		139.68 kB	Adobe PDF	View/Open

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

Edit...

Export Item

Export (migrate) Item

Export metadata