



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/3501>


Title:	Synthesis, spectroscopic and structural characterization of Co(II), Ni(II) and Cu(II) complexes of substituted 2-pyridyl amine based [N,N] chelating ligand
Authors:	Prashanth, B. (/jspui/browse?type=author&value=Prashanth%2C+B.) Karanam, M. (/jspui/browse?type=author&value=Karanam%2C+M.) Choudhury, A.R. (/jspui/browse?type=author&value=Choudhury%2C+A.R.) Singh, Sanjay (/jspui/browse?type=author&value=Singh%2C+Sanjay)
Keywords:	Cobalt complex Copper complex Hydrogen bonding Pyridyl amine
Issue Date:	2012
Publisher:	Elsevier Ltd
Citation:	Polyhedron, 47(1) PP. 112-117.
Abstract:	The reaction of N-benzylaminopyridine with the imidoylchloride of N-(2,6-iPr 2C 6H 3)acetamide in the presence of Et 3N affords a new neutral [N,N] chelating ligand, (PhCH 2)N(2-pyridyl)C((Me)(N-2,6-iPr 2C 6H 3)) (L). The reaction of equimolar quantities of L with Cu(NO 3) 2, CuCl 2 and NiBr 2, respectively, in DCM, acetonitrile and DME yields the corresponding mononuclear complexes L·Cu(NO 3) 2 (1), L·CuCl 2 (2) and L·NiBr 2 (3). Whereas, the reaction of L with CoCl 2·6H 2O leads to the formation of [HL·CoCl 3] (4) with pyridine nitrogen coordinated to cobalt. Solid state structure of L and compounds 1-4 have been investigated by single crystal X-ray structural analysis. The ligand L shows the E-anti arrangement in the solid state and its mononuclear complex 1 shows six coordinated Cu in a quasi square planar geometry with two long distanced donors; complexes 2 and 3 show distorted tetrahedral arrangement of the substituents around metal ions. Interestingly, the solid state structure of complex 4 reveals C-H...Cl intra-molecular hydrogen bonding and N-H...Cl and C-H...Cl inter-molecular hydrogen bonds. These hydrogen bonding interactions in complex 4 facilitate the formation of an extended 2D network structure.
URI:	<a href="https://www.sciencedirect.com/science/article/pii/S0277538712005931">https://www.sciencedirect.com/science/article/pii/S0277538712005931</a> ( <a href="https://www.sciencedirect.com/science/article/pii/S0277538712005931">https://www.sciencedirect.com/science/article/pii/S0277538712005931</a> ) <a href="http://hdl.handle.net/123456789/3501">http://hdl.handle.net/123456789/3501</a> ( <a href="http://hdl.handle.net/123456789/3501">http://hdl.handle.net/123456789/3501</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/3501/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/3501/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/3501?mode=full>)

 (</jspui/handle/123456789/3501/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.