



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2281>


Title:	C. elegans Locomotion: Finding Balance in Imbalance
Authors:	Thapliyal, Shruti (/jspui/browse?type=author&value=Thapliyal%2C+Shruti) Babu, Kavita (/jspui/browse?type=author&value=Babu%2C+Kavita)
Keywords:	Excitation Inhibition Acetylcholine GABA C. elegans
Issue Date:	2018
Publisher:	Springer New York LLC
Citation:	Advances in Experimental Medicine and Biology, 1112, pp. 185-196
Abstract:	<p>The excitation-inhibition (E-I) imbalance in neural circuits represents a hallmark of several neuropsychiatric disorders. The tiny nematode <i>Caenorhabditis elegans</i> has emerged as an excellent system to study the molecular mechanisms underlying this imbalance in neuronal circuits. The <i>C. elegans</i> body wall muscles receive inputs from both excitatory cholinergic and inhibitory GABAergic motor neurons at neuromuscular junctions (NMJ), making it an excellent model for studying the genetic and molecular mechanisms required for maintaining E-I balance at the NMJ. The cholinergic neurons form dyadic synapses wherein they synapse onto ipsilateral body wall muscles allowing for muscle contraction as well as onto GABAergic motor neurons that in turn synapse on the contralateral body wall muscles causing muscle relaxation. An alternating wave of contraction and relaxation mediated by excitatory and inhibitory signals maintains locomotion in <i>C. elegans</i>. This locomotory behavior requires an intricate balance between the excitatory cholinergic signaling and the inhibitory GABAergic signaling mechanisms. Studies on the <i>C. elegans</i> NMJ have provided insights into several molecular mechanisms that could regulate this balance in neural circuits. This review provides a discussion on multiple genetic factors including neuropeptides and their receptors, cell adhesion molecules, and other molecular pathways that have been associated with maintaining E-I balance in <i>C. elegans</i> motor circuits. Further, it also discusses the implications of these studies that could help us in understanding the role of E-I balance in mammalian neural circuits and how changes in this balance could give rise to brain disorders.</p>
URI:	https://link.springer.com/chapter/10.1007%2F978-981-13-3065-0_14 (https://link.springer.com/chapter/10.1007%2F978-981-13-3065-0_14) http://hdl.handle.net/123456789/2281 (http://hdl.handle.net/123456789/2281)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2281/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2281/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/2281?mode=full>)

 (</jspui/handle/123456789/2281/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.