



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Thesis & Dissertation (/jspui/handle/123456789/1)

/ Doctor of Philosophy (PhD) (/jspui/handle/123456789/268)

/ PhD-2014 (/jspui/handle/123456789/462)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1568>

Title:	Group actions on Dold and Milnor manifolds
Authors:	Dey, Pinka (/jspui/browse?type=author&value=Dey%2C+Pinka)
Keywords:	Transformation groups Spectral sequences Dold manifolds \mathbb{Z}_2 -equivariant maps
Issue Date:	Dec-2019
Publisher:	IISER Mohali
Abstract:	<p>The Dold manifold $P(m, n)$ is the quotient of $S^m \times \mathbb{C}P^n$ by the free involution that acts antipodally on the sphere S^m and by complex conjugation on the complex projective space $\mathbb{C}P^n$. In the thesis, we investigate free actions of finite groups on products of Dold manifolds. We show that if a finite group G acts freely and mod 2 cohomologically trivially on a finite-dimensional CW-complex K homotopy equivalent to $P(2m_i, n_i)$, then $G \sim (Z_2)^l$ for some $l \leq k$. This $i=1$ is achieved by first proving a similar assertion for $K \times S^{2m_i} \times \mathbb{C}P^{n_i}$. We also $i=1$ determine the possible mod 2 cohomology algebra of orbit spaces of arbitrary free involutions on Dold Manifolds, and give an application to \mathbb{Z}_2-equivariant maps from spheres to Dold manifolds. We also study free \mathbb{Z}_2 and S^1-actions on cohomology real and complex Milnor manifolds. A real Milnor manifold $RH_{r,s}$ is a non-singular hypersurface of degree $(1, 1)$ in the product $\mathbb{R}P^r \times \mathbb{R}P^s$. A complex Milnor manifold $CH_{r,s}$ is defined analogously. We compute the mod 2 cohomology algebra of the orbit space of an arbitrary free \mathbb{Z}_2 and S^1-action on a compact Hausdorff space with mod 2 cohomology algebra of a real or a complex Milnor manifold. As applications, we deduce some Borsuk-Ulam type results for equivariant maps between spheres and these spaces. For the complex case, we obtain a lower bound on the Schwarz genus, which further establishes the existence of coincidence points for maps from Milnor manifolds to the Euclidean plane.</p>
URI:	http://hdl.handle.net/123456789/1568 (http://hdl.handle.net/123456789/1568)
Appears in Collections:	PhD-2014 (/jspui/handle/123456789/462)

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

ph14072.pdf (/jspui/bitstream/123456789/1568/3/ph14072.pdf)	534.41 kB	Adobe PDF	View/Open (/jspui/bitstream/123456789/1568/3)
--	--------------	--------------	---

[Show full item record \(/jspui/handle/123456789/1568?mode=full\)](/jspui/handle/123456789/1568?mode=full)

 [\(/jspui/handle/123456789/1568/statistics\)](/jspui/handle/123456789/1568/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

Edit...

Export Item

Export (migrate) Item

Export metadata