



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/3482>

Title: Chiral Bent-Shaped Molecules Exhibiting Unusually Wide Range of Blue Liquid-Crystalline Phases and Multistimuli-Responsive Behavior

Authors: Punjani, V. (/jspui/browse?type=author&value=Punjani%2C+V.)
Mohiuddin, G. (/jspui/browse?type=author&value=Mohiuddin%2C+G.)
Kaur, Supreet (/jspui/browse?type=author&value=Kaur%2C+Supreet)
Choudhury, A.R. (/jspui/browse?type=author&value=Choudhury%2C+A.R.)
Pal, S.K. (/jspui/browse?type=author&value=Pal%2C+S.K.)

Keywords: Liquid crystals
Mechanochromism
Phase transitions
Single crystals
Thermochromism

Issue Date: 2020

Publisher: Wiley-VCH Verlag

Citation: Chemistry - A European Journal, 26(26), pp.5859-5871.

Abstract: Recently, an unprecedented observation of polar order, thermochromic behavior, and exotic mesophases in new chiral, bent-shaped systems with a $-CH_3$ moiety placed at the transverse position of the central core was reported. Herein, a homologous series of compounds with even-numbered carbon chains from $n=4$ to 18 were synthesized, in which $-Cl$ was substituted for $-CH_3$ at the kink position and a drastic modification in the phase structure of the bent-shaped molecule was observed. An unusual stabilization of the cubic blue phase (BP) over a wide range of $16.4^\circ C$ has been witnessed. Two homologues in this series (1-12 and 1-14) exhibit an interesting phase sequence consisting of BPI/III, chiral nematic, twist grain boundary, smectic A, and smectic X (SmX) phases. The higher homologues (1-16 and 1-18) stabilize the SmX phase enantiotropically over the entire temperature range. Crystal structure analysis confirmed the bent molecular architecture, with a bent angle of 148° , and revealed the presence of two different molecular conformations in an asymmetric unit of compound 1-4. A DFT study corroborated that the $-Cl$ moiety at the central core of the molecule led to an increase in the dipole moment along the transverse direction, which, in turn, facilitated the unusual stabilization of frustrated structures. Crystal polymorphism has been evidenced in three homologues (1-10, 1-12, and 1-14) of the series. On the application of mechanical pressure through grinding, compound 1-10 transformed from a bright yellow crystalline solid to a dark orange–green amorphous solid, which reversed upon dropwise addition of dichloromethane, indicating reversible mechanochromism in this class of compounds. In addition, excellent thermochromic behavior has been observed for compound 1-10 with a controlled temperature–color combination.

Description: Only IISERM authors are available in the record.


URI: <https://chemistry-europe.onlinelibrary.wiley.com/doi/full/10.1002/chem.201905707>
(<https://chemistry-europe.onlinelibrary.wiley.com/doi/full/10.1002/chem.201905707>)
<http://hdl.handle.net/123456789/3482> (<http://hdl.handle.net/123456789/3482>)

Appears in Collections: Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/3482/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/3482/1/Need%20to%20add%20pdf.odt)

[Show full item record \(/jspui/handle/123456789/3482?mode=full\)](#)

 [\(/jspui/handle/123456789/3482/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.