

## Library Indian Institute of Science Education and Research Mohali



## DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/3042

Title: Pre-pore oligomer formation by Vibrio cholerae cytolysin: Insights from a truncated variant lacking

the pore-forming pre-stem loop

Authors: Paul, Karan (/jspui/browse?type=author&value=Paul%2C+Karan)

Chattopadhyay, K. (/jspui/browse?type=author&value=Chattopadhyay%2C+K.)

Keywords: Vibrio cholerae cytolysin

Pore-forming toxin

fluorescence resonance energy transfer

Circular dichroism

Issue Date:

Publisher:

Elsevier

Citation:

Biochemical and Biophysical Research Communications, 443(1), pp.189-193.

Abstract:

Vibrio cholerae cytolysin (VCC), a  $\beta$ -barrel pore-forming toxin ( $\beta$ -PFT), induces killing of the target eukaryotic cells by forming heptameric transmembrane  $\beta$ -barrel pores. Consistent with the  $\beta$ -PFT mode of action, binding of the VCC toxin monomers with the target cell membrane triggers formation of pre-pore oligomeric intermediates, followed by membrane insertion of the  $\beta$ -strands contributed by the pre-stem motif within the central cytolysin domain of each protomer. It has been shown previously that blocking of membrane insertion of the VCC pre-stem motif arrests conversion of the pre-pore state to the functional transmembrane pore. Consistent with the generalized β-PFT mechanism, it therefore appears that the VCC pre-stem motif plays a critical role toward forming the structural scaffold of the transmembrane β-barrel pore. It is, however, still not known whether the pre-stem motif plays any role in the membrane interaction process, and subsequent pre-pore structure formation by VCC. In this direction, we have constructed a recombinant variant of VCC deleting the pre-stem region, and have characterized the effect(s) of physical absence of the pre-stem motif on the distinct steps of the membrane pore-formation process. Our results show that the deletion of the pre-stem segment does not affect membrane binding and pre-pore oligomer formation by the toxin, but it critically abrogates the functional poreforming activity of VCC. Present study extends our insights regarding the structure-function mechanism associated with the membrane pore formation by VCC, in the context of the  $\beta$ -PFT mode of action.

URI:

https://www.sciencedirect.com/science/article/pii/S0006291X13019797?via%3Dihub (https://www.sciencedirect.com/science/article/pii/S0006291X13019797?via%3Dihub) http://hdl.handle.net/123456789/3042 (http://hdl.handle.net/123456789/3042)

Appears in Collections:

Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File Description Size Format

need to add pdf....odt 8.12 OpenDocument (/jspui/bitstream/123456789/3042/1/need%20to%20add%20pdf....odt) kB Text

View/Open (/jspui/bitstream/1234

Show full item record (/jspui/handle/123456789/3042?mode=full)

■ (/jspui/handle/123456789/3042/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.