



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4628>

Title:	The Hercules cluster in X-rays with XMM-Newton and Chandra.
Authors:	Tiwari, Juhi (/jspui/browse?type=author&value=Tiwari%2C+Juhi) Singh, Kulinder Pal (/jspui/browse?type=author&value=Singh%2C+Kulinder+Pal)
Keywords:	galaxies: clusters: general X-rays: galaxies: clusters galaxies: clusters: intracluster medium
Issue Date:	2021
Citation:	Monthly Notices of the Royal Astronomical Society, 500(4), 5524–5542.
Abstract:	We present a detailed X-ray study of the central subcluster of the nearby ( $z \sim 0.0368$ ) Hercules cluster (Abell 2151) identified as A2151C that shows a bimodal structure. A bright clump of hot gas with X-ray emission extending to radius $r \sim 304$ kpc and $L_X = 3.03 \pm 0.02 - 0.04 \times 10^{43}$ erg s $^{-1}$ in the 0.4–7.0 keV energy range is seen as a fairly regular subclump towards the west (A2151C(B)). An irregular, fainter and cooler subclump with radius $r \sim 364$ kpc is seen towards the east (A2151C(F)) and has $L_X = 1.13 \pm 0.02 \times 10^{43}$ erg s $^{-1}$ in the 0.4–7.0 keV energy band. The average temperature and elemental abundance of A2151C(B) are $2.01 \pm 0.05$ keV and $0.43 \pm 0.05 Z_\odot$ , respectively, while these values are $1.17 \pm 0.04$ keV and $0.13 \pm 0.02 Z_\odot$ for A2151C(F). Low temperature ( $1.55 \pm 0.07$ keV) and a short cooling time ( $\sim 0.81$ Gyr) within the central 15 arcsec region confirm the presence of a cool core in A2151C(B). We identify several compact groups of galaxies within A2151C(F). We find that A2151C(F) is a distinct galaxy group in the process of formation and likely not a ram-pressure stripped part of the eastern subcluster in Hercules (A2151E). X-ray emission from A2151C shows a region of overlap between A2151C(B) and A2151C(F) but without any enhancement of temperature or entropy in the two-dimensional (2D) projected thermodynamic maps that could have indicated an interaction due to a merger between the two subclumps.
Description:	Only IISERM authors are available in the record.
URI:	<a href="https://doi.org/10.1093/mnras/staa3619">https://doi.org/10.1093/mnras/staa3619</a> ( <a href="https://doi.org/10.1093/mnras/staa3619">https://doi.org/10.1093/mnras/staa3619</a> ) <a href="http://hdl.handle.net/123456789/4628">http://hdl.handle.net/123456789/4628</a> ( <a href="http://hdl.handle.net/123456789/4628">http://hdl.handle.net/123456789/4628</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF. (/jspui/bitstream/123456789/4628/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF.)		15.36 kB	Unknown

[View/Open \(/jspui/\)](#)

[Show full item record \(/jspui/handle/123456789/4628?mode=full\)](#)

[Statistics \(/jspui/handle/123456789/4628/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.