



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**

**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**

**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2508>

**Title:** Does a hydrogen bonded complex with dual contacts show synergism? A matrix isolation infrared and ab-initio study of propargyl alcohol–water complex

**Authors:** Saini, J. (/jspui/browse?type=author&value=Saini%2C+J.)  
Viswanathan, K.S. (/jspui/browse?type=author&value=Viswanathan%2C+K.S.)

**Keywords:** Propargyl alcohol  
Matrix isolation  
Hydrogen bonding  
Conformations

**Issue Date:** 2016

**Publisher:** Elsevier

**Citation:** Journal of Molecular Structure, 1118, pp. 147-156.

**Abstract:** When hydrogen bonded complexes are formed with more than one contact, the question arises if these multiple contacts operate synergistically. Propargyl alcohol-H<sub>2</sub>O complex presents a good case study to address this question, which is discussed in this work. Complexes of propargyl alcohol (PA) and H<sub>2</sub>O were studied experimentally using matrix isolation infrared spectroscopy, which was supported by quantum chemical computations performed at the M06-2X and MP2 level of theories, using 6-311++G (d,p) and aug-cc-pVDZ basis sets. A 1:1 PA-H<sub>2</sub>O complex was identified in the experiments and corroborated by our computations, where the PA was in the gauche conformation. This complex, which was a global minimum, showed dual interactions, one of which was an n-σ interaction between the O–H group of PA and the O of H<sub>2</sub>O, while the second was a H···π contact between the O–H group of H<sub>2</sub>O and the π system of PA. We explored if the two interactions in the 1:1 complex exhibited synergism. We finally argue that the two interactions showed antagonism rather than synergism. Our computations indicated three other local minima for the 1:1 complexes; though these local minima were not identified in our experiments. Atoms-in-molecules and energy decomposition analysis executed through LMO-EDA were also performed to understand the nature of intermolecular interactions in the PA-H<sub>2</sub>O complexes. We have also revisited the problem of conformations of PA, with a view to understanding the reasons for gauche conformational preferences in PA.

**URI:** <https://www.sciencedirect.com/science/article/pii/S0022286016303143>  
(<https://www.sciencedirect.com/science/article/pii/S0022286016303143>)  
<http://hdl.handle.net/123456789/2508> (<http://hdl.handle.net/123456789/2508>)


**Appears in Collections:** Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need to add pdf.odt (/jspui/bitstream/123456789/2508/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text

[View/Open \(/jspui/bitstream/123456789/2508/1/Need%20to%20add%20pdf.odt\)](#)

Show full item record (</jspui/handle/123456789/2508?mode=full>)

 (</jspui/handle/123456789/2508/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.