





Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Master of Science / MS-19

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/5785

Analytic Theory of Time-Propagators in the Strong Coupling Regime

Authors: Das, Soumyadev

Keywords: Analytic

Time-Propagators

Regime

Issue Date:

Title

May-2024

Publisher:

IISER Mohali

Abstract:

NMR methodology concerns about tuning the behaviour of a spin system (considering the effects of local interactions) for achieving a desired pulsed sequence. In order to do so, it is imperative to have a complete understanding about the dynamics of the spin system (based on the principles of quantum mechanics). In general, its become difficult to formulate an analytic description of the time evolution of a spin system due to the time dependency of the Hamiltonian. The current study discusses the efficacy of an averaging scheme in deriving the time propagators (of a time dependent Hamiltonian). At first, an ideal two state spin system is considered and an analytic description of the spin system is proposed which concerns about the time evolution of the system. The primary focus of the study is to observe the effect of amplitude of the driving field on time evolution of the system. Owing to the complicacies (time dependency) of the Hamiltonian, the corresponding time propagators are derived using an averaging scheme (Floquet Magnus Expansion). After probing the efficiency of the said averaging scheme the Hamiltonian of a periodically driven (say rotating solid) system commonly encountered in NMR spec- troscopy is discussed.

URI: http://hdl.handle.net/123456789/5785

Appears in Collections:

MS-19

Files in This Item:

File	Description	Size	Format	
embargo period.pdf		6.04 kB	Adobe PDF	View/Open

Show full item record

di

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Theme by CINEC

Customized & Implemented by - Jivesna Tech