

## Library Indian Institute of Science Education and Research Mohali



## DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/4454

Title: Spatiotemporal modulations in heterotypic condensates of prion and  $\alpha$ -synuclein control phase

transitions and amyloid conversion

Authors: Agarwal, Aishwarya (/jspui/browse?type=author&value=Agarwal%2C+Aishwarya)

Arora, Lisha (/jspui/browse?type=author&value=Arora%2C+Lisha)
Rai, Sandeep K. (/jspui/browse?type=author&value=Rai%2C+Sandeep+K.)

Rai, Sandeep K. (/jspui/browse?type=author&value=Rai%2C+Sandeep+K.)

Avni, Anamika (/jspui/browse?type=author&value=Avni%2C+Anamika)

Mukhopadhyay, Samrat (/jspui/browse?type=author&value=Mukhopadhyay%2C+Samrat)

Keywords: Spatiotemporal modulations

heterotypic condensates of prion and  $\alpha$ -synuclein control

amyloid conversion

Issue Date: 2022

Publisher: Springer Nature

Citation: Nature Communications, 13(1), 1154

Abstract:

Biomolecular condensation via liquid-liquid phase separation of proteins and nucleic acids is associated with a range of critical cellular functions and neurodegenerative diseases. Here, we demonstrate that complex coacervation of the prion protein and  $\alpha$ -synuclein within narrow stoichiometry results in the formation of highly dynamic, reversible, thermo-responsive liquid droplets via domain-specific electrostatic interactions between the positively-charged intrinsically disordered N-terminal segment of prion and the acidic C-terminal tail of  $\alpha$ -synuclein. The addition of RNA to these coacervates yields multiphasic, vesicle-like, hollow condensates. Picosecond time-resolved measurements revealed the presence of transient electrostatic nanoclusters that are stable on the nanosecond timescale and can undergo breaking-and-making of interactions on slower timescales giving rise to a liquid-like behavior in the mesoscopic regime. The liquid-to-solid transition drives a rapid conversion of complex coacervates into heterotypic amyloids. Our results suggest that synergistic prion- $\alpha$ -synuclein interactions within condensates provide mechanistic underpinnings of their physiological role and overlapping neuropathological features

Description: Only IISERM authors are available in the record

URI: http://hdl.handle.net/123456789/4454 (http://hdl.handle.net/123456789/4454)

Appears in Research Articles (/jspui/handle/123456789/9) Collections:

Files in This Item:

 File
 Description
 Size
 Format

 Need To Add...Full Text\_PDF..pdf
 15.36
 Adobe (/jspui/bitstream/123456789/4454/1/Need%20To%20Add%e2%80%a6Full%20Text\_PDF..pdf)
 View/Open (/jspui/bitstream/123456789/4454/1/Need%20Text\_PDF..pdf)
 View/Open (/jspui/bitstream/123456789/4454/1/Need%20Text\_PDF..pdf)

Show full item record (/jspui/handle/123456789/4454?mode=full)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.