

## Library Indian Institute of Science Education and Research Mohali



## DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/3359

Title: Structural basis of the strong cell-cell junction formed by cadherin-23

Authors: Singaraju, G.S. (/jspui/browse?type=author&value=Singaraju%2C+G.S.)

Sagar, A. (/jspui/browse?type=author&value=Sagar%2C+A.)
Kumar, Anuj (/jspui/browse?type=author&value=Kumar%2C+Anuj)
Samuel, J.S. (/jspui/browse?type=author&value=Samuel%2C+J.S.)
Hazra, J.P. (/jspui/browse?type=author&value=Hazra%2C+J.P.)

Sannigrahi, M.K. (/jspui/browse?type=author&value=Sannigrahi%2C+M.K.)

Rakshit, S. (/jspui/browse?type=author&value=Rakshit%2C+S.)

Keywords: Atypical cadherins

Cadherin-23
Cell–cell adhesion

Single-molecule Forster resonance energy transfer small-angle X-ray scattering

Small-angle X-ray scattering

Issue Date: 2020

Publisher: Blackwell Publishing

Citation: FEBS Journal, 287(11), pp.2328-2347.

Abstract:

Cadherin-23, a giant atypical cadherin, form homophilic interactions at the cell–cell junction of epithelial cells and heterophilic interactions with protocadherin-15 at the tip links of neuroepithelial cells. While the molecular structure of the heterodimer is solved, the homodimer structure is yet to be resolved. The homodimers play an essential role in cell–cell adhesion as the downregulation of cadherin-23 in cancers loosen the intercellular junction resulting in faster migration of cancer cells and a significant drop in patient survival. In vitro studies have measured a stronger aggregation propensity of cadherin-23 compared to typical E-cadherin. Here, we deciphered the unique transhomodimer structure of cadherin-23 in solution and show that it consists of two electrostatic-based interfaces extended up to two terminal domains. The interface is robust, with a low off-rate of  $\sim 8 \times 10^{-4} \, s^{-1}$  that supports its strong aggregation propensity. We identified a point mutation, E78K, that disrupts this binding. Interestingly, a mutation at the interface was reported in skin cancer. Overall, the structural basis of the strong cadherin-23 adhesion may have far-reaching

applications in the fields of mechanobiology and cancer.

Description: Only IISERM authors are available in the record.

URI: https://febs.onlinelibrary.wiley.com/doi/full/10.1111/febs.15141 (https://febs.onlinelibrary.wiley.com/doi/full/10.1111/febs.15141)

http://hdl.handle.net/123456789/3359 (http://hdl.handle.net/123456789/3359)

Appears in Research Articles (/jspui/handle/123456789/9)

Collections:

Files in This Item:

| File   | Description | Size       | Format               |                                  |
|--|-------------|------------|----------------------|----------------------------------|
| need to add pdfodt<br>(/jspui/bitstream/123456789/3359/1/need%20to%20add%20pdfodt) |             | 8.12<br>kB | OpenDocument<br>Text | View/Open (/jspui/bitstream/1234 |

Show full item record (/jspui/handle/123456789/3359?mode=full)

■ (/jspui/handle/123456789/3359/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.