



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**  
**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**  
**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/2594>

Title:	When is $R[\theta]$ integrally closed?
Authors:	Khanduja, S.K. (/jspui/browse?type=author&value=Khanduja%2C+S.K.) Jhorar, B. (/jspui/browse?type=author&value=Jhorar%2C+B.)
Keywords:	Valued fields Irreducible polynomials Non-Archimedean valued fields
Issue Date:	2016
Publisher:	World Scientific
Citation:	Journal of Algebra and its Applications, 15(5).
Abstract:	<p>Let <math>R</math> be an integrally closed domain with quotient field <math>K</math> and <math>\theta</math> be an element of an integral domain containing <math>R</math> with <math>\theta</math> integral over <math>R</math>. Let <math>F(x)</math> be the minimal polynomial of <math>\theta</math> over <math>K</math> and <math>p</math> be a maximal ideal of <math>R</math>. Kummer proved that if <math>R[\theta]</math> is an integrally closed domain, then the maximal ideals of <math>R[\theta]</math> which lie over <math>p</math> can be explicitly determined from the irreducible factors of <math>F(x)</math> modulo <math>p</math>. In 1878, Dedekind gave a criterion known as Dedekind Criterion to be satisfied by <math>F(x)</math> for <math>R[\theta]</math> to be integrally closed in case <math>R</math> is the localization <math>Z(p)</math> of <math>Z</math> at a nonzero prime ideal <math>pZ</math> of <math>Z</math>. Indeed he proved that if <math>g_1(x)e_1 \cdots g_r(x)e_r</math> is the factorization of <math>F(x)</math> into irreducible polynomials modulo <math>p</math> with <math>g_i(x) \in Z[x]</math> monic, then <math>Z(p)[\theta]</math> is integrally closed if and only if for each <math>i</math>, either <math>e_i = 1</math> or <math>g_i(x)</math> does not divide <math>H(x)</math> modulo <math>p</math>, where <math>H(x) = 1p(F(x) - g_1(x)e_1 \cdots g_r(x)e_r)</math>. In 2006, a similar necessary and sufficient condition was given by Ershov for <math>R[\theta]</math> to be integrally closed when <math>R</math> is the valuation ring of a Krull valuation of arbitrary rank (see [Comm. Algebra. 38 (2010) 684–696]). In this paper, we deal with the above problem for more general rings besides giving some equivalent versions of Dedekind Criterion. The well-known result of Uchida in this direction proved for Dedekind domains has also been deduced (cf. [Osaka J. Math. 14 (1977) 155–157]).</p>
URI:	<a href="https://www.worldscientific.com/doi/abs/10.1142/S0219498816500912">https://www.worldscientific.com/doi/abs/10.1142/S0219498816500912</a> ( <a href="https://www.worldscientific.com/doi/abs/10.1142/S0219498816500912">https://www.worldscientific.com/doi/abs/10.1142/S0219498816500912</a> ) <a href="http://hdl.handle.net/123456789/2594">http://hdl.handle.net/123456789/2594</a> ( <a href="http://hdl.handle.net/123456789/2594">http://hdl.handle.net/123456789/2594</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

## Files in This Item:

File	Description	Size	Format	
Need to add pdf.odt (/jspui/bitstream/123456789/2594/1/Need%20to%20add%20pdf.odt)		8.63 kB	OpenDocument Text	<a href="#">View/Open (/jspui/bitstream/123456789/2594/1/Need%20to%20add%20pdf.odt)</a>

[Show full item record \(/jspui/handle/123456789/2594?mode=full\)](#)

[Statistics \(/jspui/handle/123456789/2594/statistics\)](#)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.