



# Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali / Thesis & Dissertation / Doctor of Philosophy (PhD) / PhD-2017

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/5894>

Title:	On the local constancy of certain mod $p$ Galois representations
Authors:	<a href="#">Kumar, Suneel</a>
Keywords:	Galois representations binomial identities Hecke algebras
Issue Date:	Jul-2023
Publisher:	IISER Mohali
Abstract:	In this thesis, we study local constancy in the weight space of the mod $p$ reduction of certain irreducible 2-dimensional crystalline representations $V_{k,a,p}$ of $\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ using the mod $p$ local Langlands correspondence for $\text{GL}_2(\mathbb{Q}_p)$ . Indeed all irreducible 2-dimensional crystalline representations of $\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ are a twist by a crystalline character of some $V_{k,a,p}$ . Latter is uniquely determined by the Hodge-Tate weights $(0, k-1)$ and the trace of the Frobenius $a_p$ . Let $V_{k,a,p}$ denote the mod $p$ reduction of a $\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ -stable lattice up to semisimplification. For a fixed $a_p$ , we study local constancy of $V_{k,a,p}$ in the weight space. In order to prove local constancy, we show that $V_{k',a_p}$ is constant for all $k' \equiv k + p^t (p-1)\mathbb{Z} > 0$ ( $t$ sufficiently large) by explicitly computing $V_{k',a_p}$ . Latter is obtained using the mod $p$ local Langlands correspondence for $\text{GL}_2(\mathbb{Q}_p)$ . This gives local constancy in the punctured disk $\{k' \mid k' \equiv k + p^t (p-1)\mathbb{Z} > 0\}$ around $k$ . Next, we determine the structure of $V_{k,a,p}$ by applying a result of Berger-Li-Zhu together with Berger's local constancy theorem to establish constancy in the whole disk. This gives a lower bound on the radius of local constancy solely in terms of $a_p$ . Let $k = b + c(p-1) + 2$ where $2 \leq b \leq p$ and $0 \leq c \leq p-2$ . Broadly speaking, our results fall into two regimes $(c, p-1)$ and $(1, c+1)$ of the slope $v(a_p)$ . In the former situation, we establish local constancy under certain constraints on $b, c$ and $v(a_p)$ . We also in this case improve on the lower bound for $k$ in Berger's theorem guaranteeing local constancy. When $v(a_p) < c+1$ , we establish local constancy (in the irreducible cases) for non-integral slope under the condition that $b \geq c + \lfloor v(a_p) \rfloor - 1$ holds. Lastly, assuming a certain technical conjecture (for which we provide good evidence), we establish local constancy for certain subsets of $b \leq c + \lfloor v(a_p) \rfloor - 2$ and such that the slope is non-integral, lying in $(1, c-1)$ .
URI:	<a href="http://hdl.handle.net/123456789/5894">http://hdl.handle.net/123456789/5894</a>
Appears in Collections:	<a href="#">PhD-2017</a>

## Files in This Item:

File	Description	Size	Format	
<a href="#">Suneel(PH17043)PhD_Thesis_final.pdf</a>		926.9 kB	Adobe PDF	<a href="#">View/Open</a>

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

## Admin Tools

Edit...

Export Item

Export (migrate) Item

Export metadata