



**Library**  
**Indian Institute of Science Education and Research**  
**Mohali**



**DSpace@IISERMohali (/jspui/)**

**/ Thesis & Dissertation (/jspui/handle/123456789/1)**

**/ Master of Science (/jspui/handle/123456789/2)**

**/ MS-12 (/jspui/handle/123456789/723)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/752>

Title:	Identifying New Players Involved in Ubiquinone Biosynthesis in Escherichia coli
Authors:	Balecha, Himanshi (/jspui/browse?type=author&value=Balecha%2C+Himanshi)
Keywords:	Biology E. coli Biosynthesis Quinones
Issue Date:	13-Jul-2017
Publisher:	IISER-M

**Abstract:** Quinones are lipid soluble electron carriers in the electron transport chain (ETC) and help in energy generation. They can be broadly divided into benzoquinones (e.g. ubiquinone) involved in aerobic ETC, and naphthoquinones (e.g. menaquinone and demethylmenaquinone) involved in anaerobic respiration. Our lab conducted a highthroughput genetic screen on a long chain fatty acid (LCFA), oleate, in *Escherichia coli* and compared the LCFA dataset with published genome-wide screens on other carbon sources. This comparative analysis revealed highest requirement of ubiquinone for growth in oleate. In particular, a  $\Delta$ ubil strain (Ubil is a hydroxylase involved in ubiquinone biosynthesis), which produces very low level of ubiquinone exhibited growth defect only in oleate. We thus hypothesized that some of the uncharacterized genes (ygenes) in *E. coli*, deletion of which led to significant growth defect only in oleate, might also be involved in ubiquinone biosynthesis. Detailed studies on one such candidate, yqiC (yqiC encodes a small protein with predicted scaffold function) showed that it is involved in ubiquinone biosynthesis since its deletion resulted in low ubiquinone content. Moreover, the phenotype of  $\Delta$ yqiC in terms of reduced ubiquinone levels and accumulation of an intermediate was similar to  $\Delta$ ubil. Furthermore, a  $\Delta$ yqiC $\Delta$ ubil double knockout showed a synthetic sick phenotype with no detectable ubiquinone suggesting that the two genes act at the same step in ubiquinone biosynthesis. Introduction of multicopy yqiC clone in  $\Delta$ ubil showed slight growth recovery in oleate suggesting that YqiC might promote the function of some other hydroxylase in the absence of Ubil to rescue ubiquinone levels. Since YqiC in *Salmonella typhimurium* is involved in menaquinone biosynthesis, it is also possible that overexpression of yqiC in  $\Delta$ ubil rescues the growth defect of  $\Delta$ ubil in oleate by regulating the synthesis of naphthoquinones. In addition to understanding the role of YqiC in ubiquinone biosynthesis, we are also investigating additional y-genes for their involvement in ubiquinone biosynthesis.

**URI:** <http://hdl.handle.net/123456789/752> (<http://hdl.handle.net/123456789/752>)

**Appears in** MS-12 (</jspui/handle/123456789/723>)  
**Collections:**

Files in This Item:

File	Description	Size	Format	
MS-12097.pdf ( <a href="/jspui/bitstream/123456789/752/3/MS-12097.pdf">/jspui/bitstream/123456789/752/3/MS-12097.pdf</a> )		3.83 MB	Adobe PDF	<a href="/jspui/bitstream/123456789/752/3/MS-12097.pdf">View/Open (/jspui/bitstream/123456789/752/3/MS-12097.pdf)</a>

[Show full item record \(/jspui/handle/123456789/752?mode=full\)](/jspui/handle/123456789/752?mode=full)

[📊 \(/jspui/handle/123456789/752/statistics\)](/jspui/handle/123456789/752/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.