

## Library Indian Institute of Science Education and Research Mohali



## DSpace@IISERMohali / Thesis & Dissertation / Doctor of Philosophy (PhD) / PhD-2017

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/5328

Title: Metal-free Transformations of para-Quinone Methides (p-QMs) to Oxygen-Containing Heterocycles and Carbazoles

Authors: Singh, Gurdeep

Keywords: Para-Quinone Oxvaen

Heterocycles

Issue

27-Jan-2023

Date:

Publisher: IISER Mohali

Abstract:

Oxygen and nitrogen-containing heterocycles such as benzo[b]furan, dihydrobenzofuran, coumarins, chromenes, and carbazoles are important scaffolds and widely found in many natural products and biologically active molecules, and several of them display various impressive pharmacological activities such as anticancer, anti-allergic, anti-HIV, cytotoxic, etc. Due to their fascinating and diverse range of applications in material science as well, numerous synthetic techniques have been documented to access these heterocycles, but most of them require pre-functionalized starting material(s), poor substrate scope, lack of regioselectivity, and an expensive metal catalyst. Therefore, developing practical and metal- free protocols to access these derivatives is highly desirable and advantageous. In the recent past, para-quinone methide chemistry has been widely explored for the synthesis of various organic frameworks such as heterocycles, carbocycles, spiro-cycles, etc. In the present work, we have utilized the para-quinone methides (p-QMs) as a 1,6-conjugate acceptor for the synthesis of oxygen and nitrogen-containing heterocycles under metal-free reaction conditions. The first chapter of this work involves the synthesis of 2,3-diaryl benzo[b]furans, dihydrobenzofurans, 3,4-diaryl coumarins, and chromenes by the reaction of 2-hydroxyphenyl substituted para-quinone methides with suitable coupling partners under acidic/basic conditions. The second chapter describes the [3+3] annulation of 2- indolylmethanols with para-quinone methides for the formation of tetrahydroindolo[2,3-b]carbazoles. Further, this work has been extended to synthesize different isomers of tetrahydrothieno carbazoles.

URI: http://hdl.handle.net/123456789/5328

Appears in Collections:

PhD-2017

Files in This Item:

 File
 Description
 Size
 Format

 Full thesis.pdf
 20.02 MB
 Adobe PDF

 View/Open

Show full item record



Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.

Admin Tools

Edit...

Export Item

Export (migrate) Item

Export metadata

