



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

/ Publications of IISER Mohali (/jspui/handle/123456789/4)

/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/1658>

Title:	Structural organization and molecular self-assembly of a new class of polar and non-polar four-ring based bent-core molecules
Authors:	Kaur, Supreet (/jspui/browse?type=author&value=Kaur%2C+Supreet) Mohiuddin, G. (/jspui/browse?type=author&value=Mohiuddin%2C+G.) Punjani, V. (/jspui/browse?type=author&value=Punjani%2C+V.) Pal, S.K. (/jspui/browse?type=author&value=Pal%2C+S.K.)
Keywords:	Bent-core molecules Self-assembly Cybotactic nematic phase Single crystal DFT study
Issue Date:	2019
Publisher:	Elsevier
Citation:	Journal of Molecular Liquids, 295.
Abstract:	We have studied two new series of polar and non-polar four-ring based achiral bent-core molecules containing single bondCN and single bondOCH ₃ moiety respectively at the tip of the molecular short arm and their molecular self-assembly in crystal and liquid crystal state. Crystal structure analysis revealed the bent molecular geometry of the compounds with bent angle 151° along with fundamental intermolecular interactions such as antiparallel arrangement of the polar molecules, intermolecular hydrogen bonding, different types of π - π interactions and entangled/helical molecular arrangement in 3D space. Due to the dimeric antiparallel molecular arrangement, lower homologues of the polar compounds of series 1-n (1-8, 1-10) exhibited long range nematic phase with cybotactic clustering (Ncyb). However, the lower homologues of non-polar compounds of series 2-n (2-6, 2-8) did not form any type of clustering exhibiting normal nematic phase. Dielectric spectroscopic investigations and X-ray diffraction measurements of compound 1-8 confirmed the presence of nematic cybotacticity. Thus, the formation of clusters in nematic phase of bent-core molecules was not only due to an underlying smectic phase but also due to the combined effect of the molecular structural parameters including bent angle, molecular dipole moment and its spatial arrangement. Negative bend-splay (K ₃₃ -K ₁₁) elastic constant behavior was observed in the entire nematic range.
Description:	Only IISER Mohali authors are available in the record.
URI:	https://www.sciencedirect.com/science/article/pii/S0167732219329472 (https://www.sciencedirect.com/science/article/pii/S0167732219329472) http://hdl.handle.net/123456789/1658 (http://hdl.handle.net/123456789/1658)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
------	-------------	------	--------

Need to add pdf.odt (/jspui/bitstream/123456789/1658/1/Need%20to%20add%20pdf.odt)	8.63 kB	OpenDocument Text	View/Open (/jspui/bitstream/123456789/1658/1/Need%20to%20add%20pdf.odt)
--	------------	----------------------	---

Show full item record (/jspui/handle/123456789/1658?mode=full)

 (/jspui/handle/123456789/1658/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.