

Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)

- / Publications of IISER Mohali (/jspui/handle/123456789/4)
- / Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: http://hdl.handle.net/123456789/2539

Title: Regio- and diastereoselective construction of a new set of functionalized pyrrolidine,

 $spiropyrrolidine\ and\ spiropyrrolizidine\ scaffolds\ appended\ with\ aryl-\ and\ heteroaryl\ moieties\ via\ the$

azomethine ylide cycloadditions

Authors: Rajkumar, V. (/jspui/browse?type=author&value=Rajkumar%2C+V.)

Babu, S.A. (/jspui/browse?type=author&value=Babu%2C+S.A.)

Padmavathi, R. (/jspui/browse?type=author&value=Padmavathi%2C+R.)

Keywords: Azomethine ylide

1,3-Dipolar cycloaddition Diastereoselectivity Spiropyrrolidines Spirooxindoles

Issue Date: 2016

Publisher: Elsevier Ltd

Citation: Tetrahedron, 72(36), pp. 5578-5594

Abstract:

Highly regio- and diastereoselective syntheses of a new set of functionalized pyrrolidines, spiro-pyrrolidine/pyrrolizidine oxindoles, spiroacenaphthylenolyl-pyrrolidines/pyrrolizidines and spiro-1,3-indandionolyl-pyrrolidines/pyrrolizidines appended with various aryl- and heteroaryl moieties via the azomethine ylide cycloaddition reaction are reported. The Ag-catalyzed [3+2] cycloaddition of azomethine ylides derived from N-benzylideneiminoglycinates with various arylidene/heteroarylidenemalononitriles gave C-3,C-5-aryl/heteroaryl substituted C-4,C-4-dicyanopyrrolidine-2-carboxylate scaffolds with good regio- and diastereoselectivity. Further, the [3+2] cycloaddition of azomethine ylides derived from the decarboxylative reactions of different 1,2-dicarbonyls and α -amino acids with the indole/pyrrole-based dipolarophiles were investigated. In the context of enriching the library of functionalized spiropyrrolidine- and spiropyrrolizidine scaffolds, these reactions have led to the assembling of various spiro-pyrrolidine/pyrrolizidine oxindoles, spiroacenaphthylenolyl-pyrrolidines/pyrrolizidines and spiro-1,3-indandionolyl-pyrrolidines/pyrrolizidines appended with the indolyl- and pyrrolyl moieties at the C-3 position of the spiro-pyrrolidine/pyrrolizidine rings. The stereochemistry of the cycloadducts was assigned based on the single crystal X-ray structures of representative major diastereomers 8c, 8d, 9b, 9c, 17a,

URI:

https://www.sciencedirect.com/science/article/pii/S0040402016307037 (https://www.sciencedirect.com/science/article/pii/S0040402016307037) http://hdl.handle.net/123456789/2539 (http://hdl.handle.net/123456789/2539)

17b, 17c, 18a, 18e and 19a obtained from the azomethine ylide cycloaddition reactions

Appears in Collections:

Research Articles (/jspui/handle/123456789/9)

Files in This Item:

FileDescriptionSizeFormatNeed to add pdf.odt
(/jspui/bitstream/123456789/2539/1/Need%20to%20add%20pdf.odt)7.9OpenDocument
kBView/Open (/jspui/bitstream/12345

Show full item record (/jspui/handle/123456789/2539?mode=full)

■ (/jspui/handle/123456789/2539/statistics)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.