



# Library Indian Institute of Science Education and Research Mohali



**DSpace@IISERMohali (/jspui/)**  
**/ Publications of IISER Mohali (/jspui/handle/123456789/4)**  
**/ Research Articles (/jspui/handle/123456789/9)**

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4875>


Title:	Understanding the Influence of Meteorology and Emission Sources on PM <sub>2.5</sub> Mass Concentrations Across India: First Results From the COALESCE Network
Authors:	Chaudhary, Pooja (/jspui/browse?type=author&value=Chaudhary%2C+Pooja) Sinha, Baerbel (/jspui/browse?type=author&value=Sinha%2C+Baerbel)
Keywords:	Emission Sources Mass Concentrations COALESCE Network
Issue Date:	2022
Publisher:	Wiley
Citation:	Journal of Geophysical Research: Atmospheres, 127(4), 42005
Abstract:	<p>The Carbonaceous Aerosol Emissions, Source Apportionment and Climate Impacts (COALESCE) is a multi-institutional Indian network project to better understand carbonaceous aerosol induced air quality and climate effects. This study presents time synchronized measurements of surface PM<sub>2.5</sub> concentrations made during 2019 at 11 COALESCE sites across India. The network median PM<sub>2.5</sub> concentration was 42 µg m<sup>-3</sup> with the highest median value at Rohtak (99 µg m<sup>-3</sup>) and the lowest median value at Mysuru (26 µg m<sup>-3</sup>). The influence of six meteorological parameters on PM<sub>2.5</sub> were evaluated. Causality analysis suggested that temperature, surface pressure, and relative humidity were the most important factors influencing fine PM mass, on an annual as well as seasonal scale. Further, a multivariable linear regression model showed that, on an annual basis, meteorology could explain 16%–41% of PM<sub>2.5</sub> variability across the network. Concentration Weighted Trajectories (CWT) together with the results of causality analysis revealed common regional sources affecting PM<sub>2.5</sub> concentrations at multiple regional sites. Further, CWT source locations for all sites across the network correlated with the SMOG-India emissions inventory at the 95th percentile confidence. Finally, CWT maps in conjunction with emissions inventory were used to obtain quantitative estimates of anthropogenic primary PM<sub>2.5</sub> sectoral shares from a mass-meteorology-emissions reconciliation, for all 11 pan-India network sites. These estimates can help guide immediate source reduction and mitigation actions at the national level.</p>
Description:	Only IISER Mohali authors are available in the record.
URI:	<a href="https://doi.org/10.1029/2021JD035663">https://doi.org/10.1029/2021JD035663</a> ( <a href="https://doi.org/10.1029/2021JD035663">https://doi.org/10.1029/2021JD035663</a> ) <a href="http://hdl.handle.net/123456789/4875">http://hdl.handle.net/123456789/4875</a> ( <a href="http://hdl.handle.net/123456789/4875">http://hdl.handle.net/123456789/4875</a> )
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF. (/jspui/bitstream/123456789/4875/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF.)		15.36 kB	Unknown

[View/Open \(/jspui/\)](#)

Show full item record (</jspui/handle/123456789/4875?mode=full>)

 (</jspui/handle/123456789/4875/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.