



Library Indian Institute of Science Education and Research Mohali



DSpace@IISERMohali (/jspui/)
/ Publications of IISER Mohali (/jspui/handle/123456789/4)
/ Research Articles (/jspui/handle/123456789/9)

Please use this identifier to cite or link to this item: <http://hdl.handle.net/123456789/4895>


Title:	Nickel-Catalyzed Selective Synthesis of α -Alkylated Ketones via Dehydrogenative Cross-Coupling of Primary and Secondary Alcohols
Authors:	Bains, Amreen K (/jspui/browse?type=author&value=Bains%2C+Amreen+K) Biswas, Ayanangshu (/jspui/browse?type=author&value=Biswas%2C+Ayanangshu) Adhikari, Debashis (/jspui/browse?type=author&value=Adhikari%2C+Debashis)
Keywords:	Nickel-Catalyzed α -Alkylated Ketones Dehydrogenative Cross-Coupling Primary and Secondary Alcohols
Issue Date:	2022
Publisher:	John Wiley & Sons
Citation:	Advanced Synthesis and Catalysis, 364(1), 47-52.
Abstract:	Herein, we describe an isolable, air-stable, homogeneous, nickel catalyst that performs dehydrogenative cross-coupling reaction between secondary and primary alcohols to result α -alkylated ketone products selectively. The sequence of steps involve in this one-pot reaction is dehydrogenation of both alcohols, condensation between the ketone and the aldehyde, and hydrogenation of the in situ-generated α,β -unsaturated ketone. Preliminary mechanistic investigation hints a radical mechanism following borrowing hydrogen reaction. The construction of C-C bond with structural diversity and complexity is quintessential in organic synthesis. ¹ Traditional approaches toward their fabrication include nucleophilic attack by an appropriate C-nucleophile to the alkyl halide or similar alkylating agent. ² This approach often requires cryogenic conditions, employs toxic or mutagenic alkyl halides and generates a copious amount of inorganic waste. Henceforth, alternative methods are intensely sought after that can be environmentally benign, atom- and process-efficient by utilizing cheap, nontoxic starting materials. In this regard, borrowing hydrogen (BH) approach provides a very promising solution, where biomass-derived, abundantly available alcohols can be the starting feedstocks. ³ In BH method, hydrogen is extracted from the alcohol, kept stored in the catalyst and at the final step the hydrogen is redelivered to an in situ-generated bond to accomplish efficient forging of C-C bond, giving water as the sole by-product.
Description:	Only IISER Mohali authors are available in the record.
URI:	https://doi.org/10.1002/adsc.202101077 (https://doi.org/10.1002/adsc.202101077) http://hdl.handle.net/123456789/4895 (http://hdl.handle.net/123456789/4895)
Appears in Collections:	Research Articles (/jspui/handle/123456789/9)

Files in This Item:

File	Description	Size	Format
Need To Add...Full Text_PDF (/jspui/bitstream/123456789/4895/1/Need%20To%20Add%e2%80%a6Full%20Text_PDF)		15.36 kB	Unknown

[View/Open \(/jspui/t](#)

Show full item record (</jspui/handle/123456789/4895?mode=full>)

 (</jspui/handle/123456789/4895/statistics>)

Items in DSpace are protected by copyright, with all rights reserved, unless otherwise indicated.